Supplementary Information

Impact of Eastern and Central Pacific El Niño on Lower Tropospheric Ozone in China

Zhongjing Jiang¹, Jing Li¹

¹ Department of Atmospheric and Oceanic Sciences, School of Physics, Peking University, Beijing, China

Correspondence: Jing Li (jing-li@pku.edu.cn)

Contents of this file

Figures S1 to S7
Tables S1

Figure S1. The sea surface temperature (SST) anomalies for four seasons in EP (2015-2016) and CP (2009-2010) El Niño years.

Figure S2. The composite sea surface temperature (SST) anomalies for four seasons in EP and CP El Niño years.

Figure S3. The simulated (GC) and satellite-observed (IASI) seasonal mean tropospheric column ozone concentration (0-6 km, unit: DU) for four seasons from September 2007 to August 2017.

Figure S4. The satellite-observed (IASI) and simulated (GC) tropospheric column ozone (0-6 km, unit: DU) anomalies for four seasons in EP (2015-2016) and CP (2009-2010) El Niño years.

Figure S5. The simulated tropospheric column ozone (0-6 km, unit: DU) anomalies driven by the composite meteorological fields for four seasons in EP and CP El Niño years.

Figure S6. The composite solar radiation (SR) anomalies for four seasons in EP and CP El Niño years.

Figure S7. The percentage changes (unit: %) of composite simulated (GC) tropospheric column ozone (0-6 km, unit: DU) anomalies for four seasons in EP and CP La Niña years. The selected EP and CP La Niña years are shown in Table S1.

type	years
EP La Niña	1984-1985, 1995-1996, 2005-2006
CP La Niña	1988-1989, 1998-1999, 2000-2001, 2008-2009, 2011-2012

Table S1. The classification results of EP and CP El Niño from 1980 to 2017, adopted from (Shi & Qian, 2018).

References

- Shi, J., & Qian, W. (2018). Asymmetry of two types of ENSO in the transition between the East Asian winter monsoon and the ensuing summer monsoon. *Climate Dynamics*, 51(9–10), 3907–3926. <https://doi.org/10.1007/s00382-018-4119-1>