Supplement of

Predicting the mineral composition of dust aerosols – Part 2: Model evaluation and identification of key processes with observations

J. P. Perlwitz et al.

Correspondence to: J. P. Perlwitz (jan.p.perlwitz@nasa.gov)

Supplement of “Predicting the mineral composition of dust aerosols. Part II: Model evaluation and identification of key processes with observations” by Perlwitz et al.

Contents

5	1 Supplementary Figures	S2
	2 Supplementary Tables	S8

List of Figures

	S1	Simulated versus observed fractions of illite, kaolinite, smectite, and sum of illite and smectite, and sum of illite, smectite, and kaolinite	S2
10	S2	Ratio between various minerals and illite	S3
	S3	Ratio between various minerals and kaolinite	S4
	S4	Ratio between various minerals and smectite	S5
	S5	Ratio between various minerals and sum of illite, kaolinite, and smectite	S6
15	S6	Ratio between various minerals and carbonates	S7

List of Tables

S1	Mineral fraction data used in evaluation	S9
----	--	----

1 Supplementary Figures

Figure S1. Scatter plot of mineral fractions of illite, kaolinite, smectite, the sum of illite and smectite, and the sum of illite, kaolinite, and smectite for all size groups together, as simulated with soil mineral fraction (SMF) method, aerosol mineral fraction (AMF) method, and in the simulation without soil reaggregation (AMF ($\gamma = 0$)) versus observations. The dashed lines mark a ratio of 2:1 and 1:2 between simulated and observed mineral fractions. The horizontal and vertical error bars show the 95% confidence intervals of the observation and simulation data, respectively, based on an assumed beta distribution using the method of moments, when sufficient data were available for the calculation.

Figure S2. Measured versus simulated ratios between various mineral phases and illite for the soil mineral fraction (SMF) method and the aerosol mineral fraction (AMF) method. The dashed lines mark a ratio of 2:1 and 1:2 between simulated and observed mineral ratios.

Figure S3. Measured versus simulated ratios between various mineral phases and kaolinite for the soil mineral fraction (SMF) method and the aerosol mineral fraction (AMF) method. The dashed lines mark a ratio of 2:1 and 1:2 between simulated and observed mineral ratios.

Figure S4. Measured versus simulated ratios between various mineral phases and smectite for the soil mineral fraction (SMF) method and the aerosol mineral fraction (AMF) method. The dashed lines mark a ratio of 2:1 and 1:2 between simulated and observed mineral ratios.

Figure S5. Measured versus simulated ratios between various mineral phases and the sum of illite, kaolinite, and smectite for the soil mineral fraction (SMF) method and the aerosol mineral fraction (AMF) method. The dashed lines mark a ratio of 2:1 and 1:2 between simulated and observed mineral ratios.

Figure S6. Measured versus simulated ratios between various mineral phases and carbonates for the soil mineral fraction (SMF) method and the aerosol mineral fraction (AMF) method. The dashed lines mark a ratio of 2:1 and 1:2 between simulated and observed mineral ratios.

2 Supplementary Tables

Table S1: Mineral fraction data used in evaluation of the model. Mean and standard deviation (StdDev) are calculated from temporal data in the reference or are provided by the reference. N_O is the number of data points as reported by the reference (e.g., the average). Bold faced minerals are used for the comparison with the ModelE minerals.

Reference: Adedokum et al. (1989)			Size Range: Bulk
Ile-Ife, Nigeria (Lat: 7°17'N, Lon: 4°20'E)			
Variable: Total Dry Deposition			
Note: Average of 12 sites			
Mineral	Mean (%)	StdDev (%)	N_O
Halloysite	1.45	na	1
Kaolinite	10.28	na	1
Mica	2.54	na	1
Microcline	9.74	na	1
Quartz	74.78	na	1
Reference: Alastuey et al. (2005)			Size Range: Bulk
Izaña, Tenerife, Spain (Lat: 28°19'N, Lon: 16°30'W)			
Variable: Concentration			
Note: Dust event			
Mineral	Mean (%)	StdDev (%)	N_O
Albite	5.00	na	1
Calcite	9.00	na	1
Clinochlore	8.00	na	1
Gypsum	3.50	na	1
Halite	0.00	na	1
Illite	15.00	na	1
Kaolinite	14.00	na	1
Mascagnite	6.00	na	1
Microcline	4.00	na	1
Palygorskite	10.00	na	1
Quartz	23.00	na	1
Reference: Alastuey et al. (2005)			Size Range: Bulk
Santa Cruz de Tenerife, Spain (Lat: 28°28'N, Lon: 16°15'W)			
Variable: Surface Concentration			
Note: Dust event			
Mineral	Mean (%)	StdDev (%)	N_O
Albite	5.00	na	1
Calcite	6.00	na	1
Clinochlore	9.00	na	1
Gypsum	10.00	na	1
Halite	2.00	na	1
Illite	13.00	na	1

Kaolinite	10.00	na	1
Muscagnite	5.00	na	1
Microcline	5.00	na	1
Palygorskite	9.00	na	1
Quartz	23.00	na	1

Reference: Al-Awadhi and AlShuaibi (2013) Size Range: Bulk

Kuwait City, Kuwait (Lat: 29°22'11"N, Lon: 47°58'42"E)

Variable: Total deposition

Note: Spatial-temporal sampling

Mineral	Mean (%)	StdDev (%)	N_O
Albite	9.38	2.51	10
Calcite	37.76	6.12	10
Dolomite	5.88	1.87	10
Illite	11.63	1.67	10
Orthoclase	5.27	4.79	10
Quartz	28.37	11.89	10

Reference: Al-Dousari and Al-Awadhi (2012) Size Range: Bulk

Ain, West U.A.E. (Lat: 24°12'27"N, Lon: 55°44'41"E)

Variable: Total deposition

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	34.00	na	1
Clay	1.00	na	1
Dolomite	19.00	na	1
Feldspar	20.00	na	1
Others	0.00	na	1
Quartz	26.00	na	1

Reference: Al-Dousari and Al-Awadhi (2012) Size Range: Bulk

Amman, Jordan (Lat: 31°56'59"N, Lon: 35°55'58"E)

Variable: Total deposition

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	52.00	na	1
Clay	0.00	na	1
Dolomite	16.00	na	1
Feldspar	4.00	na	1
Others	7.00	na	1
Quartz	21.00	na	1

Reference: Al-Dousari and Al-Awadhi (2012) Size Range: Bulk

Bubiyah, East Kuwait (Lat: 29°46'11"N, Lon: 48°15'20"E)

Variable: Total deposition

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	20.00	na	1
Clay	5.00	na	1
Dolomite	14.00	na	1

Feldspar	18.00	na	1
Others	14.00	na	1
Quartz	28.00	na	1
Reference: Al-Dousari and Al-Awadhi (2012)			Size Range: Bulk
Doha, Qatar (Lat: 25°17'12"N, Lon: 51°32'0"E)			
Variable: Total deposition			
Mineral	Mean (%)	StdDev (%)	N_O
Calcite	21.00	na	1
Clay	0.00	na	1
Dolomite	7.00	na	1
Feldspar	24.00	na	1
Others	0.00	na	1
Quartz	48.00	na	1
Reference: Al-Dousari and Al-Awadhi (2012)			Size Range: Bulk
Dubai, East U.A.E. (Lat: 24°57'N, Lon: 55°20'E)			
Variable: Total deposition			
Mineral	Mean (%)	StdDev (%)	N_O
Calcite	25.00	na	1
Clay	0.00	na	1
Dolomite	21.00	na	1
Feldspar	6.00	na	1
Others	27.00	na	1
Quartz	21.00	na	1
Reference: Al-Dousari and Al-Awadhi (2012)			Size Range: Bulk
Manama, Bahrain (Lat: 26°13'0"N, Lon: 50°35'0"E)			
Variable: Total deposition			
Mineral	Mean (%)	StdDev (%)	N_O
Calcite	25.00	na	1
Clay	3.00	na	1
Dolomite	16.00	na	1
Feldspar	10.00	na	1
Others	15.00	na	1
Quartz	32.00	na	1
Reference: Al-Dousari and Al-Awadhi (2012)			Size Range: Bulk
National Park, North Kuwait (Lat: 29°30'46"N, Lon: 47°48'27"E)			
Variable: Total deposition			
Mineral	Mean (%)	StdDev (%)	N_O
Calcite	38.00	na	1
Clay	2.00	na	1
Dolomite	7.00	na	1
Feldspar	10.00	na	1
Others	5.00	na	1
Quartz	38.00	na	1

Reference: Al-Dousari and Al-Awadhi (2012)			Size Range: Bulk
Sabiya, East Kuwait (Lat: 29°34'24"N, Lon: 48°10'17"E)			
Variable: Total deposition			
Mineral	Mean (%)	StdDev (%)	N_O
Calcite	26.00	na	1
Clay	6.00	na	1
Dolomite	11.00	na	1
Feldspar	12.00	na	1
Others	5.00	na	1
Quartz	39.00	na	1
Reference: Al-Dousari and Al-Awadhi (2012)			Size Range: Bulk
Wadi Dawasir, Saudi Arabia (Lat: 20°27'52"N, Lon: 44°47'14"E)			
Variable: Total deposition			
Mineral	Mean (%)	StdDev (%)	N_O
Calcite	13.00	na	1
Clay	1.00	na	1
Dolomite	0.00	na	1
Feldspar	24.00	na	1
Others	0.00	na	1
Quartz	62.00	na	1
Reference: Al-Dousari and Al-Awadhi (2012)			Size Range: Bulk
Warba, NE Kuwait (Lat: 29°59'38"N, Lon: 48°4'0"E)			
Variable: Total deposition			
Mineral	Mean (%)	StdDev (%)	N_O
Calcite	30.00	na	1
Clay	5.00	na	1
Dolomite	11.00	na	1
Feldspar	8.00	na	1
Others	9.00	na	1
Quartz	36.00	na	1
Reference: Al-Dousari et al. (2013)			Size Range: Bulk
Cairo, Egypt (Lat: 30°03'N, Lon: 31°14'E)			
Variable: Total deposition			
Mineral	Mean (%)	StdDev (%)	N_O
Calcite	20.00	na	1
Clay	0.00	na	1
Dolomite	14.00	na	1
Feldspar	15.00	na	1
Others	0.00	na	1
Quartz	51.00	na	1
Reference: Al-Dousari et al. (2013)			Size Range: Bulk
Cartagena, Colombia (Lat: 10°24'N, Lon: 75°30'W)			

Variable: Total deposition

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	0.00	na	1
Clay	0.00	na	1
Dolomite	0.00	na	1
Feldspar	33.00	na	1
Others	1.00	na	1
Quartz	66.00	na	1

Reference: Al-Dousari et al. (2013)

Size Range: Bulk

Zelfi, Saudi Arabia (Lat: 26°17'N, Lon: 44°48'E)

Variable: Total deposition

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	13.00	na	1
Clay	1.00	na	1
Dolomite	0.00	na	1
Feldspar	24.00	na	1
Others	0.00	na	1
Quartz	62.00	na	1

Reference: Arnold et al. (1998)

Size Range: < 2 μm

ADIOS Experiment, North of Hawaii (Lat: 26°00'N, Lon: 155°00'W)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	4.86	1.72	12
Illite	68.88	9.97	12
Kaolinite	10.31	7.06	12
Plagioclase	5.19	5.23	12
Quartz	8.05	1.73	12
Smectite	2.28	0.75	12

Reference: Arnold et al. (1998)

Size Range: 2 – 20 μm

ADIOS Experiment, North of Hawaii (Lat: 26°00'N, Lon: 155°00'W)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	4.66	1.26	12
Illite	60.20	11.76	12
Kaolinite	9.35	5.72	12
Plagioclase	9.88	6.74	12
Quartz	13.53	6.61	12
Smectite	2.37	0.77	12

Reference: Arnold et al. (1998)

Size Range: < 2 μm

Moana Wave Cruise, Northeastern Pacific (Lat: Various, Lon: Various)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	4.44	1.71	10

Illite	51.12	15.65	10
Kaolinite	5.70	4.71	10
Plagioclase	7.16	6.55	10
Quartz	5.15	2.44	10
Smectite	2.60	1.24	10

Reference: Arnold et al. (1998) Size Range: 2 – 20 μm

Moana Wave Cruise, Northeastern Pacific (Lat: Various, Lon: Various)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	4.72	2.65	10
Illite	33.37	14.22	10
Kaolinite	4.49	4.91	10
Plagioclase	17.02	11.09	10
Quartz	14.90	6.14	10
Smectite	2.03	0.99	10

Reference: Aston et al. (1973) Size Range: < 2 μm

Telamon/Hector Cruise, Off West African Coast (Lat: Various, Lon: Various)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	12.00	na	1
Illite	34.00	na	1
Kaolinite	29.00	na	1
Montmorillonite	25.00	na	1

Reference: Aston et al. (1973) Size Range: Bulk

Telamon/Hector Cruise, Off West African Coast (Lat: Various, Lon: Various)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Carbonates	1.05	0.24	10
Carbon	8.21	4.17	10
Quartz	7.18	5.21	10

Reference: Aston et al. (1973) Size Range: < 2 μm

Telamon/Hector Cruise, Northern Indian Ocean (Lat: Various, Lon: Various)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	19.33	1.15	3
Illite	65.67	5.03	3
Kaolinite	9.33	0.58	3
Montmorillonite	5.67	4.04	3

Reference: Aston et al. (1973) Size Range: Bulk

Telamon/Hector Cruise, Northern Indian Ocean (Lat: Various, Lon: Various)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Carbonates	1.00	0.00	6

Carbon	16.67	14.26	6
Quartz	5.33	5.00	6
Reference: Aston et al. (1973)			Size Range: Bulk
S. S. Telamon Cruise, Sea of China (Lat: Various, Lon: Various)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Carbonates	2.00	0.00	3
Carbon	16.60	5.40	3
Quartz	3.90	3.47	3
Reference: Aston et al. (1973)			Size Range: < 2 μm
Telamon/Hector Cruise, Southeastern Atlantic (Lat: Various, Lon: Various)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	13.33	2.08	3
Illite	41.33	18.50	3
Kaolinite	26.33	14.57	3
Montmorillonite	19.00	2.65	3
Reference: Aston et al. (1973)			Size Range: Bulk
Telamon/Hector Cruise, Southeastern Atlantic (Lat: Various, Lon: Various)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Carbonates	1.00	0.00	6
Carbon	12.64	9.71	6
Quartz	3.18	2.18	6
Reference: Aston et al. (1973)			Size Range: < 2 μm
Telamon/Hector Cruise, Southeast Africa Coast (Lat: Various, Lon: Various)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	29.00	22.00	2
Illite	53.00	22.00	2
Kaolinite	11.33	5.33	2
Montmorillonite	6.67	5.33	2
Reference: Aston et al. (1973)			Size Range: Bulk
Telamon/Hector Cruise, Southeast Africa Coast (Lat: Various, Lon: Various)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Carbonates	1.00	0.00	5
Carbon	21.22	17.39	5
Quartz	4.78	5.72	5
Reference: Avila et al. (1997)			Size Range: Bulk
Montseny Mountains, Spain (Lat: 41°46'N, Lon: 2°21'W)			
Variable: Bulk Deposition			
Note: Red Rain events			

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	8.68	7.37	13
Dolomite	4.54	2.56	13
Feldspar	3.38	2.43	13
Illite	38.96	11.17	13
Kaolinite	6.18	3.87	13
Palygorskite	8.85	2.38	13
Quartz	18.73	3.57	13
Smectite	11.00	9.00	12

Reference: Awadh (2012)

Size Range: Bulk

Baghdad, Iraq (Lat: 33°20'N, Lon: 44°26'E)

Variable: Total Dry Deposition

Note: No clays in samples; dust storms

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	35.56	12.22	5
Dolomite	0.80	0.30	5
Feldspar	4.98	2.26	5
Gypsum	4.88	2.39	5
Heavy Minerals	3.20	0.88	5
Quartz	48.96	8.75	5

Reference: Chester and Johnson (1971a)

Size Range: < 2 μm

M. V. Surveyor Cruise, Off Moroccan Coast (Lat: Various, Lon: Various)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	13.20	1.39	2
Illite	42.40	2.77	2
Kaolinite	25.20	2.08	2
Montmorillonite	17.20	5.54	2

Reference: Chester and Johnson (1971b)

Size Range: < 2 μm

R. R. S. Discovery Cruise, Off West African Coast (Lat: Various, Lon: Various)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	15.67	3.06	3
Illite	53.67	6.66	3
Kaolinite	21.67	2.31	3
Montmorillonite	9.00	6.24	3

Reference: Chester et al. (1971)

Size Range: < 2 μm

RMMV Good Hope Castle Cruise, Eastern Atlantic (Lat: Various, Lon: Various)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	18.00	7.68	5
Illite	34.40	10.41	5
Kaolinite	32.60	17.26	5

Montmorillonite	15.00	17.52	5
Reference: Chester et al. (1972) Size Range: < 2 μm			
M. V. Elpenor Cruise, Southeastern Atlantic (Lat: Various, Lon: Various)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	15.24	5.91	13
Illite	41.28	14.00	13
Kaolinite	30.98	12.01	13
Montmorillonite	12.29	8.59	10
Reference: Chester et al. (1977) Size Range: < 2 μm			
R. R. S. Shackleton Cruise, Eastern Mediterranean (Lat: Various, Lon: Various)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	6.81	5.05	16
Illite	44.50	10.17	16
Kaolinite	20.88	6.50	16
Smectite	27.81	12.20	16
Reference: Chester et al. (1977) Size Range: Bulk			
R. R. S. Shackleton Cruise, Eastern Mediterranean (Lat: Various, Lon: Various)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Carbonates	6.16	4.05	16
Quartz	9.06	1.69	16
Reference: Chester et al. (1977) Size Range: < 2 μm			
R. V. Chain Cruise, Eastern Mediterranean (Lat: Various, Lon: Various)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	6.40	3.91	10
Illite	39.70	6.63	10
Kaolinite	50.40	3.89	10
Smectite	3.50	0.62	10
Reference: Chester et al. (1977) Size Range: Bulk			
R. V. Chain Cruise, Eastern Mediterranean (Lat: Various, Lon: Various)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Carbonates	25.80	11.62	10
Quartz	20.00	2.31	10
Reference: Chester et al. (1984) Size Range: < 2 μm			
R. R. S. Shackleton Cruise, Tyrrhenian Sea (Lat: Various, Lon: Various)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	12.50	6.55	8
Illite	61.38	12.00	8

Kaolinite	25.25	15.28	8
Montmorillonite	0.88	1.25	8

Reference: Delany et al. (1967) Size Range: < 2 μm

Kitridge Point, Barbados, Caribbean (Lat: 13°10'N, Lon: 59°25'W)

Variable: Surface Concentration

Note: PM2: On a CaCO₃ free base

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	7.17	4.77	10
Illite	35.95	7.63	10
Kaolinite	24.64	16.13	10
Montmorillonite	18.85	3.94	10

Reference: Díaz-Hernández et al. (2011) Size Range: < 20 μm

Granada Depression, Spain (Lat: 37°10'N, Lon: 3°31'W)

Variable: Bulk Deposition

Note: Illite and Muscovite together

Mineral	Mean (%)	StdDev (%)	N_O
Amorphous Material	3.37	0.62	43
Bassanite	2.39	0.71	43
Calcite	12.46	5.41	43
Chlorite	3.73	1.15	43
Dolomite	13.53	6.15	43
Feldspar	0.81	0.32	43
Graphite	1.04	0.42	43
Gypsum	2.12	0.99	43
Halite	4.43	1.59	43
Illite	7.44	2.63	43
Kaolinite	8.00	2.83	43
Paragonite	2.58	0.85	43
Quartz	27.73	8.50	43
Smectite	10.35	3.36	43

Reference: Enete et al. (2012) Size Range: < 2 μm

Enugu, Nigeria (Lat: Various, Lon: Various)

Variable: Total deposition

Note: Average of maximum and minimum

Mineral	Mean (%)	StdDev (%)	N_O
Al-Chloride	11.00	na	1
Ca-Na-feldspar	1.50	na	1
Fe-Al Oxyhydroxide	3.50	na	1
Fe-illite	13.00	na	1
Illite	13.00	na	1
Kaolinite	24.50	na	1
K-feldspar	1.50	na	1
Quartz	7.00	na	1

Vermiculite	7.00	na	1
Reference: Enete et al. (2012)			Size Range: 2 – 50 μm
Enugu, Nigeria (Lat: Various, Lon: Various)			
Variable: Total deposition			
Note: Average of maximum and minimum			
Mineral	Mean (%)	StdDev (%)	N_O
Ca-Na-feldspar	2.00	na	1
Kaolinite	15.00	na	1
K-feldspar	21.00	na	1
Mica	15.00	na	1
Quartz	58.00	na	1
Reference: Engelbrecht et al. (2009)			Size Range: < 10 μm
Bagram, Afghanistan (Lat: 34°57'N, Lon: 69°16'E)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	13.20	na	1
Calcite	8.40	na	1
Carbon	0.50	na	1
Ca-Si	12.70	na	1
Clay	31.80	na	1
Dolomite	8.50	na	1
Iron oxides	1.30	na	1
K-feldspar	11.00	na	1
Lead	0.40	na	1
Misc	2.10	na	1
Na-feldspar	1.70	na	1
Quartz	4.60	na	1
Salts	1.60	na	1
Si-Mg	2.10	na	1
Sulfates	0.10	na	1
Reference: Engelbrecht et al. (2009)			Size Range: < 10 μm
Khowst, Afghanistan (Lat: 33°22'N, Lon: 69°58'E)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	5.10	na	1
Calcite	5.50	na	1
Carbon	1.66	na	1
Ca-Si	6.80	na	1
Clay	50.90	na	1
Dolomite	1.30	na	1
Iron oxides	0.40	na	1
K-feldspar	11.00	na	1
Lead	0.02	na	1
Misc	0.02	na	1

Na-feldspar	2.50	na	1
Quartz	5.10	na	1
Salts	1.30	na	1
Si-Mg	8.00	na	1
Sulfates	0.40	na	1

Reference: Engelbrecht et al. (2009) Size Range: < 10 μm

Djibouti (Lat: 11°33'N, Lon: 43°09'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	12.70	na	1
Calcite	2.50	na	1
Carbon	0.80	na	1
Ca-Si	10.20	na	1
Clay	49.20	na	1
Dolomite	0.90	na	1
Iron oxides	2.60	na	1
K-feldspar	5.50	na	1
Lead	0.90	na	1
Misc	0.80	na	1
Na-feldspar	1.30	na	1
Quartz	1.70	na	1
Salts	7.60	na	1
Si-Mg	1.60	na	1
Sulfates	1.70	na	1

Reference: Engelbrecht et al. (2009) Size Range: < 10 μm

Tallil, Iraq (Lat: 30°56'N, Lon: 46°5'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	6.80	na	1
Calcite	7.60	na	1
Carbon	0.10	na	1
Ca-Si	14.80	na	1
Clay	55.50	na	1
Dolomite	2.50	na	1
Iron oxides	0.70	na	1
K-feldspar	4.20	na	1
Lead	0.60	na	1
Misc	0.60	na	1
Na-feldspar	1.30	na	1
Quartz	1.70	na	1
Salts	1.70	na	1
Si-Mg	0.60	na	1
Sulfates	1.30	na	1

Reference: Engelbrecht et al. (2009) Size Range: < 10 μm

Al Asad, Iraq (Lat: 33°48'N, Lon: 42°26'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	5.90	na	1
Calcite	13.10	na	1
Carbon	0.40	na	1
Ca-Si	14.00	na	1
Clay	31.80	na	1
Dolomite	21.20	na	1
Iron oxides	0.90	na	1
K-feldspar	3.20	na	1
Lead	0.80	na	1
Misc	1.30	na	1
Na-feldspar	1.30	na	1
Quartz	1.70	na	1
Salts	1.30	na	1
Si-Mg	0.40	na	1
Sulfates	2.50	na	1

Reference: Engelbrecht et al. (2009)

Size Range: < 10 μm

Balad, Iraq (Lat: 33°56'N, Lon: 44°22'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	10.60	na	1
Calcite	8.10	na	1
Carbon	0.80	na	1
Ca-Si	15.70	na	1
Clay	45.30	na	1
Dolomite	1.70	na	1
Iron oxides	1.30	na	1
K-feldspar	5.50	na	1
Lead	1.29	na	1
Misc	0.01	na	1
Na-feldspar	1.30	na	1
Quartz	3.80	na	1
Salts	1.70	na	1
Si-Mg	2.10	na	1
Sulfates	0.80	na	1

Reference: Engelbrecht et al. (2009)

Size Range: < 10 μm

Tikrit, Iraq (Lat: 34°41'N, Lon: 43°33'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	13.50	na	1
Calcite	9.40	na	1
Carbon	0.80	na	1

Ca-Si	23.40	na	1
Clay	35.60	na	1
Dolomite	0.40	na	1
Iron oxides	0.40	na	1
K-feldspar	4.20	na	1
Lead	1.20	na	1
Misc	0.10	na	1
Na-feldspar	0.90	na	1
Quartz	3.40	na	1
Salts	0.10	na	1
Si-Mg	1.20	na	1
Sulfates	5.40	na	1

Reference: Engelbrecht et al. (2009)

Size Range: < 10 μm

Taji, Iraq (Lat: 33°32'N, Lon: 44°17'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	9.70	na	1
Calcite	6.80	na	1
Carbon	0.60	na	1
Ca-Si	17.00	na	1
Clay	44.90	na	1
Dolomite	3.00	na	1
Iron oxides	1.20	na	1
K-feldspar	4.20	na	1
Lead	0.70	na	1
Misc	0.80	na	1
Na-feldspar	1.30	na	1
Quartz	2.60	na	1
Salts	3.40	na	1
Si-Mg	1.30	na	1
Sulfates	2.50	na	1

Reference: Engelbrecht et al. (2009)

Size Range: < 10 μm

Baghdad, Iraq (Lat: 33°18'N, Lon: 44°15'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	12.30	na	1
Calcite	8.50	na	1
Carbon	1.29	na	1
Ca-Si	17.30	na	1
Clay	39.90	na	1
Dolomite	2.90	na	1
Iron oxides	1.30	na	1
K-feldspar	3.80	na	1
Lead	0.01	na	1

Misc	0.80	na	1
Na-feldspar	1.20	na	1
Quartz	2.60	na	1
Salts	3.00	na	1
Si-Mg	1.70	na	1
Sulfates	3.40	na	1

Reference: Engelbrecht et al. (2009)

Size Range: < 10 μm

Central Kuwait (Lat: 29°21'N, Lon: 47°31'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	7.60	na	1
Calcite	6.80	na	1
Carbon	0.20	na	1
Ca-Si	13.60	na	1
Clay	55.50	na	1
Dolomite	2.60	na	1
Iron oxides	1.20	na	1
K-feldspar	3.80	na	1
Lead	0.30	na	1
Misc	0.80	na	1
Na-feldspar	1.30	na	1
Quartz	2.50	na	1
Salts	1.30	na	1
Si-Mg	1.30	na	1
Sulfates	1.20	na	1

Reference: Engelbrecht et al. (2009)

Size Range: < 10 μm

Southern Kuwait (Lat: 28°53'N, Lon: 48°10'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	8.90	na	1
Calcite	9.30	na	1
Carbon	0.40	na	1
Ca-Si	12.30	na	1
Clay	40.70	na	1
Dolomite	5.50	na	1
Iron oxides	1.30	na	1
K-feldspar	5.10	na	1
Lead	0.40	na	1
Misc	1.70	na	1
Na-feldspar	1.30	na	1
Quartz	3.40	na	1
Salts	1.70	na	1
Si-Mg	6.70	na	1
Sulfates	1.30	na	1

Reference: Engelbrecht et al. (2009) Size Range: < 10 μm

Northern Kuwait (Lat: 29°42'N, Lon: 47°25'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	7.20	na	1
Calcite	9.30	na	1
Carbon	0.90	na	1
Ca-Si	18.20	na	1
Clay	44.10	na	1
Dolomite	2.10	na	1
Iron oxides	1.30	na	1
K-feldspar	3.90	na	1
Lead	0.70	na	1
Misc	0.10	na	1
Na-feldspar	1.20	na	1
Quartz	2.50	na	1
Salts	1.30	na	1
Si-Mg	1.70	na	1
Sulfates	5.50	na	1

Reference: Engelbrecht et al. (2009) Size Range: < 10 μm

Coastal Kuwait (Lat: 29°3'N, Lon: 48°9'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	7.60	na	1
Calcite	7.20	na	1
Carbon	0.60	na	1
Ca-Si	12.30	na	1
Clay	44.90	na	1
Dolomite	6.40	na	1
Iron oxides	1.70	na	1
K-feldspar	3.80	na	1
Lead	0.70	na	1
Misc	0.80	na	1
Na-feldspar	1.30	na	1
Quartz	3.00	na	1
Salts	3.40	na	1
Si-Mg	3.80	na	1
Sulfates	2.50	na	1

Reference: Engelbrecht et al. (2009) Size Range: < 10 μm

Qatar (Lat: 25°7'N, Lon: 51°19'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	5.50	na	1

Calcite	11.00	na	1
Carbon	0.40	na	1
Ca-Si	11.10	na	1
Clay	45.70	na	1
Dolomite	6.80	na	1
Iron oxides	1.70	na	1
K-feldspar	4.20	na	1
Lead	0.50	na	1
Misc	0.80	na	1
Na-feldspar	0.90	na	1
Quartz	1.70	na	1
Salts	1.30	na	1
Si-Mg	2.50	na	1
Sulfates	5.90	na	1

Reference: Engelbrecht et al. (2009)

Size Range: < 10 μm

United Arab Emirates (Lat: 24°15'N, Lon: 54°33'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Ca-feldspar	4.20	na	1
Calcite	16.60	na	1
Carbon	0.90	na	1
Ca-Si	8.90	na	1
Clay	23.30	na	1
Dolomite	15.20	na	1
Iron oxides	1.20	na	1
K-feldspar	2.50	na	1
Lead	0.80	na	1
Misc	1.70	na	1
Na-feldspar	0.90	na	1
Quartz	1.70	na	1
Salts	3.40	na	1
Si-Mg	15.30	na	1
Sulfates	3.40	na	1

Reference: Engelbrecht et al. (2014)

Size Range: < 2.5 μm

Las Palmas de Gran Canaria, Spain (Lat: 28°04'26.28"N, Lon: 15°27'08.57"W)

Variable: Surface concentration

Mineral	Mean (%)	StdDev (%)	N_O
Ammonium sulfate	13.10	10.20	16
Calcite	0.59	1.02	16
Clay	24.33	15.77	16
Elemental carbon	4.46	4.16	16
Iron oxides	4.56	3.20	16
Gypsum	6.54	3.69	16
Halite	3.70	2.51	16

Organic carbon	14.95	8.36	16
Other evaporites	17.92	13.12	16
Quartz	9.84	9.60	16

Reference: Falkovich et al. (2001) Size Range: Bulk

Tel-Aviv, Israel (Lat: 32°5'N, Lon: 34°48'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	29.00	na	1
Dolomite	26.00	na	1
Feldspar	8.00	na	1
Gypsum	5.00	na	1
Quartz	29.00	na	1

Reference: Ferguson et al. (1970) Size Range: < 2 μm

USCGC Acushnet Cruise, Northeastern Pacific (Lat: Various, Lon: Various)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	19.25	6.91	10
Illite	49.05	4.81	10
Kaolinite	14.35	3.08	10
Montmorillonite	15.45	7.03	10

Reference: Fiol et al. (2005) Size Range: Bulk

Palma de Mallorca, Spain (Lat: 39°38'N, Lon: 2°39'E)

Variable: Bulk Deposition

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	26.10	2.54	9
Dolomite	9.40	6.94	9
Illite	1.30	1.89	9
Kaolinite	2.50	2.10	9
K-feldspar	1.90	2.27	9
Palygorskite	1.00	1.28	9
Plagioclase	4.00	2.99	9
Quartz	54.10	13.69	9

Reference: Formenti et al. (2008) Size Range: < 40 μm

Banizoumbou, Niger (Lat: 13°30'N, Lon: 2°36'E)

Variable: Surface Concentration

Note: Percent of diffractive surface

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	2.59	1.84	22
Dolomite	1.41	0.75	22
Illite	9.98	1.65	22
Kaolinite	46.34	5.10	22
Other feldspar	2.32	1.03	22
Plagioclase	3.98	0.88	22

Quartz	33.39	4.13	22
Reference: Game (1964)			Size Range: Bulk
S. S. Duncan, Eastern Atlantic (Lat: 25°04'N, Lon: 20°44'W)			
Variable: Total Dry Deposition			
Note: Only non-aggregated			
Mineral	Mean (%)	StdDev (%)	N_O
Aggregates	32.00	na	1
Amphibole	2.00	na	1
Biotite	2.00	na	1
Carbonates	20.00	na	1
Feldspar	4.00	na	1
Iron oxides	3.50	na	1
Organics	17.50	na	1
Others	1.00	na	1
Quartz	18.00	na	1
Reference: Ganor (1991)			Size Range: < 10 μm
Jerusalem, Israel (Lat: 31°47'N, Lon: 35°13'E)			
Variable: Total Dry Deposition			
Note: dust storms			
Mineral	Mean (%)	StdDev (%)	N_O
Illite	28.64	na	28
Kaolinite	26.82	na	28
Montmorillonite and mixed layer	40.96	na	28
Palygorskite	3.57	na	28
Reference: Ganor et al. (2000)			Size Range: < 2 μm
Lake Kinneret, Israel (Lat: Various, Lon: Various)			
Variable: Total Deposition			
Note: Mean of 16 locations			
Mineral	Mean (%)	StdDev (%)	N_O
Illite	16.00	0.00	1
Kaolinite	34.00	0.00	1
Palygorskite	42.00	0.00	1
Smectite	8.00	0.00	1
Reference: Ganor et al. (2000)			Size Range: > 2 μm
Lake Kinneret, Israel (Lat: Various, Lon: Various)			
Variable: Total Deposition			
Note: Average Mean, StdDev, and N_O of 16 locations			
Mineral	Mean (%)	StdDev (%)	N_O
Calcite	21.09	13.98	42
Dolomite	16.67	8.26	42
Feldspar	32.89	14.05	42
Quartz	27.61	12.93	42
Reference: Gaudichet et al. (1989)			Size Range: Bulk

Amsterdam Island, TAAF (Lat: 37°47'S, Lon: 77°31'E)

Variable: Surface Concentration

Note: From Number Fractions Assuming Model Densities

Mineral	Mean (%)	StdDev (%)	N_O
Carbonates	3.43	2.27	3
Feldspar	4.78	5.10	3
Illite	30.65	9.34	3
Kaolinite	34.71	20.81	3
Quartz	23.48	16.90	3
Smectite	2.96	7.49	3

Reference: Gaudichet et al. (1992)

Size Range: < 2 μm

South Pole, Antarctica (Lat: 90°S, Lon: 0°)

Variable: Total Dry Deposition

Note: from snow

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	6.00	na	1
Illite	65.00	na	1
Kaolinite	12.00	na	1
Smectite	16.00	na	1
Vermiculite	1.00	na	1

Reference: Gaudichet et al. (1992)

Size Range: < 2 μm

Vostok, Antarctica (Lat: 78°27.87'S, Lon: 106°50.24E)

Variable: Total Dry Deposition

Note: from snow

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	6.00	na	1
Illite	65.00	na	1
Kaolinite	12.00	na	1
Smectite	16.00	na	1
Vermiculite	1.00	na	1

Reference: Glaccum and Prospero (1980)

Size Range: Bulk

Barbados (Lat: 13°10'N, Lon: 59°25'W)

Variable: Surface Concentration

Note: Assumed conversion Calcite to Gypsum

Mineral	Mean (%)	StdDev (%)	N_O
Carbonates	3.68	0.69	3
Chlorite	4.15	0.28	3
Kaolinite	8.39	0.75	3
Mica	64.28	0.97	3
Microcline	1.53	0.06	3
Plagioclase	4.07	0.54	3
Quartz	13.89	1.29	3

Reference: Glaccum and Prospero (1980)

Size Range: Bulk

Miami, Florida (Lat: 25°47'N, Lon: 80°13'W)

Variable: Surface Concentration

Note: Assumed conversion Calcite to Gypsum

Mineral	Mean (%)	StdDev (%)	N_O
Carbonates	6.97	0.76	3
Chlorite	4.30	0.36	3
Kaolinite	7.03	1.65	3
Mica	62.10	1.85	3
Microcline	1.10	0.20	3
Plagioclase	4.40	0.46	3
Quartz	14.27	0.95	3

Reference: Glaccum and Prospero (1980)

Size Range: Bulk

Sal Island, Cape Verde (Lat: 16°45'N, Lon: 22°57'W)

Variable: Surface Concentration

Note: Assumed conversion Calcite to Gypsum

Mineral	Mean (%)	StdDev (%)	N_O
Carbonates	8.11	1.21	3
Chlorite	4.35	0.21	3
Kaolinite	6.56	0.06	3
Mica	53.50	1.32	3
Microcline	2.17	0.24	3
Plagioclase	5.35	0.21	3
Quartz	19.53	1.15	3

Reference: Goldberg and Griffin (1970)

Size Range: < 2 μm

CIRCE Cruise, Bay of Bengal (Lat: Various, Lon: Various)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	27.05	3.92	23
Illite	48.45	4.61	23
Kaolinite	10.88	3.10	23
Montmorillonite	14.00	3.93	23

Reference: Goldberg and Griffin (1970)

Size Range: < 2 μm

Waltair, Eastern India (Lat: 17°45'N, Lon: 83°25'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	11.16	3.72	16
Illite	36.42	13.11	16
Kaolinite	27.29	3.63	16
Montmorillonite	25.16	14.81	16

Reference: Jeong (2008)

Size Range: < 10 μm

Seoul, South Korea (Lat: 37°30'N, Lon: 126°55'E)

Variable: Surface Concentration

Note: Mixed Layer Illite/Smectite added to Illite and Smectite

Mineral	Mean (%)	StdDev (%)	N_O
Carbonates	8.62	5.18	8
Chlorite	2.38	0.52	8
Kaolinite	1.12	0.35	8
K-feldspar	8.25	1.04	8
Interstratified illite-smectite	21.62	5.68	8
Plagioclase	10.62	2.07	8
Illite	19.50	5.04	8
Smectite	0.88	0.35	8
Quartz	27.88	5.44	8

Reference: Jeong and Achterberg (2014)

Size Range: < 60 μm

Andong, South Korea (Lat: 36°32'34"N, Lon: 128°47'56"E)

Variable: Surface concentration

Note: Dust storms

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	1.33	1.15	3
Calcite	5.33	0.58	3
Chlorite	5.67	2.31	3
Gypsum	8.33	4.04	3
Kaolinite	3.00	1.73	3
K-feldspar	1.00	1.00	3
Interstratified illite-smectite	50.67	9.02	3
Plagioclase	11.00	1.00	3
Quartz	14.00	1.00	3

Reference: Jeong and Achterberg (2014)

Size Range: < 60 μm

Deokjeok Island, South Korea (Lat: 37°13'59"N, Lon: 126°08'57"E)

Variable: Surface concentration

Note: Dust storms

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	1.00	na	1
Calcite	2.00	na	1
Chlorite	6.00	na	1
Gypsum	2.00	na	1
Kaolinite	3.00	na	1
K-feldspar	6.00	na	1
Interstratified illite-smectite	42.00	na	1
Plagioclase	15.00	na	1
Quartz	23.00	na	1

Reference: Jeong and Achterberg (2014)

Size Range: < 60 μm

Sao Vicente, Cap Verde (Lat: 16°51'50"N, Lon: 24°52'03"W)

Variable: Surface concentration

Note: Dust storms

Mineral	Mean (%)	StdDev (%)	N_O
---------	----------	------------	-------

Amphibole	0.00	0.00	2
Calcite	2.38	0.68	2
Chlorite	3.00	0.00	2
Gypsum	4.75	1.37	2
Kaolinite	6.50	2.74	2
K-feldspar	1.00	0.00	2
Interstratified illite-smectite	72.12	2.05	2
Plagioclase	2.00	0.00	2
Quartz	8.88	2.05	2

Reference: Jeong et al. (2014)

Size Range: < 60 μm

Andong, South Korea (Lat: 36°32'34"N, Lon: 128°47'56"E)

Variable: Surface concentration

Note: Dust storms

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	0.60	0.17	2
Apatite	0.24	0.17	2
Barite	0.08	0.12	2
Biotite	4.06	1.81	2
Calcite	5.84	0.07	2
Chlorite	5.63	0.94	2
Dolomite	0.59	0.53	2
Fe-Ti oxides	0.36	0.25	2
Gypsum	0.78	0.25	2
Illite, smectite, mixed layers	47.02	0.82	2
Kaolinite	1.27	0.05	2
K-feldspar	3.54	0.89	2
Muscovite	0.64	0.45	2
Plagioclase	10.33	0.35	2
Pyrite	0.18	0.26	2
Pyroxene	0.00	0.00	2
Quartz	16.76	0.07	2
Talc	0.10	0.14	2
Ti oxides	0.24	0.11	2
Titanite	0.00	0.00	2
Zoisite	0.36	0.17	2

Reference: Jeong et al. (2014)

Size Range: 10 – 20 μm

Deokjeok Island, South Korea (Lat: 37°13'59"N, Lon: 126°08'57"E)

Variable: Surface concentration

Note: Dust storms

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	0.84	na	1
Apatite	0.12	na	1
Biotite	4.97	na	1

Calcite	7.17	na	1
Chlorite	3.85	na	1
Dolomite	0.85	na	1
Fe-Ti oxides	0.35	na	1
Gypsum	0.00	na	1
Illite, smectite, mixed layers	36.35	na	1
Kaolinite	0.92	na	1
K-feldspar	6.95	na	1
Muscovite	1.37	na	1
Plagioclase	13.84	na	1
Pyrite	0.00	na	1
Pyroxene	0.00	na	1
Quartz	20.59	na	1
Talc	0.00	na	1
Ti oxides	0.00	na	1
Titanite	0.00	na	1
Zoisite	0.24	na	1

Reference: Jeong et al. (2014)

Size Range: 20 – 60 μm

Deokjeok Island, South Korea (Lat: 37°13'59"N, Lon: 126°08'57"E)

Variable: Surface concentration

Note: Dust storms

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	0.97	na	1
Apatite	0.00	na	1
Biotite	7.61	na	1
Calcite	5.22	na	1
Chlorite	1.89	na	1
Dolomite	0.00	na	1
Fe-Ti oxides	0.00	na	1
Gypsum	0.00	na	1
Illite, smectite, mixed layers	32.62	na	1
Kaolinite	5.45	na	1
K-feldspar	2.83	na	1
Muscovite	5.45	na	1
Plagioclase	7.92	na	1
Pyrite	0.00	na	1
Pyroxene	0.67	na	1
Quartz	28.56	na	1
Talc	0.00	na	1
Ti oxides	0.00	na	1
Titanite	0.80	na	1
Zoisite	0.00	na	1

Reference: Jeong et al. (2014)

Size Range: < 5 μm

Deokjeok Island, South Korea (Lat: 37°13'59"N, Lon: 126°08'57"E)

Variable: Surface concentration

Note: Dust storms

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	0.95	na	1
Apatite	0.24	na	1
Biotite	3.21	na	1
Calcite	6.51	na	1
Chlorite	4.03	na	1
Dolomite	1.79	na	1
Fe-Ti oxides	0.70	na	1
Gypsum	1.02	na	1
Illite, smectite, mixed layers	41.90	na	1
Kaolinite	2.21	na	1
K-feldspar	4.88	na	1
Muscovite	0.73	na	1
Plagioclase	9.04	na	1
Pyrite	0.36	na	1
Pyroxene	0.00	na	1
Quartz	17.35	na	1
Talc	0.10	na	1
Ti oxides	0.47	na	1
Titanite	0.39	na	1
Zoisite	0.24	na	1

Reference: Jeong et al. (2014)

Size Range: 5 – 10 μm

Deokjeok Island, South Korea (Lat: 37°13'59"N, Lon: 126°08'57"E)

Variable: Surface concentration

Note: Dust storms

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	1.08	na	1
Apatite	0.12	na	1
Biotite	3.93	na	1
Calcite	6.16	na	1
Chlorite	4.62	na	1
Dolomite	0.74	na	1
Fe-Ti oxides	0.71	na	1
Gypsum	0.00	na	1
Illite, smectite, mixed layers	40.74	na	1
Kaolinite	1.31	na	1
K-feldspar	3.86	na	1
Muscovite	2.64	na	1
Plagioclase	11.43	na	1
Pyrite	0.00	na	1
Pyroxene	0.13	na	1
Quartz	20.48	na	1

Talc	0.00	na	1
Ti oxides	0.16	na	1
Titanite	0.13	na	1
Zoisite	0.60	na	1

Reference: Jeong et al. (2014)

Size Range: < 60 μm

Deokjeok Island, South Korea (Lat: 37°13'59"N, Lon: 126°08'57"E)

Variable: Surface concentration

Note: Dust storms

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	0.97	na	1
Apatite	0.00	na	1
Biotite	6.29	na	1
Calcite	6.01	na	1
Chlorite	2.88	na	1
Dolomite	0.43	na	1
Fe-Ti oxides	0.18	na	1
Gypsum	0.00	na	1
Illite, smectite, mixed layers	34.89	na	1
Kaolinite	3.49	na	1
K-feldspar	4.28	na	1
Muscovite	3.72	na	1
Plagioclase	10.21	na	1
Pyrite	0.00	na	1
Pyroxene	0.40	na	1
Quartz	24.94	na	1
Talc	0.00	na	1
Ti oxides	0.00	na	1
Titanite	0.40	na	1
Zoisite	0.12	na	1

Reference: Johnson (1976)

Size Range: < 2 μm

BM,1969,0,87, Barbados (Lat: 13°10'N, Lon: 59°25'W)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Illite	35.00	na	1
Kaolinite+chlorite	41.00	na	1
Montmorillonite	24.00	na	1

Reference: Johnson (1976)

Size Range: Bulk

BM,1969,0,87, Barbados (Lat: 13°10'N, Lon: 59°25'W)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Clay	62.00	na	1
K-feldspar	1.50	na	1
Plagioclase	3.40	na	1

Quartz	11.00	na	1
Reference: Johnson (1976)			
BM,1973,0,288, Eastern Atlantic (Lat: 12°35'N, Lon: 17°35'W)			
Variable: Total Dry Deposition			
Mineral	Mean (%)	StdDev (%)	N_O
Illite	37.00	na	1
Kaolinite+chlorite	29.00	na	1
Montmorillonite	34.00	na	1
Reference: Johnson (1976)			
BM,1973,0,288, Eastern Atlantic (Lat: 12°35'N, Lon: 17°35'W)			
Variable: Total Dry Deposition			
Mineral	Mean (%)	StdDev (%)	N_O
Clay	27.00	na	1
K-feldspar	5.50	na	1
Plagioclase	8.20	na	1
Quartz	32.30	na	1
Reference: Johnson (1976)			
BM,M7398, Eastern Atlantic (Lat: 21°00'N, Lon: 21°00'W)			
Variable: Total Dry Deposition			
Mineral	Mean (%)	StdDev (%)	N_O
Illite	49.00	na	1
Kaolinite+chlorite	21.00	na	1
Montmorillonite	30.00	na	1
Reference: Johnson (1976)			
BM,M7398, Eastern Atlantic (Lat: 21°00'N, Lon: 21°00'W)			
Variable: Total Dry Deposition			
Mineral	Mean (%)	StdDev (%)	N_O
Clay	40.00	na	1
K-feldspar	1.90	na	1
Plagioclase	9.50	na	1
Quartz	17.80	na	1
Reference: Johnson (1976)			
E3, Central Atlantic (Lat: 11°37'S, Lon: 16°55'W)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Illite	55.00	na	1
Kaolinite+chlorite	34.00	na	1
Montmorillonite	13.00	na	1
Reference: Johnson (1976)			
E3, Central Atlantic (Lat: 11°37'S, Lon: 16°55'W)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Clay	48.00	na	1

K-feldspar	1.10	na	1
Plagioclase	5.50	na	1
Quartz	15.00	na	1

Reference: Johnson (1976) Size Range: < 2 μm

EG25, Eastern Atlantic (Lat: 14°22'N, Lon: 26°24'W)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Illite	56.00	na	1
Kaolinite+chlorite	26.00	na	1
Montmorillonite	18.00	na	1

Reference: Johnson (1976) Size Range: Bulk

EG25, Eastern Atlantic (Lat: 14°22'N, Lon: 26°24'W)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Clay	44.00	na	1
K-feldspar	1.40	na	1
Plagioclase	4.70	na	1
Quartz	10.00	na	1

Reference: Kandler et al. (2007) Size Range: 10 – 20 μm

Izaña, Tenerife, Spain (Lat: 28°19'N, Lon: 16°30'W)

Variable: Concentration

Note: From Volume Fractions Times Assumed Densities

Mineral	Mean (%)	StdDev (%)	N_O
Albite	14.56	na	1
Bloedite	0.77	na	1
Calcite	5.64	na	1
Carbonaceous	0.00	na	1
Chloritoid	23.71	na	1
Dolomite	0.99	na	1
Glauberite	0.00	na	1
Gypsum	0.80	na	1
Halite	0.75	na	1
Hematite	0.00	na	1
Illite	43.59	na	1
Ilmenite	0.00	na	1
Quartz	9.20	na	1
Rutile	0.00	na	1

Reference: Kandler et al. (2007) Size Range: 1 – 2.5 μm

Izaña, Tenerife, Spain (Lat: 28°19'N, Lon: 16°30'W)

Variable: Concentration

Note: From Volume Fractions Times Assumed Densities

Mineral	Mean (%)	StdDev (%)	N_O
Albite	13.17	na	1

Bloedite	2.40	na	1
Calcite	4.86	na	1
Carbonaceous	1.15	na	1
Chloritoid	23.22	na	1
Dolomite	1.02	na	1
Glauberite	2.01	na	1
Gypsum	0.83	na	1
Halite	0.78	na	1
Hematite	1.89	na	1
Illite	41.07	na	1
Ilmenite	0.00	na	1
Quartz	7.61	na	1
Rutile	0.00	na	1

Reference: Kandler et al. (2007)

Size Range: < 20 μm

Izaña, Tenerife, Spain (Lat: 28°19'N, Lon: 16°30'W)

Variable: Concentration

Note: From Volume Fractions Times Assumed Densities

Mineral	Mean (%)	StdDev (%)	N_O
Albite	11.90	na	1
Bloedite	7.58	na	1
Calcite	3.07	na	1
Carbonaceous	2.72	na	1
Chloritoid	20.40	na	1
Dolomite	1.08	na	1
Glauberite	5.29	na	1
Gypsum	0.87	na	1
Halite	0.00	na	0
Hematite	1.99	na	1
Illite	39.07	na	1
Ilmenite	0.00	na	0
Quartz	6.02	na	1
Rutile	0.00	na	0

Reference: Kandler et al. (2007)

Size Range: 2.5 – 5 μm

Izaña, Tenerife, Spain (Lat: 28°19'N, Lon: 16°30'W)

Variable: Concentration

Note: From Volume Fractions Times Assumed Densities

Mineral	Mean (%)	StdDev (%)	N_O
Albite	14.39	na	1
Bloedite	0.76	na	1
Calcite	5.57	na	1
Carbonaceous	0.27	na	1
Chloritoid	23.43	na	1
Dolomite	0.98	na	1
Glauberite	0.00	na	1

Gypsum	0.79	na	1
Halite	0.74	na	1
Hematite	1.80	na	1
Illite	43.08	na	1
Ilmenite	0.00	na	1
Quartz	8.18	na	1
Rutile	0.00	na	1

Reference: Kandler et al. (2007) Size Range: 5 – 10 μm

Izaña, Tenerife, Spain (Lat: 28°19'N, Lon: 16°30'W)

Variable: Concentration

Note: From Volume Fractions Times Assumed Densities

Mineral	Mean (%)	StdDev (%)	N_O
Albite	15.45	na	1
Bloedite	0.00	na	1
Calcite	4.69	na	1
Carbonaceous	0.00	na	1
Chloritoid	26.17	na	1
Dolomite	0.99	na	1
Glauberite	0.00	na	1
Gypsum	0.80	na	1
Halite	0.00	na	1
Hematite	0.00	na	1
Illite	45.47	na	1
Ilmenite	0.00	na	1
Quartz	6.43	na	1
Rutile	0.00	na	1

Reference: Kandler et al. (2009) Size Range: < 10 μm

Tinfou, Morocco (Lat: 30.24°N, Lon: 5.6°W)

Variable: Surface Concentration

Note: calc. from volume and dust distr.

Mineral	Mean (%)	StdDev (%)	N_O
Calcite+Dolomite	12.64	na	1
Feldspar	33.85	na	1
Hematite	1.56	na	1
Gypsum	3.38	na	1
Illite	30.21	na	1
Kaolinite	6.01	na	1
Quartz	12.35	na	1

Reference: Kandler et al. (2009) Size Range: < 2 μm

Tinfou, Morocco (Lat: 30.24°N, Lon: 5.6°W)

Variable: Surface Concentration

Note: calc. from volume and dust distr.

Mineral	Mean (%)	StdDev (%)	N_O
Calcite+Dolomite	10.31	na	1

Feldspar	35.27	na	1
Hematite	1.88	na	1
Gypsum	3.32	na	1
Illite	31.49	na	1
Kaolinite	6.26	na	1
Quartz	11.47	na	1

Reference: Kandler et al. (2009) Size Range: < 20 μm

Tinfou, Morocco (Lat: 30.24°N, Lon: 5.6°W)

Variable: Surface Concentration

Note: calc. from volume and dust distr.

Mineral	Mean (%)	StdDev (%)	N_O
Calcite+Dolomite	11.51	na	1
Feldspar	33.42	na	1
Hematite	1.52	na	1
Gypsum	3.01	na	1
Illite	29.86	na	1
Kaolinite	5.95	na	1
Quartz	14.73	na	1

Reference: Kandler et al. (2009) Size Range: 20 – 32 μm

Tinfou, Morocco (Lat: 30.24°N, Lon: 5.6°W)

Variable: Surface Concentration

Note: calc. from volume and dust distr.

Mineral	Mean (%)	StdDev (%)	N_O
Calcite+Dolomite	8.16	na	1
Feldspar	32.39	na	1
Hematite	1.60	na	1
Gypsum	0.87	na	1
Illite	28.96	na	1
Kaolinite	5.78	na	1
Quartz	22.23	na	1

Reference: Kandler et al. (2009) Size Range: 2 – 20 μm

Tinfou, Morocco (Lat: 30.24°N, Lon: 5.6°W)

Variable: Surface Concentration

Note: calc. from volume and dust distr.

Mineral	Mean (%)	StdDev (%)	N_O
Calcite+Dolomite	11.72	na	1
Feldspar	33.10	na	1
Hematite	1.45	na	1
Gypsum	2.96	na	1
Illite	29.58	na	1
Kaolinite	5.90	na	1
Quartz	15.30	na	1

Reference: Kandler et al. (2009) Size Range: < 32 μm

Tinfou, Morocco (Lat: 30.24°N, Lon: 5.6°W)

Variable: Surface Concentration

Note: calc. from volume and dust distr.

Mineral	Mean (%)	StdDev (%)	N_O
Calcite+Dolomite	10.35	na	1
Feldspar	33.06	na	1
Hematite	1.55	na	1
Gypsum	2.27	na	1
Illite	29.55	na	1
Kaolinite	5.89	na	1
Quartz	17.33	na	1

Reference: Kandler et al. (2011)

Size Range: Bulk

Praia, Cape Verde (Lat: 14°56'50.89"N, Lon: 23°29'4.31"W)

Variable: Surface concentration

Note: Weighted average of maritime and dust episodes means

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	2.18	0.41	2
Gypsum	4.37	1.47	2
Halite	4.02	7.13	2
Kaolinite	33.15	4.26	2
K-feldspar	20.65	2.38	2
Mixed-layer illite-smectite	13.88	2.21	2
Plagioclase	5.23	2.30	2
Quartz	10.63	1.47	2
Smectite	6.09	1.06	2

Reference: Khalaf et al. (1985)

Size Range: < 4 μm

Kuwait (Lat: Various, Lon: Various)

Variable: Total Dry Deposition

Note: Average of 7 sites

Mineral	Mean (%)	StdDev (%)	N_O
Albite	7.17	na	1
Calcite	14.62	na	1
Chlorite	5.07	na	1
Dolomite	4.57	na	1
Gypsum	2.88	na	1
Illite+mixed-layer illite-smectite	22.60	na	1
Kaolinite	1.98	na	1
K-feldspar	8.09	na	1
Palygorskite	25.35	na	1
Quartz	7.66	na	1

Reference: Leinen et al. (1994)

Size Range: < 2 μm

Glomar Challenger Cruise, Eastern Pacific (Lat: Various, Lon: Various)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	2.33	0.89	16

Illite	38.02	5.09	16
Kaolinite	18.20	5.11	16
Plagioclase	10.20	3.19	16
Quartz	6.89	2.05	16
Smectite	4.76	3.19	16

Reference: Leinen et al. (1994) Size Range: 2 – 20 μm
 Glomar Challenger Cruise, Eastern Pacific (Lat: Various, Lon: Various)
 Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	2.65	1.30	14
Illite	29.69	10.41	14
Kaolinite	7.73	3.17	14
Plagioclase	36.42	15.21	14
Quartz	20.64	4.04	14
Smectite	2.88	1.86	14

Reference: Leinen et al. (1994) Size Range: < 2 μm
 Glomar Challenger Cruise, Northwestern Pacific (Lat: Various, Lon: Various)
 Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	2.33	0.89	16
Illite	38.02	5.09	16
Kaolinite	18.20	5.11	16
Plagioclase	10.20	3.19	16
Quartz	6.89	2.05	16
Smectite	4.76	3.19	16

Reference: Leinen et al. (1994) Size Range: 2 – 20 μm
 Glomar Challenger Cruise, Northwestern Pacific (Lat: Various, Lon: Various)
 Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	2.65	1.30	14
Illite	29.69	10.41	14
Kaolinite	7.73	3.17	14
Plagioclase	36.42	15.21	14
Quartz	20.64	4.04	14
Smectite	2.88	1.86	14

Reference: Lu et al. (2006) Size Range: < 10 μm
 BeijingI, China (Lat: 40°00'N, Lon: 116°46'E)
 Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	10.91	9.30	12
Carbon particles	10.31	9.90	12
Compounds	11.95	8.23	12
Dolomite	3.79	1.85	12

Feldspar	7.55	4.36	12
Gypsum	9.49	9.91	12
Other minerals	2.65	4.07	12
Phyllosilicates	30.07	14.11	12
Quartz	13.54	3.36	12

Reference: Menéndez et al. (2007)

Size Range: Bulk

Gran Canaria, Spain (Lat: 27°58'N, Lon: 15°36'W)

Variable: Total Deposition

Mineral	Mean (%)	StdDev (%)	N_O
Aragonite	2.00	5.00	52
Calcite	11.00	8.00	52
Dolomite	8.00	6.00	52
Feldspar	11.00	8.00	52
Halite	1.10	1.80	52
Illite	0.10	0.10	52
Kaolinite-chlorite	0.10	0.10	52
Magnetite	7.00	8.00	52
Mg calcite	19.00	13.00	52
Palygorskite	0.02	0.07	52
Quartz	42.00	22.00	52

Reference: Møberg et al. (1991)

Size Range: < 2 μm

Zaria, Nigeria (Lat: 11°04'S, Lon: 7°42'E)

Variable: Total Dry Deposition

Note: Average of maximum and minimum

Mineral	Mean (%)	StdDev (%)	N_O
Al-chlorite	11.50	na	1
Al-Fe Oxyhydroxides	3.50	na	1
Ca-Na-feldspar	1.50	na	1
Fe-rich illite	15.50	na	1
Kaolinite	25.00	na	1
K-feldspar	1.50	na	1
Quartz	10.50	na	1
Smectite	20.00	na	1
Vermiculite	7.00	na	1

Reference: O'Hara et al. (2006)

Size Range: Bulk

Libya North (Lat: Various, Lon: Various)

Variable: Total Deposition

Mineral	Mean (%)	StdDev (%)	N_O
Albite	1.89	1.78	12
Anorthocite	1.55	1.70	12
Calcite	24.67	5.96	12
Chlorite	2.06	2.55	12
Dolomite	3.12	1.42	12
Gypsum	3.20	1.46	12

Halite	11.49	7.40	12
Illite	2.43	1.69	12
Kaolinite	2.69	0.97	12
Orthoclase	2.66	2.64	12
Palygorskite	1.65	0.87	12
Quartz	42.27	11.17	12

Reference: O'Hara et al. (2006) Size Range: Bulk
 Libya South (Lat: Various, Lon: Various)
 Variable: Total Deposition

Mineral	Mean (%)	StdDev (%)	N_O
Albite	2.05	2.50	12
Anorthocite	1.00	1.27	12
Calcite	9.84	6.56	12
Chlorite	2.80	2.03	12
Dolomite	1.65	1.33	12
Gypsum	3.99	2.47	12
Halite	4.18	2.30	12
Illite	2.84	1.79	12
Kaolinite	6.61	2.49	12
Orthoclase	2.17	2.35	12
Palygorskite	2.35	1.30	12
Quartz	60.31	14.49	12

Reference: Parkin et al. (1970) Size Range: Bulk
 M. V. Nova Scotia (Lat: Various, Lon: Various)
 Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Illite	13.17	1.39	2
Kaolinite+chlorite	41.42	7.67	2
Montmorillonite	10.42	0.70	2
Quartz	12.25	2.09	2
Talc	22.75	6.27	2

Reference: Parkin et al. (1970) Size Range: Bulk
 M. V. Samaria (Lat: Various, Lon: Various)
 Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Illite	22.41	5.97	10
Kaolinite+chlorite	24.83	8.37	10
Montmorillonite	5.43	2.26	10
Quartz	41.19	11.64	10
Talc	5.99	5.07	10

Reference: Parkin et al. (1972) Size Range: Bulk
 S. S. Hardwicke Grange (Lat: Various, Lon: Various)
 Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Illite	21.78	8.04	28
Kaolinite+chlorite	32.99	12.21	28
Montmorillonite	4.95	3.45	28
Quartz	36.61	11.20	28
Talc	4.09	11.39	28

Reference: Prospero and Bonatti (1969)

Size Range: < 20 μm

R.V. Pillsbury (Lat: Various, Lon: Various)

Variable: Surface Concentration

Note: Legs 3/1 and 11/2

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	2.00	1.41	2
Illite	2.50	2.12	2
Kaolinite	6.75	6.01	2
Muscovite	2.50	2.12	2
Plagioclase	26.95	28.07	2
Quartz	6.05	3.61	2
Smectite	16.50	19.80	2

Reference: Prospero et al. (1981)

Size Range: Bulk

Barbados (Lat: 13°10'N, Lon: 59°30'W)

Variable: Surface Concentration

Note: Mica includes illite

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	5.00	na	1
Chlorite	4.00	na	1
Gibbsite	0.00	na	1
Goethite	2.00	na	1
Gypsum	3.00	na	1
Kaolinite	6.00	na	1
Mica	60.00	na	1
Microcline	3.00	na	1
Plagioclase	6.00	na	1
Quartz	10.00	na	1

Reference: Prospero et al. (1981)

Size Range: Bulk

Cayenne, French Guiana (Lat: 4°50'N, Lon: 52°22'W)

Variable: Surface Concentration

Note: Mica includes illite

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	4.00	na	1
Chlorite	5.00	na	1
Gibbsite	1.00	na	1
Goethite	3.00	na	1
Gypsum	3.00	na	1

Kaolinite	8.00	na	1
Mica	59.00	na	1
Microcline	3.00	na	1
Plagioclase	5.00	na	1
Quartz	10.00	na	1

Reference: Prospero et al. (1981) Size Range: Bulk

Cayenne Mult, French Guiana (Lat: 4°50'N, Lon: 52°22'W)

Variable: Surface Concentration

Note: Mica includes illite

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	2.92	2.83	25
Chlorite	4.52	0.71	25
Gibbsite	7.72	9.89	25
Goethite	8.80	7.07	25
Gypsum	1.52	0.71	25
Kaolinite	7.60	3.53	25
Mica	44.72	25.44	25
Microcline	1.52	0.71	25
Plagioclase	6.44	2.12	25
Quartz	14.24	9.19	25

Reference: Prospero et al. (1981) Size Range: Bulk

Dakar, Senegal (Lat: 14°44'N, Lon: 17°33'W)

Variable: Surface Concentration

Note: Mica includes illite

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	5.00	na	1
Chlorite	7.00	na	1
Gibbsite	0.00	na	1
Goethite	2.00	na	1
Gypsum	1.00	na	1
Kaolinite	7.00	na	1
Mica	57.00	na	1
Microcline	2.00	na	1
Plagioclase	5.00	na	1
Quartz	14.00	na	1

Reference: Queralt-Mitjans et al. (1993) Size Range: Bulk

Filabres Range, Spain (Lat: Various, Lon: Various)

Variable: Bulk Deposition

Note: Date for Event 6 guessed

Mineral	Mean (%)	StdDev (%)	N_O
Aragonite	0.26	0.61	25
Calcite	0.58	1.00	25
Chlorite	13.52	9.24	25
Dolomite	0.94	1.33	25

Feldspar	4.04	2.17	25
Gypsum	1.32	2.49	25
Kaolinite	9.16	8.63	25
Mica-illite	54.60	12.80	25
Palygorskite	3.68	5.84	25
Paragonite	6.92	7.53	25
Quartz	4.92	2.93	25

Reference: Rashki et al. (2013)

Size Range: < 75 μm

Hamoun A, Iran (Lat: 31.10N, Lon: 61.51E)

Variable: Total Dry Deposition

Note: derived from figures; dust storms

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	22.07	2.63	15
Chlorite	6.33	1.40	15
Diopside	0.67	1.11	15
Dolomite	2.13	1.68	15
Enstatite	3.33	3.54	15
Gypsum	1.00	1.60	15
Halite	1.60	1.88	15
Hornblende	0.80	0.77	15
Illite	12.93	3.49	15
Microcline	1.33	1.50	15
Orthoclase	1.13	1.25	15
Plagioclase	11.20	1.32	15
Quartz	35.27	3.97	15

Reference: Rashki et al. (2013)

Size Range: < 75 μm

Hamoun B, Iran (Lat: 31.20N, Lon: 61.61E)

Variable: Total Dry Deposition

Note: derived from figures; dust storms

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	18.56	3.78	9
Chlorite	5.89	0.93	9
Diopside	0.17	0.35	9
Dolomite	4.22	1.56	9
Enstatite	1.22	1.46	9
Gypsum	0.78	0.71	9
Halite	0.83	0.61	9
Hornblende	0.83	0.87	9
Illite	10.00	3.50	9
Microcline	1.39	0.82	9
Orthoclase	3.17	2.63	9
Plagioclase	12.72	1.64	9
Quartz	40.22	4.41	9

Reference: Shao et al. (2008)

Size Range: < 2 μm

Beijing, China (Lat: 39°59'37.1"N, Lon: 116°20'45.6"E)

Variable: Total dry deposition

Note: Dust events

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	9.50	0.71	2
Illite	34.50	0.71	2
Kaolinite	8.50	0.71	2
Interstratified illite-smectite	47.50	2.12	2

Reference: Shao et al. (2008)

Size Range: Bulk

Beijing, China (Lat: 39°59'37.1"N, Lon: 116°20'45.6"E)

Variable: Total dry deposition

Note: Dust events

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	12.88	4.07	4
Clay	31.02	2.10	4
Dolomite	6.88	3.29	4
Gypsum	1.25	0.65	4
Hornblende	1.18	0.76	4
K-feldspar	3.82	0.88	4
Plagioclase	11.95	1.70	4
Quartz	31.02	4.31	4

Reference: Shen et al. (2006)

Size Range: Bulk

Dunhuang, China (Lat: 40°30'N, Lon: 94°49'E)

Variable: Surface Concentration

Note: Dust Storms

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	8.64	2.71	11
Chlorite	11.66	1.47	11
Dolomite	4.57	1.17	11
Illite	41.17	5.68	11
Kaolinite	5.94	1.11	11
K-feldspar	1.14	0.37	11
Plagioclase	5.29	1.65	11
Quartz	21.50	2.93	11

Reference: Shen et al. (2009)

Size Range: Bulk

Aksu, China (Lat: 40°16'N, Lon: 80°28'E)

Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	11.00	1.00	11
Dolomite	3.00	1.00	11
Illite	38.00	3.00	11
Kaolinite+chlorite	15.00	2.00	11
K-feldspar	2.00	1.00	11
Plagioclase	7.00	2.00	11

Quartz	24.00	3.00	11
Reference: Shen et al. (2009)			Size Range: Bulk
Changwu, China (Lat: 35°12'N, Lon: 107°40'30"N)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Calcite	5.00	2.00	16
Dolomite	2.00	1.00	16
Illite	50.00	3.00	16
Kaolinite+chlorite	23.00	2.00	16
K-feldspar	1.00	1.00	16
Plagioclase	4.00	2.00	16
Quartz	15.00	3.00	16
Reference: Shen et al. (2009)			Size Range: Bulk
Dunhuang, China (Lat: 40°30'N, Lon: 94°49'E)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Calcite	9.00	3.00	35
Dolomite	4.00	4.00	35
Illite	43.00	2.00	35
Kaolinite+chlorite	18.00	2.00	35
K-feldspar	1.00	1.00	35
Plagioclase	6.00	2.00	35
Quartz	19.00	4.00	35
Reference: Shen et al. (2009)			Size Range: Bulk
Tongliao, China (Lat: 43°14'20"N, Lon: 122°14'E)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Calcite	1.00	1.00	11
Dolomite	0.00	1.00	11
Illite	37.00	7.00	11
Kaolinite+chlorite	13.00	2.00	11
K-feldspar	6.00	2.00	11
Plagioclase	15.00	8.00	11
Quartz	26.00	6.00	11
Reference: Shen et al. (2009)			Size Range: Bulk
Zhenbeitai, China (Lat: 38°17'23"N, Lon: 109°42'18"E)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Calcite	5.00	3.00	37
Dolomite	2.00	1.00	37
Illite	43.00	5.00	37
Kaolinite+chlorite	20.00	2.00	37
K-feldspar	3.00	1.00	37

Plagioclase	8.00	2.00	37
Quartz	20.00	5.00	37

Reference: Shi et al. (2005) Size Range: < 10 μm
Beijing, China (Lat: 40°00'N, Lon: 116°46'E)
Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Carbonates	8.30	1.13	2
Iron oxides	0.45	0.64	2
Gypsum	0.15	0.21	2
Hornblende	0.35	0.07	2
K-feldspar	1.80	0.42	2
Noncrystal	20.25	0.35	2
Phyllosilicates	39.55	0.78	2
Plagioclase	8.10	0.42	2
Pyrite	0.95	0.07	2
Quartz	20.10	0.85	2

Reference: Shi et al. (2005) Size Range: < 2 μm
Beijing, China (Lat: 40°00'N, Lon: 116°46'E)
Variable: Surface Concentration
Note: PM2: Clays only

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	6.50	0.71	2
Kaolinite	5.50	0.71	2
Mixed-layer illite-smectite	78.50	0.71	2
Illite	9.50	0.71	2

Reference: Shi et al. (2005) Size Range: Bulk
Beijing, China (Lat: 40°00'N, Lon: 116°46'E)
Variable: Surface Concentration

Mineral	Mean (%)	StdDev (%)	N_O
Carbonates	14.00	na	1
K-feldspar	1.70	na	1
Noncrystal	10.10	na	1
Phyllosilicates	35.50	na	1
Plagioclase	7.00	na	1
Pyrite	1.00	na	1
Quartz	30.30	na	1

Reference: Skonieczny et al. (2013) Size Range: < 30 μm
Mbour, Senegal (Lat: 14°24'38"N, Lon: 16°57'32"W)
Variable: Total deposition
Note: Al-Si mineralogy

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	13.91	3.86	139
Illite	14.30	5.47	139

Kaolinite	25.95	5.34	139
Palygorskite	6.38	2.48	139
Smectite	39.47	10.39	139
Reference: Tomadin et al. (1984)			Size Range: < 2 μm
Adria to Strait of Messina (Lat: Various, Lon: Various)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	16.86	7.59	5
Illite	63.28	5.17	5
Kaolinite	13.89	4.58	5
Serpentine	3.67	3.41	5
Smectite	2.15	3.96	5
Reference: Tomadin et al. (1984)			Size Range: < 2 μm
Gulf of Saronikis to Adria (Lat: Various, Lon: Various)			
Variable: Surface Concentration			
Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	13.44	5.01	8
Illite	68.79	6.42	8
Kaolinite	9.11	2.15	8
Serpentine	6.45	4.01	8
Smectite	2.21	3.94	8
Reference: Tomadin et al. (1984)			Size Range: < 2 μm
Italy (Lat: Various, Lon: Various)			
Variable: Total Deposition			
Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	11.00	6.56	3
Illite	62.67	8.62	3
Kaolinite	20.67	13.32	3
Serpentine	0.00	0.00	3
Smectite	5.67	2.08	3
Reference: Windom (1969)			Size Range: Bulk
Camp Century, Greenland (Lat: 77°10'N, Lon: 61°08'W)			
Variable: Total Deposition			
Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	6.00	na	1
Chlorite	11.00	na	1
Feldspar	23.00	na	1
Kaolinite	9.00	na	1
Mica	27.00	na	1
Montmorillonite	0.00	na	1
Quartz	24.00	na	1
Reference: Windom (1969)			Size Range: Bulk
Franz Josef Glacier, New Zealand (Lat: 43°28'N, Lon: 170°11'E)			

Variable: Total Deposition

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	0.00	na	1
Chlorite	5.00	na	1
Feldspar	28.14	na	1
Kaolinite	2.51	na	1
Mica	25.94	na	1
Montmorillonite	0.00	na	1
Quartz	38.40	na	1

Reference: Windom (1969)

Size Range: Bulk

Mt. Olympus, Washington (Lat: 47°52'N, Lon: 123°35'W)

Variable: Total Deposition

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	0.85	na	1
Chlorite	7.71	na	1
Feldspar	12.73	na	1
Kaolinite	0.85	na	1
Mica	42.24	na	1
Montmorillonite	0.00	na	1
Quartz	35.04	na	1

Reference: Windom (1969)

Size Range: Bulk

Mt. Orizaba, Mexico (Lat: 19°01'N, Lon: 97°15'W)

Variable: Total Deposition

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	11.00	na	1
Chlorite	3.00	na	1
Feldspar	0.00	na	1
Kaolinite	6.00	na	1
Mica	16.00	na	1
Montmorillonite	6.00	na	1
Quartz	56.00	na	1

Reference: Windom (1969)

Size Range: Bulk

Mt. Popocatepetl, Mexico (Lat: 19°01'N, Lon: 98°37'W)

Variable: Total Deposition

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	8.00	na	1
Chlorite	0.00	na	1
Feldspar	0.00	na	1
Kaolinite	11.00	na	1
Mica	18.00	na	1
Montmorillonite	9.00	na	1
Quartz	53.00	na	1

Reference: Windom (1969)

Size Range: Bulk

Site 2, Greenland (Lat: 76°59'N, Lon: 56°04'W)

Variable: Total Deposition

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	5.00	na	1
Chlorite	11.00	na	1
Feldspar	16.00	na	1
Kaolinite	8.00	na	1
Mica	37.00	na	1
Montmorillonite	0.00	na	1
Quartz	23.00	na	1

Reference: Windom (1969)

Size Range: Bulk

St. Elias Range, Canada (Lat: 60°45'N, Lon: 139°30'W)

Variable: Total Deposition

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	13.45	na	1
Chlorite	5.81	na	1
Feldspar	10.10	na	1
Kaolinite	2.77	na	1
Mica	34.06	na	1
Montmorillonite	0.00	na	1
Quartz	32.84	na	1

Reference: Windom (1969)

Size Range: Bulk

Tasman Glacier, New Zealand (Lat: 43°31'S, Lon: 170°20'E)

Variable: Total Deposition

Mineral	Mean (%)	StdDev (%)	N_O
Amphibole	0.00	na	1
Chlorite	4.32	na	1
Feldspar	25.32	na	1
Kaolinite	3.34	na	1
Mica	26.50	na	1
Montmorillonite	0.00	na	1
Quartz	40.72	na	1

Reference: Zdanowicz et al. (2006)

Size Range: Bulk

St. Elias Mountains, Canada (Lat: Various, Lon: Various)

Variable: Total Deposition

Mineral	Mean (%)	StdDev (%)	N_O
Chlorite	27.00	0.00	2
Illite	65.50	7.78	2
Kaolinite	6.00	8.49	2
Smectite	0.75	0.35	2

Reference: Zhou and Tazaki (1996)

Size Range: Bulk

Matsue, Shimane Prefecture, Japan (Lat: 35°29'N, Lon: 133°04'E)

Variable: Surface concentration

Mineral	Mean (%)	StdDev (%)	N_O
Calcite	5.53	1.58	4
Clay	13.33	5.23	4
Gypsum	3.60	1.40	4
Halite	10.12	3.50	4
Quartz	10.35	6.81	4

20 References

- Adedokun, J. A., Emofurieta, W. O., and Adedeji, O. A.: Physical, Mineralogical and Chemical Properties of Harmattan Dust at Ile-Ife, Nigeria, *Theor. Appl. Climatol.*, 40, 161–169, doi:10.1007/BF00866179, <http://dx.doi.org/10.1007/BF00866179>, 1989.
- Al-Awadhi, J. M. and AlShuaibi, A. A.: Dust fallout in Kuwait city: Deposition and characterization, *Science of The Total Environment*, 461–462, 139–148, doi:10.1016/j.scitotenv.2013.03.052, <http://dx.doi.org/10.1016/j.scitotenv.2013.03.052>, 2013.
- Al-Dousari, A. M. and Al-Awadhi, J.: Dust fallout in Northern Kuwait, major sources and characteristics, *Kuwait J. Sci.*, 39, 171–187, 2012.
- Al-Dousari, A. M., Al-Awadhi, J., and Ahmed, M.: Dust fallout characteristics within global dust storm major trajectories, *Arab. J. Geosci.*, 6, 3877–3884, doi:10.1007/s12517-012-0644-0, <http://dx.doi.org/10.1007/s12517-012-0644-0>, 2013.
- Alastuey, A., Querol, X., Castillo, S., Escudero, M., Avila, A., Cuevas, E., Torres, C., Romero, P.-M., Exposito, F., García, O., Diaz, J. P., Van Dingenen, R., and Putaud, J. P.: Characterisation of TSP and PM_{2.5} at Izaña and Sta. Cruz de Tenerife (Canary Islands, Spain) during a Saharan Dust Episode (July 2002), *Atmos. Environ.*, 39, 4715–4728, doi:10.1016/j.atmosenv.2005.04.018, <http://dx.doi.org/10.1016/j.atmosenv.2005.04.018>, 2005.
- Arnold, E., Merrill, J., Leinen, M., and King, J.: The effect of source area and atmospheric transport on mineral aerosol collected over the North Pacific Ocean, *Global and Planetary Change*, 18, 137–159, doi:10.1016/S0921-8181(98)00013-7, [http://dx.doi.org/10.1016/S0921-8181\(98\)00013-7](http://dx.doi.org/10.1016/S0921-8181(98)00013-7), 1998.
- Aston, S. R., Chester, R., Johnson, L. R., and Padgham, R. C.: Eolian Dust From The Lower Atmosphere Of The Eastern Atlantic And Indian Oceans, China Sea And Sea Of Japan, *Marine Geology*, 14, 15–28, doi:10.1016/0025-3227(73)90040-6, [http://dx.doi.org/10.1016/0025-3227\(73\)90040-6](http://dx.doi.org/10.1016/0025-3227(73)90040-6), 1973.
- Avila, A., Queralt-Mitjans, I., and Alcarcón, M.: Mineralogical composition of African dust delivered by red rains over northeastern Spain, *J. Geophys. Res.*, 102, 21 977–21 996, doi:10.1029/97JD00485, 1997.
- Awadh, S. M.: Geochemistry and mineralogical composition of the airborne particles of sand dunes and dust storms settled in Iraq and their environmental impact, *Environ. Earth. Sci.*, 66, 2247–2256, doi:10.1007/s12665-011-1445-6, <http://dx.doi.org/10.1007/s12665-011-1445-6>, 2012.
- Chester, R. and Johnson, L. R.: Atmospheric Dusts Collected off the Atlantic Coast of North Africa and the Iberian Peninsula, *Marine Geology*, 11, 251–260, doi:10.1016/0025-3227(71)90027-2, [http://dx.doi.org/10.1016/0025-3227\(71\)90027-2](http://dx.doi.org/10.1016/0025-3227(71)90027-2), 1971a.
- Chester, R. and Johnson, L. R.: Atmospheric Dust Collected Off the West African Coast, *Nature*, 229, 105–107, doi:10.1038/229105b0, <http://dx.doi.org/10.1038/229105b0>, 1971b.
- Chester, R., Elderfield, H., and Griffin, J. J.: Dust transported in the North-east and South-east Trade Winds in the Atlantic Ocean, *Nature*, 233, 474–476, doi:10.1038/233474a0, <http://dx.doi.org/10.1038/233474a0>, 1971.
- Chester, R., Elderfield, H., Griffin, J. J., Johnson, L., and Padgham, R. C.: Eolian dust along the eastern margins of the Atlantic Ocean, *Marine Geology*, 13, 91–105, doi:10.1016/0025-3227(72)90048-5, [http://dx.doi.org/10.1016/0025-3227\(72\)90048-5](http://dx.doi.org/10.1016/0025-3227(72)90048-5), 1972.
- Chester, R., Baxter, G. G., Behairy, A. K. A., Connor, K., Cross, D., Elderfield, H., and Padgham, R. C.: Soil-sized eolian dusts from the lower troposphere of the eastern Mediterranean Sea, *Marine Geology*, 24, 201–217, doi:10.1016/0025-3227(77)90028-7, [http://dx.doi.org/10.1016/0025-3227\(77\)90028-7](http://dx.doi.org/10.1016/0025-3227(77)90028-7), 1977.
- Chester, R., Sharples, E. J., Sanders, G. S., and Saydam, A. C.: Saharan Dust Incursion Over The Tyrrhenian Sea, *Atmos. Environ.*, 18, 929–935, doi:10.1016/0004-6981(84)90069-6, [http://dx.doi.org/10.1016/0004-6981\(84\)90069-6](http://dx.doi.org/10.1016/0004-6981(84)90069-6), 1984.
- Delany, A. C., Delany, A. C., Parkin, D. W., Griffin, J. J., Goldberg, E. D., and Reimann, B. E. F.: Airborne dust collected at Barbados, *Geochimica et Cosmochimica Acta*, 31, 885–909, doi:10.1016/S0016-7037(67)80037-1, [http://dx.doi.org/10.1016/S0016-7037\(67\)80037-1](http://dx.doi.org/10.1016/S0016-7037(67)80037-1), 1967.
- Díaz-Hernández, J. L., Martín-Ramos, J. D., and López-Galindo, A.: Quantitative analysis of mineral phases in atmospheric dust deposited in the south-eastern Iberian Peninsula, *Atmos. Environ.*, 45, 3015–3024, doi:10.1016/j.atmosenv.2011.03.024, <http://dx.doi.org/10.1016/j.atmosenv.2011.03.024>, 2011.
- Enete, I. C., Obienusi, E. A., Igu, I. N., and Ayadiulo, R.: Harmattan Dust: Composition, Characteristics and Effects on Soil Fertility in Enugu, Nigeria, *British Journal of Applied Science & Technology*, 2, 72–81, doi:10.9734/BJAST/2012/950, <http://dx.doi.org/10.9734/BJAST/2012/950>, 2012.

- Engelbrecht, J. P., McDonald, E. V., Gillies, J. A., Jayanty, R. K. M., Casuccio, G., and Gertler, A. W.: Characterizing Mineral Dusts and Other Aerosols from the Middle East—Part 1: Ambient Sampling, *Inhalation Toxicology*, 21, 297–326, doi:10.1080/08958370802464273, <http://dx.doi.org/10.1080/08958370802464273>, 2009.
- Engelbrecht, J. P., Menéndez, I., and Derbyshire, E.: Sources of $PM_{2.5}$ impacting on Gran Canaria, Spain, *CATENA*, 117, 119–132, doi:10.1016/j.catena.2013.06.017, <http://dx.doi.org/10.1016/j.catena.2013.06.017>, 2014.
- Falkovich, A. H., Ganor, E., Levin, Z., Formenti, P., and Rudich, Y.: Chemical and mineralogical analysis of individual mineral dust particles, *J. Geophys. Res.*, 106, 18,029–18,036, doi:10.1029/2000JD900430, <http://dx.doi.org/10.1029/2000JD900430>, 2001.
- Ferguson, W. S., Griffin, J. J., and Goldberg, E. D.: Atmospheric Dust from the North Pacific – A short note on a long range eolian transport, *J. Geophys. Res.*, 75, 1137–1139, doi:10.1029/JC075i006p01137, <http://dx.doi.org/10.1029/JC075i006p01137>, 1970.
- Fiol, L. A., Fornós, J. J., Gelabert, B., and Guijarro, J. A.: Dust rains in Mallorca (Western Mediterranean): Their occurrence and role in some recent geological processes, *Catena*, 63, 64–84, doi:10.1016/j.catena.2005.06.012, <http://dx.doi.org/10.1016/j.catena.2005.06.012>, 2005.
- Formenti, P., Rajot, J. L., Desboeufs, K., Caquineau, S., Chevaillier, S., Nava, S., Gaudichet, A., Journet, E., Triquet, S., Alfaro, S., Chiari, M., Haywood, J., Coe, H., and Highwood, E.: Regional variability of the composition of mineral dust from western Africa: Results from the AMMA SOP0/DABEX and DODO field campaigns, *J. Geophys. Res.*, 113, D00C13, doi:10.1029/2008JD009903, <http://dx.doi.org/10.1029/2008JD009903>, 2008.
- Game, P. M.: Observations on a dustfall in the Eastern Atlantic, February, 1962, *Journal of Sedimentary Petrology*, 34, 355–359, doi:10.1306/74D7105F-2B21-11D7-8648000102C1865D, <http://jsedres.geoscienceworld.org/content/34/2/355.abstract>, 1964.
- Ganor, E.: The composition of clay minerals transported to Israel as indicators of Saharan dust emission, *Atmospheric Environment*, 25, 2657–2664, doi:10.1016/0960-1686(91)90195-D, [http://dx.doi.org/10.1016/0960-1686\(91\)90195-D](http://dx.doi.org/10.1016/0960-1686(91)90195-D), 1991.
- Ganor, E., Deutsch, Y., and Foner, H. A.: Mineralogical Composition and Sources of Airborne Settling Particles on Lake Kinneret (the Sea of Galilee), Israel, *Water, Air, and Soil Pollution*, 118, 245–262, doi:10.1023/A:1005167230795, <http://dx.doi.org/10.1023/A:1005167230795>, 2000.
- Gaudichet, A., Lefèvre, R., Gaudry, A., Ardouin, B., Lambert, G., and Miller, J. M.: Mineralogical composition of aerosols at Amsterdam Island, *Tellus*, 41, 344–352, doi:10.1111/j.1600-0889.1989.tb00313.x, <http://dx.doi.org/10.1111/j.1600-0889.1989.tb00313.x>, 1989.
- Gaudichet, A., Anglis, M. D., Joussaume, S., Petit, J. R., Korotkevitch, Y. S., and Petrov, V. N.: Comments on the origin of dust in East Antarctica for present and ice-age conditions, *J. Atmos. Chem.*, 14, 192–142, doi:10.1007/BF00115229, <http://dx.doi.org/10.1007/BF00115229>, 1992.
- Glaccum, R. A. and Prospero, J. M.: Saharan aerosols over the tropical North Atlantic — Mineralogy, *Marine Geology*, 37, 295–321, doi:10.1016/0025-3227(80)90107-3, [http://dx.doi.org/10.1016/0025-3227\(80\)90107-3](http://dx.doi.org/10.1016/0025-3227(80)90107-3), 1980.
- Goldberg, E. D. and Griffin, J. J.: The sediments of the Northern Indian Ocean, *Deep Sea Res.*, 17, 513–537, doi:10.1016/0011-7471(70)90065-3, [http://dx.doi.org/10.1016/0011-7471\(70\)90065-3](http://dx.doi.org/10.1016/0011-7471(70)90065-3), 1970.
- Jeong, G. Y.: Bulk and single-particle mineralogy of Asian dust and a comparison with its source soils, *J. Geophys. Res.*, 113, D02208, doi:10.1029/2007JD008606, <http://dx.doi.org/10.1029/2007JD008606>, 2008.
- Jeong, G. Y. and Achterberg, E. P.: Chemistry and mineralogy of clay minerals in Asian and Saharan dusts and the implications for iron availability, *Atmos. Chem. Phys. Discuss.*, 14, 15 735–15 770, doi:10.5194/acpd-14-15735-2014, <http://dx.doi.org/10.5194/acpd-14-15735-2014>, 2014.
- Jeong, G. Y., Kim, J. Y., Seo, J., Kim, G. M., Jin, H. C., and Chun, Y.: Long-range transport of giant particles in Asian dust identified by physical, mineralogical, and meteorological analysis, *Atmos. Chem. Phys.*, 14, 505–521, doi:10.5194/acp-14-505-2014, <http://dx.doi.org/10.5194/acp-14-505-2014>, 2014.
- Johnson, L. R.: Particle-Size Fractionation of Eolian Dust During Transport and Sampling, *Marine Geology*, 21, M17–M21, doi:10.1016/0025-3227(76)90099-2, [http://dx.doi.org/10.1016/0025-3227\(76\)90099-2](http://dx.doi.org/10.1016/0025-3227(76)90099-2), 1976.
- Kandler, K., Benker, N., Bundke, U., Cuevas, E., Ebert, M., Knippertz, P., Rodríguez, S., Schütz, L., and Weinbruch, S.: Chemical composition and complex refractive index of Saharan Mineral Dust at Izan a, Tenerife (Spain) derived by electron microscopy, *Atmospheric Environment*, 41, 8058–8074, doi:10.1016/j.atmosenv.2007.06.047, <http://dx.doi.org/10.1016/j.atmosenv.2007.06.047>, 2007.

- Kandler, K., Schütz, L., Deutscher, C., Ebert, M., Hofmann, H., Jäckel, S., Jaenicke, R., Knippertz, P., Lieke, K., Massling, A., Petzold, A., Schladitz, A., Weinzierl, B., Wiedensohler, A., Zorn, S., and Weinbruch, S.: Size distribution, mass concentration, chemical and mineralogical composition and derived optical parameters of the boundary layer aerosol at Tinfou, Morocco, during SAMUM 2006, *Tellus B*, 61, 32–50, doi:10.1111/j.1600-0889.2008.00385.x, <http://dx.doi.org/10.1111/j.1600-0889.2008.00385.x>, 2009.
- Kandler, K., Schütz, L., Jäckel, S., Lieke, K., Emmel, C., Müller-Ebert, D., Ebert, M., Scheuven, D., Schladitz, A., Šegvić, B., Wiedensohler, A., and Weinbruch, S.: Ground-based off-line aerosol measurements at Praia, Cape Verde, during the Saharan Mineral Dust Experiment: microphysical properties and mineralogy, *Tellus B*, 63, 459–474, doi:10.1111/j.1600-0889.2011.00550.x, <http://dx.doi.org/10.1111/j.1600-0889.2011.00550.x>, 2011.
- Khalaf, F. I., Al-Kadi, A., and Al-Saleh, S.: Mineralogical composition and potential sources of dust fall-out deposits in Kuwait, northern Arabian Gulf, *Sedimentary Geology*, 42, 255–278, doi:10.1016/0037-0738(85)90047-8, [http://dx.doi.org/10.1016/0037-0738\(85\)90047-8](http://dx.doi.org/10.1016/0037-0738(85)90047-8), 1985.
- Leinen, M., Prospero, J. M., Arnold, E., and Blank, M.: Mineralogy of aeolian dust reaching the North Pacific Ocean 1. Sampling and analysis, *J. Geophys. Res.*, 99, 21,017–21,023, doi:10.1029/94JD01735, <http://dx.doi.org/10.1029/94JD01735>, 1994.
- Lu, S., Shao, L., Wu, M., and Jiao, Z.: Mineralogical characterization of airborne individual particulates in Beijing PM_{10} , *J. Environ. Sci.*, 18, 90–95, <http://iospress.metapress.com/content/52ganfy7xb25a0ma/>, 2006.
- Menéndez, I., Díaz-Hernández, J. L., Mangas, J., Alonso, I., and Sánchez-Soto, P. J.: Airborne dust accumulation and soil development in the North-East sector of Gran Canaria (Canary Islands, Spain), *J. Arid Environ.*, 71, 57–81, doi:10.1016/j.jaridenv.2007.03.011, <http://dx.doi.org/10.1016/j.jaridenv.2007.03.011>, 2007.
- Møberg, J. P., Esu, I. E., and Malgwi, W. B.: Characteristics and constituent composition of Harmattan dust falling in Northern Nigeria, *Geoderma*, 48, 73–81, doi:10.1016/0016-7061(91)90007-G, [http://dx.doi.org/10.1016/0016-7061\(91\)90007-G](http://dx.doi.org/10.1016/0016-7061(91)90007-G), 1991.
- O’Hara, S. L., Clarke, M. L., and Elatrash, M. S.: Field measurements of desert dust deposition in Libya, *Atmos. Environ.*, 40, 3881–3897, doi:10.1016/j.atmosenv.2006.02.020, <http://dx.doi.org/10.1016/j.atmosenv.2006.02.020>, 2006.
- Parkin, D. W., Phillips, D. R., Sullivan, R. A. L., and Johnson, L.: Airborne dust collections over the North Atlantic, *J. Geophys. Res.*, 75, 1782–1793, doi:10.1029/JC075i009p01782, <http://dx.doi.org/10.1029/JC075i009p01782>, 1970.
- Parkin, D. W., Phillips, D. R., Sullivan, R., and Johnson, L.: Airborne dust collections down the Atlantic, *Quarterly Journal of the Royal Meteorological Society*, 98, 798–808, doi:10.1002/qj.49709841807, <http://dx.doi.org/10.1002/qj.49709841807>, 1972.
- Prospero, J. M. and Bonatti, E.: Continental Dust in the Atmosphere of the Eastern Equatorial Pacific, *J. Geophys. Res.*, 74, 3362–3371, doi:10.1029/JC074i013p03362, <http://dx.doi.org/10.1029/JC074i013p03362>, 1969.
- Prospero, J. M., Glaccum, R. A., and Nees, R. T.: Atmospheric transport of soil dust from Africa to South America, *Nature*, 289, 570–572, doi:10.1038/289570a0, <http://dx.doi.org/10.1038/289570a0>, 1981.
- Queralt-Mitjans, I., Domingo, F., and Sole-Benet, A.: The influence of local sources on the mineral content of bulk deposition over an altitudinal gradient in the Filabres Range (SE Spain), *J. Geop.*, 98, 16,761–16,768, doi:10.1029/93JD01281, <http://dx.doi.org/10.1029/93JD01281>, 1993.
- Rashki, A., Eriksson, P. G., de W. Rautenbach, C. J., Kaskaoutis, D. G., Grote, W., and Dykstra, J.: Assessment of chemical and mineralogical characteristics of airborne dust in the Sistan region, Iran, *Chemosphere*, 90, 227–236, doi:10.1016/j.chemosphere.2012.06.059, <http://dx.doi.org/10.1016/j.chemosphere.2012.06.059>, 2013.
- Shao, L., Li, W., Xiao, Z., and Sun, Z.: The mineralogy and possible sources of spring dust particles over Beijing, *Adv. Atmos. Sci.*, 25, 395–403, doi:10.1007/s00376-008-0395-8, <http://dx.doi.org/10.1007/s00376-008-0395-8>, 2008.
- Shen, Z., Cao, J., Li, X., Okuda, T., Wang, Y., and Wang, X.: Mass concentration and mineralogical characteristics of aerosol particles collected at Dunhuang during ACE-Asia, *Adv. Atmos. Sci.*, 23, 291–298, doi:10.1007/s00376-006-0291-z, <http://dx.doi.org/10.1007/s00376-006-0291-z>, 2006.
- Shen, Z., Caquineau, S., Cao, J., Zhang, X., Han, Y., Gaudichet, A., and Gomes, L.: Mineralogical characteristics of soil dust from source regions in northern China, *Particuology*, 7, 507–512, doi:10.1016/j.partic.2009.10.001, <http://dx.doi.org/10.1016/j.partic.2009.10.001>, 2009.

- Shi, Z., Shao, L., Jones, T. P., and Lu, S.: Microscopy and mineralogy of airborne particles collected during severe dust storm episodes in Beijing, China, *J. Geophys. Res.*, 110, D01303, doi:10.1029/2004JD005073, <http://dx.doi.org/10.1029/2004JD005073>, 2005.
- Skonieczny, C., Bory, A., Bout-Roumazeilles, V., Abouchami, W., Galer, S. J. G., Crosta, X., Diallo, A., and Ndiaye, T.: A three-year time series of mineral dust deposits on the West African margin: Sedimentological and geochemical signatures and implications for interpretation of marine paleo-dust records, *Earth and Planetary Science Letters*, 364, 145–156, doi:10.1016/j.epsl.2012.12.039, <http://dx.doi.org/10.1016/j.epsl.2012.12.039>, 2013.
- Tomadin, L., Lenaz, R., Landuzzi, V., Mazzucotelli, A., and Vannucci, R.: Wind-blown dust over the Central Mediterranean, *Oceanologica Acta*, 7, 13–23, 1984.
- Windom, H. L.: Atmospheric Dust Records in Permanent Snowfields: Implications to Marine Sedimentation, *Geol. Soc. Amer. Bull.*, 80, 761–782, doi:10.1130/0016-7606(1969)80[761:ADRIPS]2.0.CO;2, [http://dx.doi.org/10.1130/0016-7606\(1969\)80\[761:ADRIPS\]2.0.CO;2](http://dx.doi.org/10.1130/0016-7606(1969)80[761:ADRIPS]2.0.CO;2), 1969.
- Zdanowicz, C., Hall, G., Vaive, J., Amelin, Y., Percival, J., Girard, I., Biscaye, P., and Bory, A.: Asian dustfall in the St. Elias Mountains, Yukon, Canada, *Geochimica et Cosmochimica Acta*, 70, 3493–3507, doi:10.1016/j.gca.2006.05.005, <http://dx.doi.org/10.1016/j.gca.2006.05.005>, 2006.
- Zhou, G. and Tazaki, K.: Seasonal variation of gypsum in aerosol and its effect on the acidity of wet precipitation on the Japan Sea side of Japan, *Atmospheric Environment*, 30, 3301–3308, doi:10.1016/1352-2310(96)00071-4, [http://dx.doi.org/10.1016/1352-2310\(96\)00071-4](http://dx.doi.org/10.1016/1352-2310(96)00071-4), 1996.