Supplement to the manuscript of Lossow et al.:

THE SPARC WATER VAPOUR ASSESSMENT II: COMPARISON OF ANNUAL, SEMI-ANNUAL AND QBO VARIATIONS IN STRATOSPHERIC AND LOWER MESOSPHERIC WATER VAPOUR AS OBSERVED FROM SATELLITE

Version: Wednesday 23rd November, 2016 13:51 CET

In the main manuscript we have shown in Figs. 1, 6 and 11 examples of the latitude-altitude distribution of amplitude and phase for the different variability patterns that are the focus of this work. For the sake of completeness we show here these distributions for all data sets. Figure 1 provides the overview for the annual variation, Fig. 2 focuses on the semi-annual variation and Fig. 3 correspondingly on the QBO variation. All figures extend over multiple pages to fit the panels for all data sets. For the annual and semi-annual variation all 32 data sets that are part of the this study are shown. For the QBO variation in total 28 data sets are included, lacking the MIPAS V5H NOM data sets which cover only a period of 21 months. The key features for the different variability patterns described in the main manuscript (see Sects. 4.1.1, 4.2.1 and 4.3.1) are well depicted in many data sets. The MIPAS and MLS data sets provide results over almost the entire latitude-altitude domain considered in this work. Hence all the features can be addressed. Also the HALOE, SAGE II and SMR 489 GHz data sets cover large parts of the domain. The remaining data sets provide a spatial snapshot covering some of the key features, but not in all cases. Besides providing a good overview of the spatial coverage of the individual data sets the figures also exhibit the kind of outliers in the variability amplitudes, often towards their vertical boundaries, which we screened to provide reasonable estimates of the uncertainties of the variability characteristics (see Sect. 3.3).

annual variation

Continued on next page

annual variation

Continued on next page

annual variation

annual variation

Continued on next page

annual variation

Figure 1: Latitude-altitude cross sections of amplitude (left panels) and phase (right panels) of the annual variation in water vapour for the satellite data sets considered in this analysis. The phase is given by the month in which water vapour exhibits its annual maximum in the regression fit. The light grey and white line indicates the mean tropopause (2000 – 2014) as derived from the MERRA reanalysis data. White boxes indicate that there are no data.

semi-annual variation

Continued on next page

semi-annual variation

Continued on next page

semi-annual variation

Continued on next page

semi-annual variation

Continued on next page

semi-annual variation

Continued on next page

Figure 2: Latitude-pressure cross-section of amplitude (left panels) and phase (right panels) of the semi-annual variation in water vapour for the different satellite data sets. The phase represents the month of a calendar year were the first water vapour maximum is found in the semi-annual regression fit. The light grey and white line indicates the mean tropopause (2000 – 2014) as derived from MERRA. Note that the amplitude scale is different than in Fig. 1.

quasi-biennial variation

Continued on next page

quasi-biennial variation

Continued on next page

quasi-biennial variation

Continued on next page

quasi-biennial variation

Continued on next page

Figure 3: Amplitude (left panels) and phase (right panels) of the quasi-biennial variation in water vapour. The phase is defined as the shift of the QBO regression fit for which the correlation with the Singapore (1°N , 104°E) winds at 50 hPa maximises.