Supplemental Information

Detection of Ship Plumes from Residual Fuel Operation in Emission Control Areas using Single-Particle Mass Spectrometry

by J. Passig et al, 2020

Table S1: Assignment of the top-300 clusters of the ART-2a analysis to main particle classes.

EC-OC	OC-EC	Sulfate-nitrate	K-CN	aged sea salt	V-Fe-Ni	other	remarks
3	1	24	2	57	72		
5	7	30	4	80	97	43	NO+
8	14	31	6	82	110	44	NO+
10	18	32	9	106	111	45	NO+
11	20	39	12	120	127	89	Al+ Fe+ org
13	26	41	21	132	138	152	low quality
15	27	51	23	146	150	174	Ca+, large
16	28	52	34	155	151	184	Al+ org.
17	33	62	35	159	161	206	Fe+43
19	40	64	42	177	164	212	EC-S
22	43	69	47	180	183	222	MGO?
25	44	70	49	193	196	236	Fe+ Ni+
29	45	73	51	225	226	256	low quality
36	53	75	56	234	230	287	Fe+
37	54	95	65	246	250	296	Fe+
38	59	96	67	252	265		
44	66	100	84	253			
46	68	101	86	254			
48	74	104	90	257			
50	77	105	83	259			
55	78	112	85	260			
58	87	115	91	266			
60	88	117	92	267			
63	102	121	98	287			
71	107	124	114				
76	109	126	118				
79	113	128	122				
81	116	130	125				
93	123	131	129				
94	135	136	133				
99	139	137	140				
103	141	143	149				
108	182	147	153				
119	187	148	154				
134	199	156	157				
142	200	158	192				
144	205	160	210				
145	215	162	211				
163	216	165	213				
170	219	166	224				
171	223	167	235				
178	228	168	261				
185	229	169	282				
186	231	172	281				
190	232	173					
194	236	175					
201	238	176					
202	242	179					
204	243	181					

207	244	188
208	247	189
214	249	191
218	251	195
221	258	197
222	264	198
227	268	203
233	275	209
239	279	217
245	283	220
248	284	240
255	280	241
262	286	263
269	289	270
273	290	272
276	292	274
277	293	291
278	297	295
285		
283		
284		
288		
294		
298		
299		
300

Figure S2: HYSPLIT back trajectories for 24h run time, ending at the measurement site during the experiments.

NOAA HYSPLIT MODEL
Backward trajectories ending at 2000 UTC 27 Jun 18
GFSG Meteorological Data

NOAA HYSPLIT MODEL
Backward trajectories ending at 2000 UTC 28 Jun 18
GFSG Meteorological Data

NOAA HYSPLIT MODEL
Backward trajectories ending at 2000 UTC 29 Jun 18
GFSG Meteorological Data

NOAA HYSPLIT MODEL
Backward trajectories ending at 2000 UTC 30 Jun 18
GFSG Meteorological Data

NOAA HYSPLIT MODEL
Backward trajectories ending at 2000 UTC 01 Jul 18
GFSG Meteorological Data

Figure S3: HYSPLIT back trajectories for 48h run time, ending at the measurement site during the experiments.

Figure S4: Weight matrices (neg. and pos. mass spectra) of the top 300 cluster centers from the ART-2a analysis.

Cluster 81 to 100

1 #cluster 2 number of particles 3 median of aerodynamic diameter

Cluster 221 to 240

1 #cluster 2 number of particles 3 median of aerodynamic diameter

Cluster 241 to 260

1 #cluster 2 number of particles 3 median of aerodynamic diameter

Cluster 261 to 280

1. #cluster 2. number of particles 3. median of aerodynamic diameter

Cluster 301 to 320

1 #cluster 2 number of particles 3 median of aerodynamic diameter

Figure S5: Time series of the top 300 clusters from the ART-2a analysis.