Supplemental material:
Coastal processes modify projections of some climate-driven stressors in the California
Current System

S.A. Siedlecki^{1*}, D. Pilcher², E.M. Howard³, C. Deutsch³, P. MacCready³, E.L. Norton², H. Frenzel³, J. Newton⁴, R.A. Feely⁵, S.R. Alin⁵, and T. Klinger⁶

¹Department of Marine Sciences, University of Connecticut, 1080 Shennecossett Road, Groton, CT 06340, USA

²CICOES, University of Washington, 3737 Brooklyn Ave NE, Seattle, WA 98195, USA.

³School of Oceanography, University of Washington, 1503 NE Boat Street, Seattle, WA 98195, USA.

⁴APL, University of Washington, 1013 NE 40th Street, Seattle, WA 98105, USA.

⁵NOAA Pacific Marine Environmental Lab (PMEL), 7600 Sand Point Way NE, Seattle, WA 98115, USA.

⁶School of Marine Environment and Affairs, University of Washington, 3707 Brooklyn Ave NE, Seattle, WA 98105, USA.

Corresponding author: Samantha Siedlecki (samantha.siedlecki@uconn.edu)

Figure S1. Global changes from the CMIP5 ensemble average for the CCS region - future minus base conditions (a) temperature (deg C) (b) O₂ (ml/l) (c) pCO₂ (μ atm) (d) pH.