



Nitrate assimilation and regeneration in the Barents Sea: insights from nitrogen isotopes

Robyn E. Tuerena^{1*}, Joanne Hopkins², Raja S. Ganeshram¹, Louisa Norman³, Camille de la Vega³, Rachel Jeffreys³ and Claire Mahaffey³

¹School of GeoSciences, University of Edinburgh, James Hutton Rd, Edinburgh, EH9 3FE, UK.

²National Oceanography Centre, 6 Brownlow Street, Liverpool, L3 5DA, UK.

³School of Environmental Sciences, University of Liverpool, 4 Brownlow St, Liverpool, L69 3GP, UK.

10 *Correspondence to:* Robyn E. Tuerena (r.tuerena@ed.ac.uk)

Abstract. While the entire Arctic Ocean is warming rapidly, the Barents Sea in particular is experiencing significant warming and sea ice retreat. An increase in ocean heat transport from the Atlantic is causing the Barents Sea to be transformed from a cold, salinity stratified system into a warmer, less-stratified Atlantic-dominated climate regime. Productivity in the Barents Sea shelf is fuelled by waters of Atlantic origin (AW), which are ultimately exported to the Arctic basin. The consequences of this current regime shift on the nutrient characteristics of the Barents Sea are poorly defined. Here we use the stable isotopic ratios of nitrate ($\delta^{15}\text{N-NO}_3$, $\delta^{18}\text{O-NO}_3$), to determine the uptake and modification of AW nutrients in the Barents Sea. In summer months, phytoplankton consume nitrate, surface waters become nitrate depleted, and particulate nitrogen ($\delta^{15}\text{N-PN}$) reflects the AW nitrate source. The ammonification of organic matter in shallow sediments resupplies N to the water column through the season. Low $\delta^{18}\text{O-NO}_3$ in the northern Barents Sea reveals that the nitrate in lower temperature Arctic Waters is >80% regenerated through seasonal nitrification. During on shelf nutrient uptake and regeneration, there is no significant change to $\delta^{15}\text{N-NO}_3$ or N^* , suggesting benthic denitrification does not impart an isotopic imprint on pelagic nitrate. Our results demonstrate that the Barents Sea is distinct from other Arctic shelves, where coupled partial nitrification-denitrification enriches $\delta^{15}\text{N-NO}_3$ and decreases N^* . Our results suggest that any current or future changes to productivity on the Barents Sea shelf are unlikely to alter the magnitude or isotopic signature of nutrient supply exported to the central Arctic basin. However, we suggest that the AW nutrient source ultimately determines Barents Sea productivity and changes to this supply may alter Barents Sea primary production and subsequent nutrient supply to the central Arctic Ocean.



30 1 Introduction

The Arctic Ocean is warming (Huang et al., 2017) and experiencing sea ice loss (Notz and Stroeve, 2016), and freshening (Coupel et al., 2015) as a direct response to climate change. It is an enclosed basin filled with waters from the Atlantic and Pacific Oceans, which provide varying concentrations of nutrients (Torres-Valdes et al., 2013). In turn, these nutrient supply pathways influence the distribution and extent of primary production throughout the Arctic Ocean (Lewis et al., 2020).
35 Approximately 50% of the Arctic Ocean is made up of productive shelves that support large fisheries and diverse habitats (Dalpadado et al., 2014; Friedland and Todd, 2012). As the Arctic continues to warm and more sea ice is lost, phytoplankton growth will become less limited by light availability. Instead, nutrient availability, principally nitrate (Codispoti et al., 2013), may become the primary control on phytoplankton growth (Arrigo and van Dijken, 2015; Lewis et al., 2020). A greater understanding of how nitrate is supplied to the Arctic inflow shelves and on shelf nutrient dynamics is therefore imperative in
40 order to predict changes to Arctic primary production and food web dynamics.

Atlantic Water (AW) is supplied to the Arctic via Fram Strait and the Barents Sea Opening and fills most of the deep basins of the Arctic. It supplies nutrients to the Eurasian shelves with nitrate and phosphate concentrations close to Redfield (15-16N:1P), and low concentrations of silicate, which can limit the extent of diatom growth (Hatun et al., 2017). AW is a mixture
45 of nutrient-rich North Atlantic subpolar and nutrient-poor subtropical origin water advected into the Norwegian Sea. Over the last two decades there has been a 7% and 20% decrease in nitrate and silicate concentrations respectively in the Barents Sea (Rey, 2012). This has been driven by shallower winter mixing in the subpolar gyre, coupled with weakening and westward retraction of the gyre which has increased the proportion of subtropical origin water entering the Norwegian Sea (Rey, 2012; Hatun et al., 2017).

50 As warm and saline AW inflow water is transported across the Barents Sea, it is modified by atmospheric cooling and is mixed with cold, fresh Arctic origin water (ArW) and the Norwegian Coastal Current (NCC). ArW found across the northern Barents Sea comprises fresh Arctic river runoff, sea ice melt and precipitation, and contains the remnants of the winter mixed layer (Rudels et al., 1996). It isolates the sea surface and ice cover from warm AW below (Lind et al., 2016) and during the summer
55 is capped by a well-mixed surface layer of fresh melt water (Polar Surface Water). Sea ice import from the Nansen Basin and Kara Sea is the most important source of freshwater in the northern Barents Sea (Lind et al., 2016; Ellingsen et al., 2009). The transition between AW and ArW is marked by the Polar Front, which can be identified from the sea surface temperature gradient (Barton et al., 2018; Oziel et al., 2016).



- 60 Intense cooling of AW across the Barents Sea, reinforced by brine rejection due to ice formation creates dense Barents Sea Water (BSW) that cascades into the deeper troughs of the central and eastern Barents Sea (Arthun et al., 2011; Oziel et al., 2016). BSW eventually leaves the shelf, mainly through St. Anna Trough (Smedsrud et al., 2013), where it is entrained into Arctic Intermediate Water and spreads further into the Arctic basin (Schauer et al., 1997).
- 65 The Barents Sea is experiencing a rapid decline in winter and summer sea ice cover (Onarheim and Arthun, 2017; Arthun et al., 2012), full-depth warming driven by both increased ocean heat transport from the Atlantic and amplified atmospheric warming over the Arctic (Arthun et al., 2012; Onarheim et al., 2015; Serreze et al., 2009), alongside increases in salinity (Lind et al., 2018; Barton et al., 2018). The area occupied by AW is increasing and the southern expression of the Polar Front is moving north (Oziel et al., 2016; Oziel et al., 2020). In the northern Barents Sea, a reduction in sea-ice import and therefore a
- 70 loss of freshwater is weakening stratification and enhancing vertical mixing (Lind et al., 2018). The northern Barents Sea is therefore transitioning from a cold, salinity stratified shelf into a warmer, less stratified Atlantic dominated climate regime (Lind et al., 2018), a process described as ‘Atlantification’.

On the other side of the Arctic, the Pacific Ocean supplies high concentrations of nutrients onto the Chukchi and East Siberian

75 shelves, fuelling productivity and nutrient uptake (Granger et al., 2011). Increases in volume transport through the Bering Strait in recent years (Woodgate, 2018) have increased Pacific nutrient supply to the Arctic basin. These waters are relatively depleted in nitrate (in comparison to phosphate), and combined with sedimentary denitrification on the shallow shelves (Fripiat et al., 2018; Granger et al., 2018), promote nitrogen limitation in the western Arctic Ocean (Mills et al., 2018).

- 80 Although many studies have found the western Arctic Ocean to be strongly N limited (Mills et al., 2018; Granger et al., 2018; Brown et al., 2015), we know less about the extent of N limitation and occurrence of sedimentary denitrification in the eastern Arctic Ocean. Nitrate isotope measurements can give integrated estimates of nitrogen cycling processes, yet there is currently no data on the North Atlantic inputs which provide nutrients to the Arctic basin via the Barents Sea. The $^{15/14}\text{N}$ and $^{18/16}\text{O}$ in nitrate ($\delta^{15}\text{N-NO}_3$ and $\delta^{18}\text{O-NO}_3$, respectively) provide complementary information about nitrate uptake by
- 85 phytoplankton and regeneration processes (Sigman et al., 2009b) and can be used to determine the relevant N cycling processes.

Nitrate consumption through algal uptake fractionates both N and O in a 1:1 ratio (Granger et al., 2004) with an isotope effect close to $\sim 5\text{‰}$ (Sigman et al., 2009b). Nitrogen loss through denitrification in the water column leads to enrichment in N and O isotopes in the residual nitrate pool with a fractionation of 25-30 ‰ (Sigman et al., 2009a). In sediments, denitrification does

90 not usually impart a signature on nitrate isotopes as N is efficiently lost to N_2 before fractionation is expressed (Sigman et al., 2003). However, multiple studies from the western Arctic and Bering Sea show that benthic denitrification can impart a signature on the overlying water column ($\epsilon = 0\text{--}5\text{‰}$) when high $\delta^{15}\text{N-NH}_4$ is released, a process termed coupled partial nitrification-denitrification or CPND (Brown et al., 2015).



95 Over most of the ocean fixed nitrogen is efficiently recycled in surface waters and the regeneration of nitrate in the water column retains a $\delta^{15}\text{N}$ signature from the N source (Sigman et al., 2000). The $\delta^{15}\text{N}$ signature imparted on nitrate can therefore be used to identify nutrient sources from partial utilization of nutrients (Rafter et al., 2012), different water masses (Sigman et al., 2000; Tuerena et al., 2015), new N inputs (Knapp et al., 2008; Marconi et al., 2017), atmospheric inputs (Altieri et al., 2016) and rivers (Thibodeau et al., 2017).

100

In contrast to $\delta^{15}\text{N}\text{-NO}_3$, where N atoms are internally recycled during nitrification, oxygen atoms are sourced from ambient O_2 and seawater, in general providing a nitrification signature of $\delta^{18}\text{O}\text{-H}_2\text{O}$ plus 1.1‰ (Buchwald et al., 2012; Sigman et al., 2009b). The contrasting sources of N and O atoms and thus their distinct isotopic signatures allow the relative importance of preformed and regenerated nitrate to be investigated (Rafter et al., 2013). $\delta^{18}\text{O}\text{-NO}_3$ has been used to quantify the extent of
 105 regeneration on the Bering Sea shelf (Granger et al., 2011) and as evidence for the significance of nitrate regeneration in sustaining nutrient stocks on Arctic Shelves (Fripiat et al., 2018; Granger et al., 2018).

When nitrate is not fully consumed in surface waters, $\delta^{15}\text{N}\text{-NO}_3$ and $\delta^{18}\text{O}\text{-NO}_3$ can indicate the extent of seasonal nitrate uptake by phytoplankton (DiFiore et al., 2009). The $\delta^{15}\text{N}$ of surface water nitrate increases as nitrate is progressively utilized
 110 by phytoplankton, through the preferential consumption of ^{14}N (Sigman et al., 1999). Together with other N cycling processes this can be described by Rayleigh fractionation systematics (Mariotti et al., 1981). Nitrate utilisation by phytoplankton in an environment where there is no resupply of nutrients, i.e. a stratified upper ocean in summer, follows Rayleigh fractionation systematics for a closed system, with $\delta^{15}\text{N}\text{-NO}_3$ falling on a fractionation trend for its isotopic effect (ϵ) (Granger et al., 2004). In combination with dissolved nutrients, the $\delta^{15}\text{N}$ of particulate nitrogen ($\delta^{15}\text{N}\text{-PN}$) can track the extent of biological utilisation,
 115 contrasting nutrient sources, and the significance of new versus regenerated nutrients (Altabet and Francois, 1994).

In this study we report the first stable isotope measurements of dissolved and particulate N in the Barents Sea and use them to understand the relative sources of nutrients fuelling contemporary Barents Sea productivity. We use stable isotope tracers to investigate how N cycling processes vary across the Barents Sea, in contrast to other Arctic shelves, and reflect upon the
 120 susceptibility of the ecosystem to climate change.

2 Materials and Methods

Samples were collected in the Barents Sea as part of the ARISE project (NERC Changing Arctic Ocean programme). Shipboard measurements were taken from the RRS James Clark Ross during July-August 2017 (JR16006). A 2,200 km transect was completed, comprising 59 full depth CTD casts, starting from the northern tip of Norway and ending at the shelf edge north-



125 east of Svalbard (Figure 1a). The transect crossed the Barents Sea Opening (BSO), between Norway and the southern tip of
 Svalbard. Then, from Hopen Trench it continued north towards Kong Karls Land (along 30°E), and to the shelf edge and
 Nansen Basin.

Standard CTD measurements and water sampling were performed using a stainless steel rosette equipped with a full sensor
 130 array and twenty-four 20-litre OTE bottles. Conductivity, temperature, and pressure were measured using a CTD system
 (Seabird 911+). Derived salinity was calibrated on-board with discrete samples using an Autosal 8400B salinometer
 (Guildline) (Dumont et al., 2019) and an SBE43 oxygen sensor was calibrated against oxygen samples analysed using the
 Winkler method. A Biospherical QCP Cosine PAR sensor measured down-welling photosynthetically available radiation
 (PAR; 400-700 nm). We define the base of the euphotic zone to be the depth where PAR decreased to 1% of its surface value.
 135 The mean depth of the euphotic zone was 34.3 ± 11.9 m.

Dissolved inorganic nutrient concentrations were determined using a Bran and Luebbe QuAatro 5-channel auto analyser
 (SEAL Analytical) and AACE operating platform (V 6.1) following standard colorimetric methods with a CRM precision of
 0.3%, 0.8% and 1.9% for nitrate+nitrite, phosphate and nitrite, respectively. Nitrate isotope samples were collected and filtered
 140 inline from the CTD using an Acropak and were frozen at -20°C until analysis. $\delta^{15}\text{N}$ -PN samples were collected by gently
 vacuum filtering through combusted GF/F filters (450 °C, 4hr, Whatman, 48 mm or 25 mm, nominal pore size 0.7 µm) until a
 good colour was obtained on the filter (8 to 12 L for the 48 mm diameter filters and 2 to 5 L for the 25 mm diameter filters).
 The filters were dried at 60 °C to remove all moisture and were stored folded and wrapped in combusted aluminium foil until
 return to the home laboratory where they were placed in a -80 °C freezer until analysis. Of the 59 CTD casts in 2017, 23 were
 145 sampled for nitrate isotopes.

To quantify the distinct nutrient concentrations, nutrient ratios and isotopic values of Atlantic Water (AW), Barents Sea Water
 (BSW) and Arctic Water (ArW), we define the water mass type of each sample using the water mass properties in Oziel et al.
 (2016). These are summarised in Table 1.

150 Additional sampling was conducted in collaboration with the Norsk Institutt for Vannforskning (NIVA, Oslo) during transits
 made by the general cargo vessel M/S Norbjørn between Tromsø, Norway and Longyearbyen, Svalbard. The M/S Norbjørn
 is a ‘ship of opportunity’ onto which NIVA has fitted a FerryBox system that measures a variety of parameters including,
 temperature and salinity at approximately 4 m depth. In addition, seawater can be collected directly from the system for further
 155 analysis. During each 4 day transit in March, June, August and November 2018, surface seawater samples were collected for
 the analysis of $\delta^{15}\text{N}$ of particulate organic nitrogen ($\delta^{15}\text{N}$ -PN), the $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ of nitrate ($\delta^{15}\text{N}$ -NO₃, $\delta^{18}\text{O}$ -NO₃) and
 inorganic nutrients (nitrate+nitrite, nitrite, silicate, and phosphate) from 15 stations at pre-determined latitudes (Figure 1a).



Seawater was filtered through combusted GF/F filters for $\delta^{15}\text{N}$ -PN analysis and aliquots of the filtrate were placed into acid cleaned HDPE bottles and stored at $-20\text{ }^{\circ}\text{C}$ for the analysis of $\delta^{15}\text{N}$ -NO₃ and inorganic nutrients.

160

The isotopic composition of nitrate+nitrite ($\delta^{15}\text{N}$ -NO₃ and $\delta^{18}\text{O}$ -NO₃) was determined by the denitrifier method (Sigman et al., 2001; Casciotti et al., 2002) and following GEOTRACES protocols (Schlitzer et al., 2018). Samples were corrected using international reference standards N3 and USGS-34 (Weigand et al., 2016) and expressed in delta notation ($\delta^{15}\text{N}$ -NO₃ (‰ vs AIR) = $(R_{\text{sam}}/R_{\text{std}} - 1) \times 1000$, $\delta^{18}\text{O}$ -NO₃ (‰ vs VSMOW) = $(R_{\text{sam}}/R_{\text{std}} - 1) \times 1000$). Standards were run in triplicate with a reproducibility (σ) $\delta^{15}\text{N} \pm 0.1\text{‰}$ and $\delta^{18}\text{O} \pm 0.3\text{‰}$. Internal standards were analysed in each run and corrected using N3 and U34, with an inter-run standard deviation of $\delta^{15}\text{N} \pm 0.1\text{‰}$ and $\delta^{18}\text{O} \pm 0.3\text{‰}$. As the $\delta^{18}\text{O}$ in N₂O generated from nitrite is 25‰ lower than nitrate (Casciotti et al., 2007), we correct our $\delta^{18}\text{O}$ -NO₃ data following Kemeny et al., (2016). Our isotopic measurements are compared to studies where the nitrite in a sample has been removed using sulphamic acid (Granger and Sigman, 2009), to account for this, where nitrite was >2.5% of nitrate+nitrite, samples were re-run with sulphamic acid removal. For samples where nitrite was <2.5% of nitrate+nitrite, $\delta^{15}\text{N}$ -NO₃+NO₂ samples were corrected assuming a $\delta^{15}\text{N}$ -NO₂ of -24‰ (Kemeny et al., 2016).

170

$\delta^{15}\text{N}$ -PN was determined by EA-IRMS using a Costech Instruments Elemental Analyser coupled to Thermo Scientific Delta V Advantage mass spectrometer fitted with Conflo IV gas handling system. The instrumentation was operated using ISODAT 3.0 isotope ratio MS software. Prior to analysis the filters were wrapped in tin foil cones (OEA Laboratories) and pelletised.

175

3 Results

The Barents Sea Opening (BSO), between Norway and Svalbard, was dominated by saline ($S > 34.8$) Atlantic inflow, notably in Bear Island Trough (BIT, Figure 1a, Figure 2b). South of Spitzbergen Bank the water column was thermally stratified, with temperature exceeding $6\text{ }^{\circ}\text{C}$ in the euphotic zone (Figure 2a). South of $72\text{ }^{\circ}\text{N}$, low salinity ($S < 34.7$) water from the Norwegian Coast Current (NCC) occupied the near surface layer (Figure 2b). The water column was fresher, colder and well mixed over the shallow Spitzbergen Bank. This marks the western-most extent of ArW and the Polar Front (Figure 1b) and coincides with strong tidal currents and topographically steered flows (Oziel et al., 2016; Sundfjord et al., 2007; Vage et al., 2014).

180

Dense Barents Sea Water (BSW) was observed near the seabed in Hopen Trench (Figure 2a, HT). Above it lay cooled Atlantic origin water and a thermally stratified surface layer. North of the narrow sill joining the Spitzbergen and Great Banks (Figure 2a, SB-GB), the approximate location of the Polar Front, colder ($< 0\text{ }^{\circ}\text{C}$) and fresher ($S < 37.7$) ArW occupied depths below 50 m (Figure 2b). This was capped with an even fresher ($S < 34$) layer of sub-zero temperature melt water. This Polar Surface Layer extended southwards from the Nansen Basin, becoming progressively thinner. Below 100 m depth, over the shelf break

185



and continental slope of the Nansen Basin, high salinity (cooled) Atlantic origin water within the Boundary Current that had entered the Arctic via the Fram Strait was observed.

In the AW, nitrate concentrations were relatively homogenous below the mixed layer ($11.8 \pm 1.8 \mu\text{M}$) but low or below the limits of detection in the euphotic layer (Figure 2c). NH_4^+ concentrations were highest close to the seafloor over the Spitsbergen Bank (Figure 2d, Figure 4). $\delta^{15}\text{N}\text{-NO}_3$ and $\delta^{18}\text{O}\text{-NO}_3$ were relatively homogenous in the deeper AW ($\delta^{15}\text{N}\text{-NO}_3 = 5.1 \pm 0.1\text{‰}$, $\delta^{18}\text{O}\text{-NO}_3 = 2.8 \pm 0.3\text{‰}$, Figure 3, Table 1) and N^* was close to Redfield ($\text{N}^* = -0.2 \pm 0.8 \mu\text{M}$). As nitrate concentrations decreased into the euphotic zone, both $\delta^{15}\text{N}\text{-NO}_3$ and $\delta^{18}\text{O}\text{-NO}_3$ increased as a result of nitrate utilisation by phytoplankton (Figure 4 d and e).

The cooler ArW in the north of the Barents Sea had slightly lower nitrate concentrations of $10 \pm 1.1 \mu\text{M}$ (Table 1). NH_4^+ concentrations were high close to the seafloor in Hopen Trench but decreased with increasing latitude. There was no significant difference in $\delta^{15}\text{N}\text{-NO}_3$ or N^* between AW and ArW (ArW $\delta^{15}\text{N}\text{-NO}_3 = 5.1 \pm 0.1\text{‰}$, $\text{N}^* = -1.0 \pm 0.7 \mu\text{M}$, Table 1), suggesting these nutrients also originated from the Atlantic. In contrast, $\delta^{18}\text{O}\text{-NO}_3$ was 1.2‰ lower in ArW compared to AW (Figure 3b, Figure 4).

In the BSW, $\delta^{15}\text{N}\text{-NO}_3$, nitrate and N^* were comparable to AW and ArW ($\delta^{15}\text{N}\text{-NO}_3 = 5.1 \pm 0.4\text{‰}$, Nitrate = $10.4 \pm 1.2 \mu\text{M}$, $\text{N}^* = -1.1 \pm 1.1 \mu\text{M}$). BSW $\delta^{18}\text{O}\text{-NO}_3$ was $2.1 \pm 0.5\text{‰}$, lower than AW but higher than ArW, reflecting a mix between these two water masses.

4 Discussion

4.1 Origin of Atlantic Water supplied to the Barents Sea

The nutrient concentration within the AW that is advected into the Barents Sea is controlled by the relative contribution of nutrient-rich North Atlantic subpolar water and nutrient-poor subtropical waters that reach the Norwegian Sea together with the biological and physical transformations en-route. (Hatun et al., 2017; Rey, 2012; Johnson et al., 2013). Pre-bloom nutrient concentrations set the upper limit on seasonal primary productivity. Therefore, the origin of AW is important. Here we consider the contribution of subtropical and subpolar water to the AW sampled in the Barents Sea based on known nitrate isotope end members. We discuss the processes that the source waters are likely to have undergone and consider historical and future long-term trends in $\delta^{15}\text{N}\text{-NO}_3$.

Atlantic Water sampled in the Barents Sea during this study had a nitrate concentration of $11.8 \pm 1.8 \mu\text{M}$ (below 100m; Table 1) and a $\delta^{15}\text{N}\text{-NO}_3$ of $5.1 \pm 0.1\text{‰}$ (Table 1). This $\delta^{15}\text{N}\text{-NO}_3$ is comparable to subpolar gyre thermocline nitrate of $4.8 \pm 0.1\text{‰}$



(Peng et al., 2018), when compared over the same depth range (this study $>200\text{m}$ $5.0 \pm 0.1\text{‰}$). Subtropical sourced $\delta^{15}\text{N}\text{-NO}_3$ ($\sim 3.9\text{‰}$) is lower than subpolar $\delta^{15}\text{N}\text{-NO}_3$ (Van Oostende et al., 2017) as there are significant inputs from N_2 fixation producing a lower $\delta^{15}\text{N}$ nitrate source (Knapp et al., 2008). In comparison, isotopic measurements of subpolar nitrate and particulate nitrogen (PN) reveal the dominance of new production with local phytoplankton utilising nitrate sources from the subpolar thermocline (Peng et al., 2018; Van Oostende et al., 2017, this study), which is comparable to North Atlantic Deep Water (NADW) ($4.75\text{-}5\text{‰}$) (Marconi et al., 2015).

The $\delta^{18}\text{O}\text{-NO}_3$ of AW in the Barents Sea ($2.8 \pm 0.3\text{‰}$; Table 1) is high compared to NADW ($1.67\text{-}2.02\text{‰}$) (Marconi et al., 2015). The $\delta^{18}\text{O}\text{-NO}_3$ of NADW results from regeneration leading to $\delta^{18}\text{O}\text{-NO}_3$ close to the $\delta^{18}\text{O}\text{-H}_2\text{O}$ source plus 1.1‰ . Our characterisation of the AW nitrate that enters the Barents Sea reveals an enrichment in $\delta^{18}\text{O}\text{-NO}_3$ above a purely regenerated signal, which is also present in the subpolar gyre (Van Oostende et al., 2017). We measured a greater elevation in $\delta^{18}\text{O}\text{-NO}_3$ (by 0.8‰) than $\delta^{15}\text{N}\text{-NO}_3$ (0.1‰) compared to NADW. An elevation in $\delta^{18}\text{O}\text{-NO}_3$ relative to deep water values from the North Atlantic suggests that partial nitrate assimilation followed by nitrification is occurring on transit from the North Atlantic and into the Barents Sea (Van Oostende et al., 2017; Peng et al., 2018). This process leads to the regeneration of nitrate which decreases $\delta^{15}\text{N}\text{-NO}_3$ to a greater extent than $\delta^{18}\text{O}\text{-NO}_3$.

In the North Atlantic, low $\delta^{15}\text{N}\text{-NO}_3$ is associated with high salinity of the subtropical gyre (Knapp et al., 2008). The salinity of the AW supplied to the Barents Sea has increased in recent years (Barton et al., 2018; Oziel et al., 2016). We suggest that continued increases in salinity and the associated decrease in nitrate supply (Rey, 2012), has the potential to decrease $\delta^{15}\text{N}\text{-NO}_3$ of Arctic nitrate supply albeit to a small degree. Based upon the salinity- $\delta^{15}\text{N}\text{-NO}_3$ relationship established in the wider Atlantic (Marconi et al., 2015; Schlitzer et al., 2018), the $0.05\text{-}0.1$ psu change in salinity between the periods 1985-2005 and 2005-2016 (Barton et al., 2018) implies that there has been a $0.06\text{-}0.13$ decrease in $\delta^{15}\text{N}\text{-NO}_3$ (Pearson corr. = -0.82 , $\text{df} = 12$, $p\text{-value} = 0.0003$).

4.2 Nitrate utilisation and limitation in the Barents Sea

In July 2017, nitrate was depleted in the euphotic zone, coinciding with an increase in both $\delta^{15}\text{N}\text{-NO}_3$ and $\delta^{18}\text{O}\text{-NO}_3$. The seasonal uptake of nitrate by phytoplankton fractionates $\delta^{15}\text{N}\text{-NO}_3$ and $\delta^{18}\text{O}\text{-NO}_3$ with an isotope effect (ϵ), close to 5‰ (Sigman et al., 2009b). This relationship can determine the relative importance of algal uptake versus other processes such as dilution and regeneration (DiFiore et al., 2006; Rafter et al., 2012). Here we find that in the Arctic, ϵ is often muted in surface waters through dilution with nitrate-deplete freshwater.

The southern Barents Sea remains ice-free all year round and away from the Norwegian Coastal Current, the near surface salinity remains high. During the spring and summer months a warm surface mixed layer is established which triggers



phytoplankton growth. As nitrate decreases, both $\delta^{15}\text{N}\text{-NO}_3$ and $\delta^{18}\text{O}\text{-NO}_3$ increase and algal uptake of nitrate is the dominant N cycling process occurring in the euphotic zone and is fuelled by new production (nitrate). We estimate a $\delta^{15}\text{N}\text{-NO}_3$ uptake fractionation of 4.7–4.9‰ (Figure 5a and c), with isotopic data following a trend for Rayleigh fractionation, or a closed system (Mariotti et al., 1981). This finding is anticipated since strong stratification isolated the euphotic zone from deeper waters during the time of sampling.

In the northern Barents Sea, $\delta^{15}\text{N}\text{-NO}_3$ and $\delta^{18}\text{O}\text{-NO}_3$ increase as nitrate decreases in the euphotic zone. These waters are cooler and fresher and are likely to have undergone at least one seasonal cycle on the Barents Sea shelf, where there is evidence for nutrient regeneration (Section 4.3). We find a muted uptake fractionation in this region of 1.8‰ which is likely due to dilution of the nitrate concentration by fresh, nutrient depleted surface water (Figure 5a and c).

The stable isotopic signal recorded in the Arctic marine food web is primarily dependent upon the particulate organic material produced by phytoplankton, representing the base of the food web, whose ^{15}N is controlled by the dissolved nutrient source. With knowledge of the mechanism behind isotopic fractionation during nitrate uptake, and if nitrate uptake is the primary N cycling process occurring in the euphotic zone, then $\delta^{15}\text{N}\text{-PN}$ may be predicted.

The JR16006 cruise was conducted during summer when the southern Barents Sea was thermally-stratified. Further north, sea-ice melt had established a fresh surface mixed layer resulting in salinity-driven stratification. Throughout the Barents Sea, particulate organic matter load was highest in the euphotic zone (average of $31.7 \pm 14.7 \mu\text{g L}^{-1}$), and decreased to $9.5 \pm 3.4 \mu\text{g L}^{-1}$ below 70 m. We found that $\delta^{15}\text{N}\text{-PN}$ largely followed nitrate concentration, falling close to the trend for the integrated product of N uptake (Figure 6a). In areas where there was still nitrate available to phytoplankton, $\delta^{15}\text{N}\text{-PN}$ was lower representing the preferential consumption of the lighter isotope. $\delta^{15}\text{N}\text{-PN}$ increased to $\sim 5\text{‰}$ as the nitrate concentration approached zero, matching the AW source. Using this information, we predict how the $\delta^{15}\text{N}\text{-PN}$ is likely to change in the euphotic layer following Rayleigh fractionation systematics ($\delta^{15}\text{N}_{\text{mod}}$) from the nutrient sources of AW and ArW.

$$\delta^{15}\text{N}_{\text{mod}} = \delta^{15}\text{N}_{\text{initial}} + \varepsilon \left(\frac{u}{(1-u)} \right) \times \ln(u) \quad (1)$$

where $u = \text{NO}_3_{\text{observed}} / \text{NO}_3_{\text{initial}}$, $\delta^{15}\text{N}_{\text{initial}} = 5.1\text{‰}$, $\varepsilon = 5\text{‰}$, and $\text{NO}_3_{\text{initial}} = 11.8 \mu\text{M}$. We find the spatial trends are captured in the modelled data with the highest modelled $\delta^{15}\text{N}\text{-PN}$ where the concentrations are the lowest and vice versa (Figure 6c). Deviations from the trend, representing a lower isotopic effect, are in lower temperature samples from ArW (Figure 6c). At these locations the upper euphotic zone is salinity stratified and polar surface water dilutes the nitrate concentration. If the ArW samples are corrected to the lower isotope effect of 1.8‰ and nitrate concentration ($10 \mu\text{M}$), as predicted from our $\delta^{15}\text{N}\text{-NO}_3$ data, we find a Pearson's correlation, $r=0.87$, $\text{df}=33$, $p=0$ (Figure 6c).



285 These results support the finding that nitrate from the Atlantic is the primary source of nutrients to phytoplankton in surface waters and that the organic matter in the euphotic zone is principally autotrophic. When there is still nitrate readily available in surface waters the phytoplankton preferentially take up ^{14}N and a lower $\delta^{15}\text{N}$ is expressed in particulate N. As there is full utilisation of nutrients over the growing season, we suggest that the integrated source of organic matter to the sediments and food web is $\sim 5\text{‰}$ throughout the Barents Sea.

290

In order to investigate any seasonal changes in the organic matter source in surface waters we consider the measurements of nitrate, PON and their isotopic ratios on the repeat transects across the BSO (Figure 7). Nitrate concentrations were highest in March, from replenishment over winter months. Nitrite remained below $0.25\text{ }\mu\text{M}$ throughout all seasons. The nitrite concentrations were lowest in March, suggesting that the intermediate products, NH_4^+ and NO_2^- had been nitrified to nitrate
 295 (Figure 7). The highest nitrite concentrations were sampled in June and August, during or following the spring bloom, and remained high into November. $\delta^{15}\text{N}\text{-NO}_3$ and $\delta^{18}\text{O}\text{-NO}_3$ showed uptake driven changes in the nutrients during June and August, whereby the low nitrate concentrations coincided with heavy isotope values from the preferential consumption of the lighter isotope (Figure 5d, Figure 7d, e).

300 PON concentrations were low in winter and markedly increased in June and August (Figure 7c). $\delta^{15}\text{N}\text{-PN}$ values from August, November and March were relatively constant at around 5‰ . In June $\delta^{15}\text{N}\text{-PN}$ decreased as the lighter isotope is preferentially consumed by phytoplankton (Figure 7f). The relatively constant value of 5‰ for the rest of the annual cycle reflected the AW source value of $\sim 5\text{‰}$, suggesting that there is limited new production occurring over the winter months and that $\delta^{15}\text{N}\text{-PN}$ represents the integrated product of nitrate uptake from the previous growing season.

305 4.3 Nitrogen cycling processes occurring in the Barents Sea

4.3.1 Nitrification

As inflowing AW cools and freshens across the Barents Sea, $\delta^{18}\text{O}\text{-NO}_3$ decreases from its AW source value of $2.8 \pm 0.3\text{‰}$ to $1.6 \pm 0.3\text{‰}$ (Figure 3). This decline represents N recycling and nitrification. A range in nitrified nitrate values of -1.5 to 1.3‰ have been reported from nitrifier cocultures (Buchwald et al., 2012). Previous field and modelling studies have used a nitrifying
 310 $\delta^{18}\text{O}$ value of 1.1‰ plus $\delta^{18}\text{O}_{\text{H}_2\text{O}}$ (Granger et al., 2013; Sigman et al., 2009b). As nitrate is regenerated, newly nitrified nitrate tracks the $\delta^{18}\text{O}$ of seawater, which, in the Barents Sea is $\sim 0.2\text{‰}$ (Schlitzer et al., 2018), therefore as the proportion of regenerated nitrate increases, $\delta^{18}\text{O}$ will decrease towards $\sim 1.3\text{‰}$.

The recycling of nitrate in-situ is a common feature on Arctic shelves, evidenced using nitrate isotopes on the West Siberian
 315 Shelf (Fripiat et al., 2018) and the Canadian Shelf (Granger et al., 2018). In these regions, $\delta^{18}\text{O}\text{-NO}_3$ tracks $\delta^{18}\text{O}\text{-H}_2\text{O}$ showing



the importance of N recycling in sustaining the N-limited primary production the following season. As we have characterised the Atlantic source $\delta^{18}\text{O}-\text{NO}_3$, we can track the extent of nitrification across the Barents Sea to give an estimate of the proportion of regenerated nitrate on the Barents Sea shelf (Granger et al., 2013).

320

$$\frac{\text{NO}_3_{\text{reg}}}{\text{NO}_3_{\text{tot}}} = \frac{(\delta^{18}\text{O}_{\text{meas}} - \delta^{18}\text{O}_{\text{AW}})}{(\delta^{18}\text{O}_{\text{reg}} - \delta^{18}\text{O}_{\text{AW}})} \quad (2)$$

Where $\delta^{18}\text{O}_{\text{meas}}$ = measured $\delta^{18}\text{O}-\text{NO}_3$, $\delta^{18}\text{O}_{\text{AW}}$ =2.8‰, $\delta^{18}\text{O}_{\text{reg}}$ =1.3‰ and $\text{NO}_3_{\text{reg}}/\text{NO}_3_{\text{tot}}$ = proportion of regenerated nitrate. In Figure 3d the proportion of nitrate regenerated follows the trend of $\delta^{18}\text{O}-\text{NO}_3$. The proportion of nitrate regenerated remains relatively unmodified between the BSO and the Spitzbergen-Grand Banks Sill (the approximate location of the Polar Front), north of which the proportion of regenerated nitrate increases from <10% in the south, to >80% in ArW. The nutrient concentration of ArW is coupled to winter mixing, driven by atmospheric cooling and brine release during sea ice formation. The ArW experiences nutrient regeneration and nitrification over winter which works to decrease $\delta^{18}\text{O}-\text{NO}_3$. As the shelf waters on the Barents Sea cool and freshen, the nitrate inventory is also replenished from the nitrification of ammonium which is supplied to the water column from sediments. The resupply and mixing of nutrients from the sediments is an important component in replenishing the N inventory.

4.3.2 Nitrogen resupply from sediments

Organic matter produced in surface waters will ultimately be regenerated in the water column or sink to the seafloor. The release of NH_4^+ from relatively shallow Arctic sediments has been noted in previous work, where the organic rich shelf sediments provide a source of NH_4^+ to the water column (Brown et al., 2015). Studies from the Chukchi Sea suggest there are annually varying rates of nitrification, with much higher rates in winter (Christman et al., 2011). This suggests that there is a build-up of NH_4^+ in summer and NH_4^+ concentrations decrease into winter as nitrification rates exceed NH_4^+ release. We found NH_4^+ was enhanced over the Spitzbergen Bank and in the Hopen Trough, with the highest concentrations close to the sediment rather than the euphotic zone, indicating that the sediments are releasing NH_4^+ to the water column.

NH_4^+ is generated in sediments by the ammonification of organic material and can be released by diffusive and non-diffusive fluxes (Granger et al., 2011). Previous studies have suggested that the NH_4^+ produced during ammonification should be similar to the organic matter source, but that there is a large isotopic effect (~14‰) associated with the nitrification of NH_4^+ to NO_2^- (Casciotti et al., 2003). Over the course of the season, once all NH_4^+ that has been released from the sediments and oxidised, $\delta^{15}\text{N}-\text{NO}_3$ should reflect the N source (in this study: $5.1 \pm 0.1\text{‰}$). There were a few samples with low (<4.8 ‰) $\delta^{15}\text{N}-\text{NO}_3$ on the flanks of the Spitzbergen Bank, near the seabed at the head of Hopen Trench and over the SB-GB sill, which may be associated with partial N recycling processes and the retention of ^{15}N in NH_4^+ (Casciotti et al., 2003). However, nitrate $\delta^{15}\text{N}$



below the nitricline was relatively homogenous across our sampled transect, reflecting the AW source value of $5.1 \pm 0.1\text{‰}$ (Table 1).

In the western Arctic, Bering Sea and East Siberian Sea the release of NH_4^+ from sediments leads to a decrease in $\delta^{18}\text{O}\text{-NO}_3$ and an enrichment in $\delta^{15}\text{N}\text{-NO}_3$ over the timescales of water mass transit (Fripiat et al., 2018; Granger et al., 2018). In these regions, remineralisation is greater than nitrification, therefore NH_4^+ diffuses out of the sediments which is higher in ^{15}N , as low $\delta^{15}\text{N}$ nitrified nitrate is lost to benthic denitrification in sediments (Granger et al., 2011). This process enriches NH_4^+ in ^{15}N , a signature which is subsequently imparted on the overlying water column when NH_4^+ is released from the sediments and oxidized by nitrifiers (Brown et al., 2015). These trends are combined with concomitant decreases in N^* , demonstrating the prevalence of benthic denitrification on Arctic shelves, where $\delta^{15}\text{N}$ increases from $\sim 6.5\text{‰}$ at the Bering Strait (Brown et al., 2015; Lehmann et al., 2007) to 8‰ on the Canadian and Siberian Shelves (Fripiat et al., 2018; Granger et al., 2018).

If coupled partial nitrification-denitrification was occurring in the Barents Sea sediments, there should be an observed increase in $\delta^{15}\text{N}\text{-NO}_3$ with the decrease in $\delta^{18}\text{O}\text{-NO}_3$ through NH_4^+ release and nitrification. This is not evident in our dataset. (Table 1, Figure 4). Instead, we found no clear increase in $\delta^{15}\text{N}\text{-NO}_3$ or decrease in N^* from the AW entering the shelf, to ArW and BSW further north and east (Table 1). This finding suggests that either the process of NH_4^+ release from the sediments is insignificant to the water column nitrate inventory, or that in contrast to the Canadian and Siberian Shelves, the layer of low oxygen (and thus denitrification), is separated from the layer of ammonification and NH_4^+ release from sediments. The high NH_4^+ , which exceeds 25% of the dissolved inorganic N inventory at the base of some profiles, suggests that NH_4^+ was accumulating in the water column at the time of our study.

In the Barents Sea, the shallow banks and slopes (e.g. Spitzbergen Bank) experience strong tidal and frontal currents which induce significant mixing (Sundfjord et al., 2007), and in shallower water winter convection is able to overturn the whole water column, processes that are able to remobilise and increase the oxygenation of surficial sediments. In the shallow regions we would therefore predict a deeper depth of denitrification within sediments and the faster release of NH_4^+ to the water column, largely by advective rather than diffusive fluxes.

The contrasting findings between this study in the Barents Sea and other Arctic shelves may result from a number of factors. The Pacific inflow supplies the much shallower Chukchi and Beaufort shelves ($< 60\text{m}$), with higher concentrations of macronutrients (Granger et al., 2013). In contrast, the Atlantic inflow to the Barents Sea provides lower concentrations of macronutrients to a deeper shelf ($> 100\text{m}$), therefore the organic load to sediments and thus benthic denitrification is expected to be lower (Chang and Devol, 2009).



5 Summary

We show that nitrogen availability in the Barents Sea is supported through AW supply and the efficient replenishment of nutrients through seasonal cycling processes. By the end of the growing season, all nitrate is consumed in surface waters and the $\delta^{15}\text{N}$ of PON reflects the AW source. The N inventory is dependent on the NH_4^+ release from sediments and nitrification. In contrast to other Arctic shelf regions, we find no evidence for benthic denitrification communicating with the water column (and no loss of N relative to P). Our results indicate that although nutrients are regenerated in the western Barents Sea, $\delta^{15}\text{N}$ - NO_3 does not increase, suggesting that $\delta^{15}\text{N}$ - NO_3 supplied to Arctic Intermediate Water may be comparable to the AW source values. Our findings suggest $\delta^{15}\text{N}$ - NO_3 is unmodified in transit through the western Barents Sea. Additional samples collected in the Eastern Barents Sea and at the primary export gateway (St Anna Trough) are needed to confirm this.

Previous work has suggested that increasing NPP on Arctic shelves would increase organic matter supply to sediments and thus increase sedimentary denitrification rates (Arrigo and van Dijken, 2015). As N is the primary limiting nutrient to Arctic phytoplankton (Mills et al., 2018), this would have downstream consequences to NPP in the central Arctic basin. Our findings suggest that increased organic matter supply to sediments through increasing NPP is unlikely to exacerbate N limitation in the Barents Sea. We conclude that changes to productivity across the Barents Sea are unlikely to have a significant impact on the nutrient inventory in the Barents Sea and thus the magnitude of nutrients transported to the Arctic waters, which is more dependent on variability in the Atlantic Water inflow.

400



References

- Altabet, M. A., and Francois, R.: Sedimentary nitrogen isotopic ratio as a recorder for surface ocean nitrate utilization, *Global Biogeochemical Cycles*, 8, 103-116, 10.1029/93gb03396, 1994.
- Altieri, K. E., Fawcett, S. E., Peters, A. J., Sigman, D. M., and Hastings, M. G.: Marine biogenic source of atmospheric organic nitrogen in the subtropical North Atlantic, *Proceedings of the National Academy of Sciences of the United States of America*, 113, 925-930, 10.1073/pnas.1516847113, 2016.
- Arrigo, K. R., and van Dijken, G. L.: Continued increases in Arctic Ocean primary production, *Progress in Oceanography*, 136, 60-70, 10.1016/j.pocean.2015.05.002, 2015.
- Arthun, M., Ingvaldsen, R. B., Smedsrud, L. H., and Schrum, C.: Dense water formation and circulation in the Barents Sea, *Deep-Sea Research Part I-Oceanographic Research Papers*, 58, 801-817, 10.1016/j.dsr.2011.06.001, 2011.
- Arthun, M., Eldevik, T., Smedsrud, L. H., Skagseth, O., and Ingvaldsen, R. B.: Quantifying the Influence of Atlantic Heat on Barents Sea Ice Variability and Retreat, *Journal of Climate*, 25, 4736-4743, 10.1175/jcli-d-11-00466.1, 2012.
- Barton, B. I., Lenn, Y. D., and Lique, C.: Observed Atlantification of the Barents Sea Causes the Polar Front to Limit the Expansion of Winter Sea Ice, *Journal of Physical Oceanography*, 48, 1849-1866, 10.1175/jpo-d-18-0003.1, 2018.
- Brown, Z. W., Casciotti, K. L., Pickart, R. S., Swift, J. H., and Arrigo, K. R.: Aspects of the marine nitrogen cycle of the Chukchi Sea shelf and Canada Basin, *Deep-Sea Research Part II-Topical Studies in Oceanography*, 118, 73-87, 10.1016/j.dsr2.2015.02.009, 2015.
- Buchwald, C., Santoro, A. E., McIlvin, M. R., and Casciotti, K. L.: Oxygen isotopic composition of nitrate and nitrite produced by nitrifying cocultures and natural marine assemblages, *Limnology and Oceanography*, 57, 1361-1375, 10.4319/lo.2012.57.5.1361, 2012.
- Casciotti, K. L., Sigman, D. M., Hastings, M. G., Bohlke, J. K., and Hilkert, A.: Measurement of the oxygen isotopic composition of nitrate in seawater and freshwater using the denitrifier method, *Analytical Chemistry*, 74, 4905-4912, 10.1021/ac020113w, 2002.
- Casciotti, K. L., Sigman, D. M., and Ward, B. B.: Linking diversity and stable isotope fractionation in ammonia-oxidizing bacteria, *Geomicrobiology Journal*, 20, 335-353, 10.1080/01490450303895, 2003.
- Casciotti, K. L., Boehlke, J. K., McIlvin, M. R., Mroczkowski, S. J., and Hannon, J. E.: Oxygen isotopes in nitrite: Analysis, calibration, and equilibration, *Analytical Chemistry*, 79, 2427-2436, 10.1021/ac061598h, 2007.
- Chang, B. X., and Devol, A. H.: Seasonal and spatial patterns of sedimentary denitrification rates in the Chukchi sea, *Deep-Sea Research Part II-Topical Studies in Oceanography*, 56, 1339-1350, 10.1016/j.dsr2.2008.10.024, 2009.
- Christman, G. D., Cottrell, M. T., Popp, B. N., Gier, E., and Kirchman, D. L.: Abundance, Diversity, and Activity of Ammonia-Oxidizing Prokaryotes in the Coastal Arctic Ocean in Summer and Winter, *Applied and Environmental Microbiology*, 77, 2026-2034, 10.1128/aem.01907-10, 2011.
- Codispoti, L. A., Kelly, V., Thessen, A., Matrai, P., Suttles, S., Hill, V., Steele, M., and Light, B.: Synthesis of primary production in the Arctic Ocean: III. Nitrate and phosphate based estimates of net community production, *Progress in Oceanography*, 110, 126-150, 10.1016/j.pocean.2012.11.006, 2013.
- Coupe, P., Ruiz-Pino, D., Sicre, M. A., Chen, J. F., Lee, S. H., Schiffrine, N., Li, H. L., and Gascard, J. C.: The impact of freshening on phytoplankton production in the Pacific Arctic Ocean, *Progress in Oceanography*, 131, 113-125, 10.1016/j.pocean.2014.12.003, 2015.
- Dalpadado, P., Arrigo, K. R., Hjollo, S. S., Rey, F., Ingvaldsen, R. B., Sperfeld, E., van Dijken, G. L., Stige, L. C., Olsen, A., and Ottersen, G.: Productivity in the Barents Sea - Response to Recent Climate Variability, *Plos One*, 9, 15, 10.1371/journal.pone.0095273, 2014.
- DiFiore, P. J., Sigman, D. M., Trull, T. W., Lourey, M. J., Karsh, K., Cane, G., and Ho, R.: Nitrogen isotope constraints on subantarctic biogeochemistry, *Journal of Geophysical Research-Oceans*, 111, 10.1029/2005jc003216, 2006.



- DiFiore, P. J., Sigman, D. M., and Dunbar, R. B.: Upper ocean nitrogen fluxes in the Polar Antarctic Zone: Constraints from the nitrogen and oxygen isotopes of nitrate, *Geochemistry Geophysics Geosystems*, 10, Q11016
460 10.1029/2009gc002468, 2009.
- Dumont, Brand, and Hopkins: CTD data from NERC Changing Arctic Ocean Cruise JR16006 on the RRS James Clark Ross, June-August 2017., British Oceanographic Data Centre, National Oceanography Centre, NERC, UK. , 2019.
- 465 Ellingsen, I., Slagstad, D., and Sundfjord, A.: Modification of water masses in the Barents Sea and its coupling to ice dynamics: a model study, *Ocean Dynamics*, 59, 1095-1108, 10.1007/s10236-009-0230-5, 2009.
- Friedland, K. D., and Todd, C. D.: Changes in Northwest Atlantic Arctic and Subarctic conditions and the growth response of Atlantic salmon, *Polar Biology*, 35, 593-609, 10.1007/s00300-011-1105-z, 2012.
- 470 Fripiat, F., Declercq, M., Sapart, C. J., Anderson, L. G., Bruechert, V., Deman, F., Fonseca-Batista, D., Humborg, C., Roukaerts, A., Semiletov, I. P., and Dehairs, F.: Influence of the bordering shelves on nutrient distribution in the Arctic halocline inferred from water column nitrate isotopes, *Limnology and Oceanography*, 63, 2154-2170, 10.1002/lno.10930, 2018.
- 475 Granger, J., Sigman, D. M., Needoba, J. A., and Harrison, P. J.: Coupled nitrogen and oxygen isotope fractionation of nitrate during assimilation by cultures of marine phytoplankton, *Limnology and Oceanography*, 49, 1763-1773, 2004.
- Granger, J., and Sigman, D. M.: Removal of nitrite with sulfamic acid for nitrate N and O isotope analysis with the denitrifier method, *Rapid Communications in Mass Spectrometry*, 23, 3753-3762, 10.1002/rcm.4307, 2009.
- 480 Granger, J., Prokopenko, M. G., Sigman, D. M., Mordy, C. W., Morse, Z. M., Morales, L. V., Sambrotto, R. N., and Plessen, B.: Coupled nitrification-denitrification in sediment of the eastern Bering Sea shelf leads to N-15 enrichment of fixed N in shelf waters, *Journal of Geophysical Research-Oceans*, 116, 10.1029/2010jc006751, 2011.
- 485 Granger, J., Prokopenko, M. G., Mordy, C. W., and Sigman, D. M.: The proportion of remineralized nitrate on the ice-covered eastern Bering Sea shelf evidenced from the oxygen isotope ratio of nitrate, *Global Biogeochemical Cycles*, 27, 962-971, 10.1002/gbc.20075, 2013.
- Granger, J., Sigman, D. M., Gagnon, J., Tremblay, J. E., and Mucci, A.: On the Properties of the Arctic Halocline and Deep Water Masses of the Canada Basin from Nitrate Isotope Ratios, *Journal of Geophysical Research-Oceans*, 123, 5443-5458, 10.1029/2018jc014110, 2018.
- 490 Hatun, H., Azetsu-Scott, K., Somavilla, R., Rey, F., Johnson, C., Mathis, M., Mikolajewicz, U., CoupeI, P., Tremblay, J. E., Hartman, S., Pacariz, S. V., Salter, I., and Olafsson, J.: The subpolar gyre regulates silicate concentrations in the North Atlantic, *Scientific Reports*, 7, 10.1038/s41598-017-14837-4, 2017.
- 495 Huang, J. B., Zhang, X. D., Zhang, Q. Y., Lin, Y. L., Hao, M. J., Luo, Y., Zhao, Z. C., Yao, Y., Chen, X., Wang, L., Nie, S. P., Yin, Y. Z., Xu, Y., and Zhang, J. S.: Recently amplified arctic warming has contributed to a continual global warming trend, *Nature Climate Change*, 7, 875-+, 10.1038/s41558-017-0009-5, 2017.
- Johnson, C., Inall, M., and Hakkinen, S.: Declining nutrient concentrations in the northeast Atlantic as a result of a weakening Subpolar Gyre, *Deep-Sea Research Part I-Oceanographic Research Papers*, 82, 95-107, 10.1016/j.dsr.2013.08.007, 2013.
- 500 Kemeny, P. C., Weigand, M. A., Zhang, R., Carter, B. R., Karsh, K. L., Fawcett, S. E., and Sigman, D. M.: Enzyme-level interconversion of nitrate and nitrite in the fall mixed layer of the Antarctic Ocean, *Global Biogeochemical Cycles*, 30, 1069-1085, 10.1002/2015gb005350, 2016.
- 505 Knapp, A. N., DiFiore, P. J., Deutsch, C., Sigman, D. M., and Lipschultz, F.: Nitrate isotopic composition between Bermuda and Puerto Rico: Implications for N(2) fixation in the Atlantic Ocean, *Global Biogeochemical Cycles*, 22, Gb3014 10.1029/2007gb003107, 2008.
- 510 Lehmann, M. F., Sigman, D. M., McCorkle, D. C., Granger, J., Hoffmann, S., Cane, G., and Brunelle, B. G.: The distribution of nitrate N-15/N-14 in marine sediments and the impact of benthic nitrogen loss on the isotopic composition of oceanic nitrate, *Geochimica Et Cosmochimica Acta*, 71, 5384-5404, 10.1016/j.gca.2007.07.025, 2007.



- 515 Lewis, K. M., Van Dijken, G. L., and Arrigo, K. R.: Changes in phytoplankton concentration now drive increased Arctic Ocean primary production, 369, 198-202, 10.1126/science.aay8380, 2020.
- Lind, S., Ingvaldsen, R. B., and Furevik, T.: Arctic layer salinity controls heat loss from deep Atlantic layer in seasonally ice-covered areas of the Barents Sea, *Geophysical Research Letters*, 43, 5233-5242, 10.1002/2016gl068421, 2016.
- 520 Lind, S., Ingvaldsen, R. B., and Furevik, T.: Arctic warming hotspot in the northern Barents Sea linked to declining sea-ice import, *Nature Climate Change*, 8, 634-+, 10.1038/s41558-018-0205-y, 2018.
- Marconi, D., Weigand, M. A., Rafter, P. A., McIlvin, M. R., Forbes, M., Casciotti, K. L., and Sigman, D. M.: Nitrate isotope distributions on the US GEOTRACES North Atlantic cross-basin section: Signals of polar nitrate sources and low latitude nitrogen cycling, *Marine Chemistry*, 177, 143-156, 10.1016/j.marchem.2015.06.007, 2015.
- 525 Marconi, D., Sigman, D. M., Casciotti, K. L., Campbell, E. C., Weigand, M. A., Fawcett, S. E., Knapp, A. N., Rafter, P. A., Ward, B. B., and Haug, G. H.: Tropical Dominance of N₂ Fixation in the North Atlantic Ocean, *Global Biogeochemical Cycles*, 31, 1608-1623, 10.1002/2016gb005613, 2017.
- 530 Mariotti, A., Germon, J. C., Hubert, P., Kaiser, P., Letolle, R., Tardieux, A., and Tardieux, P.: Experimental - determination of nitrogen kinetic isotope fractionation - some principles - Illustration for the denitrification and nitrification processes, *Plant and Soil*, 62, 413-430, 10.1007/bf02374138, 1981.
- 535 Mills, M. M., Brown, Z. W., Laney, S. R., Ortega-Retuerta, E., Lowry, K. E., van Dijken, G. L., and Arrigo, K. R.: Nitrogen Limitation of the Summer Phytoplankton and Heterotrophic Prokaryote Communities in the Chukchi Sea, *Frontiers in Marine Science*, 5, 10.3389/fmars.2018.00362, 2018.
- Notz, D., and Stroeve, J.: Observed Arctic sea-ice loss directly follows anthropogenic CO₂ emission, *Science*, 354, 747-750, 10.1126/science.aag2345, 2016.
- 540 Onarheim, I. H., Eldevik, T., Arthun, M., Ingvaldsen, R. B., and Smedsrud, L. H.: Skillful prediction of Barents Sea ice cover, *Geophysical Research Letters*, 42, 5364-5371, 10.1002/2015gl064359, 2015.
- 545 Onarheim, I. H., and Arthun, M.: Toward an ice-free Barents Sea, *Geophysical Research Letters*, 44, 8387-8395, 10.1002/2017gl074304, 2017.
- Oziel, L., Sirven, J., and Gascard, J. C.: The Barents Sea frontal zones and water masses variability (1980-2011), *Ocean Science*, 12, 169-184, 10.5194/os-12-169-2016, 2016.
- 550 Oziel, L., Baudena, A., Ardyna, M., Massicotte, P., Randelhoff, A., Sallée, J. B., Ingvaldsen, R. B., Devred, E., and Babin, M.: Faster Atlantic currents drive poleward expansion of temperate phytoplankton in the Arctic Ocean. *Nat Commun* 11, 1705 (2020). <https://doi.org/10.1038/s41467-020-15485-5>. *Nature Communications*, 2020.
- 555 Peng, X. F., Fawcett, S. E., van Oostende, N., Wolf, M. J., Marconi, D., Sigman, D. M., and Ward, B. B.: Nitrogen uptake and nitrification in the subarctic North Atlantic Ocean, *Limnology and Oceanography*, 63, 1462-1487, 10.1002/lno.10784, 2018.
- Rafter, P. A., Sigman, D. M., Charles, C. D., Kaiser, J., and Haug, G. H.: Subsurface tropical Pacific nitrogen isotopic composition of nitrate: Biogeochemical signals and their transport, *Global Biogeochemical Cycles*, 26, 10.1029/2010gb003979, 2012.
- 560 Rafter, P. A., DiFiore, P. J., and Sigman, D. M.: Coupled nitrate nitrogen and oxygen isotopes and organic matter remineralization in the Southern and Pacific Oceans, *Journal of Geophysical Research-Oceans*, 118, 4781-4794, 10.1002/jgrc.20316, 2013.
- Rey, F.: Declining silicate concentrations in the Norwegian and Barents Seas, *Ices Journal of Marine Science*, 69, 208-212, 10.1093/icesjms/fss007, 2012.
- 565 Rudels, B., Anderson, L. G., and Jones, E. P.: Formation and evolution of the surface mixed layer and halocline of the Arctic Ocean, *Journal of Geophysical Research-Oceans*, 101, 8807-8821, 10.1029/96jc00143, 1996.



- 570 Schlitzer, R., Anderson, R. F., Dodas, E. M., Lohan, M., Geibere, W., Tagliabue, A., Bowie, A., Jeandel, C., Maldonado, M. T., Landing, W. M., Cockwell, D., Abadie, C., Abouchami, W., Achterberg, E. P., Agather, A., Aguiar-Islas, A., van Aken, H. M., Andersen, M., Archer, C., Auro, M., de Baar, H. J., Baars, O., Baker, A. R., Bakker, K., Basak, C., Baskaran, M., Bates, N. R., Bauch, D., van Beek, P., Behrens, M. K., Black, E., Bluhm, K., Bopp, L., Bouman, H., Bowman, K., Bown, J., Boyd, P., Boye, M., Boyle, E. A., Branellec, P., Bridgestock, L., Brissebrat, G., Browning, T., Bruland, K. W., Brumsack, H. J., Brzezinski, M., Buck, C. S., Buck, K. N., Buesseler, K., Bull, A., Butler, E., Cai, P., Mor, P. C., Cardinal, D., Carlson, C., Carrasco, G., Casacuberta, N., Casciotti, K. L., Castrillejo, M., Chamizo, E., Chance, R., Charette, M. A., Chaves, J. E., Cheng, H., Chever, F., Christl, M., Church, T. M., Closset, I., Colman, A., Conway, T. M., Cossa, D., Croot, P., Cullen, J. T., Cutter, G. A., Daniels, C., Dehairs, F., Deng, F. F., Dieu, H. T., Duggan, B., Dulaquais, G., Dumousseaud, C., Echegoyen-Sanz, Y., Edwards, R. L., Ellwood, M., Fahrbach, E., Fitzsimmons, J. N., Flegal, A. R., Fleisher, M. Q., van de Flierdt, T., Frank, M., Friedrich, J., Fripiat, F., Frollje, H., Galer, S. J. G., Gamo, T., Ganeshram, R. S., Garcia-Orellana, J., Garcia-Solsona, E., Gault-Ringold, M., George, E., Gerringa, L. J. A., Gilbert, M., Godoy, J. M., Goldstein, S. L., Gonzalez, S. R., Grissom, K., Hammerschmidt, C., Hartman, A., Hassler, C. S., Hathorne, E. C., Hatta, M., Hawco, N., Hayes, C. T., Heimbürger, L. E., Helgoe, J., Heller, M., Henderson, G. M., Henderson, P. B., van Heuven, S., Ho, P., Horner, T. J., Hsieh, Y. T., Huang, K. F., Humphreys, M. P., Isshiki, K., Jacquot, J. E., Janssen, D. J., Jenkins, W. J., John, S., Jones, E. M., Jones, J. L., Kadko, D. C., Kayser, R., Kenna, T. C., Khondoker, R., Kim, T., Kipp, L., Klar, J. K., Klunder, M., Kretschmer, S., Kumamoto, Y., Laan, P., Labatut, M., Lacan, F., Lam, P. J., Lambelet, M., Lamborg, C. H., Le Moigne, F. A. C., Le Roy, E., Lechtenfeld, O. J., Lee, J. M., Lherminier, P., Little, S., Lopez-Lora, M., Lu, Y. B., Masque, P., Mawji, E., McClain, C. R., Measures, C., Mehic, S., Menzel Barraqueta, J. L., van der Merwe, P., Middag, R., Mieruch, S., Milne, A., Minami, T., Moffett, J. W., Moncoiffe, G., Moore, W. S., Morris, P. J., Morton, P. L., Nakaguchi, Y., Nakayama, N., Niedermiller, J., Nishioka, J., Nishiuchi, A., Noble, A., Obata, H., Ober, S., Ohnemus, D. C., van Ooijen, J., O'Sullivan, J., Owens, S., Pahnke, K., Paul, M., Pavia, F., Pena, L. D., Petersh, B., Planchon, F., Planquette, H., Pradoux, C., Puigcorbe, V., Quay, P., Queroue, F., Radic, A., Rauschenberg, S., Rehkamper, M., Rember, R., Remenyi, T., Resing, J. A., Rickli, J., Rigaud, S., Rijkenberg, M. J. A., Rintoul, S., Robinson, L. F., Roca-Marti, M., Rodellas, V., Roeske, T., Rolison, J. M., Rosenberg, M., Roshan, S., van der Loeff, M. M. R., Ryabenko, E., Saito, M. A., Salt, L. A., Sanial, V., Sarthou, G., Schallenberg, C., Schauer, U., Scher, H., Schlosser, C., Schnetger, B., Scott, P., Sedwick, P. N., Semiletov, I., Shelley, R., Sherrell, R. M., Shiller, A. M., Sigman, D. M., Singh, S. K., Slagter, H. A., Slater, E., Smethie, W. M., Snaith, H., Sohrin, Y., Sohst, B., Sonke, J. E., Speich, S., Steinfeldt, R., Stewart, G., Stichel, T., Stirling, C. H., Stutsman, J., Swarr, G. J., Swift, J. H., Thomas, A., Thorne, K., Till, C. P., Till, R., Townsend, A. T., Townsend, E., Tuerena, R., Twining, B. S., Vance, D., Velazquez, S., Venchiarutti, C., Villa-Alfageme, M., Vivancos, S. M., Voelker, A. H. L., Wake, B., Warner, M. J., Watson, R., van Weerlee, E., Weigand, M. A., Weinstein, Y., Weiss, D., Wisotzki, A., Woodward, E. M. S., Wu, J. F., Wu, Y. Z., Wuttig, K., Wyatt, N., Xiang, Y., Xie, R. F. C., Xue, Z. C., Yoshikawa, H., Zhang, J., Zhang, P., Zhao, Y., Zheng, L. J., Zheng, X. Y., Zieringer, M., Zimmer, L. A., Ziveri, P., Zunino, P., and Zurbick, C.: The GEOTRACES Intermediate Data Product 2017, *Chemical Geology*, 493, 210-223, 10.1016/j.chemgeo.2018.05.040, 2018.
- 600 Serreze, M. C., Barrett, A. P., Stroeve, J. C., Kindig, D. N., and Holland, M. M.: The emergence of surface-based Arctic amplification, *Cryosphere*, 3, 11-19, 10.5194/tc-3-11-2009, 2009.
- Sigman, D. M., Altabet, M. A., McCorkle, D. C., Francois, R., and Fischer, G.: The delta N-15 of nitrate in the Southern Ocean: Consumption of nitrate in surface waters, *Global Biogeochemical Cycles*, 13, 1149-1166, 10.1029/1999gb900038, 1999.
- 605 Sigman, D. M., Altabet, M. A., McCorkle, D. C., Francois, R., and Fischer, G.: The delta N-15 of nitrate in the Southern Ocean: Nitrogen cycling and circulation in the ocean interior, *Journal of Geophysical Research-Oceans*, 105, 19599-19614, 10.1029/2000jc000265, 2000.
- Sigman, D. M., Casciotti, K. L., Andreani, M., Barford, C., Galanter, M., and Bohlke, J. K.: A bacterial method for the nitrogen isotopic analysis of nitrate in seawater and freshwater, *Analytical Chemistry*, 73, 4145-4153, 10.1021/ac010088e, 2001.
- 610 Sigman, D. M., Robinson, R., Knapp, A. N., van Geen, A., McCorkle, D. C., Brandes, J. A., and Thunell, R. C.: Distinguishing between water column and sedimentary denitrification in the Santa Barbara Basin using the stable isotopes of nitrate, *Geochemistry Geophysics Geosystems*, 4, 1040, 10.1029/2002gc000384, 2003.
- 615 Sigman, D. M., DiFiore, P. J., Hain, M. P., Deutsch, C., and Karl, D. M.: Sinking organic matter spreads the nitrogen isotope signal of pelagic denitrification in the North Pacific, *Geophysical Research Letters*, 36, L08605, 10.1029/2008gl035784, 2009a.
- 620 Sigman, D. M., DiFiore, P. J., Hain, M. P., Deutsch, C., Wang, Y., Karl, D. M., Knapp, A. N., Lehmann, M. F., and Pantoja, S.: The dual isotopes of deep nitrate as a constraint on the cycle and budget of oceanic fixed nitrogen, *Deep-Sea Research Part I-Oceanographic Research Papers*, 56, 1419-1439, 10.1016/j.dsr.2009.04.007, 2009b.



- 625 Smedsrud, L. H., Esau, I., Ingvaldsen, R. B., Eldevik, T., Haugan, P. M., Li, C., Lien, V. S., Olsen, A., Omar, A. M., Ottera, O. H.,
 Risebrobakken, B., Sando, A. B., Semenov, V. A., and Sorokina, S. A.: The role of the Barents Sea in the Arctic climate system, *Reviews*
of Geophysics, 51, 415-449, 10.1002/rog.20017, 2013.
- 630 Sundfjord, A., Fer, I., Kasajima, Y., and Svendsen, H.: Observations of turbulent mixing and hydrography in the marginal ice zone of the
 Barents Sea, *Journal of Geophysical Research-Oceans*, 112, 10.1029/2006jc003524, 2007.
- Thibodeau, B., Bauch, D., and Voss, M.: Nitrogen dynamic in Eurasian coastal Arctic ecosystem: Insight from nitrogen isotope, *Global*
Biogeochemical Cycles, 31, 836-849, 10.1002/2016gb005593, 2017.
- 635 Torres-Valdes, S., Tsubouchi, T., Bacon, S., Naveira-Garabato, A. C., Sanders, R., McLaughlin, F. A., Petrie, B., Kattner, G., Azetsu-Scott,
 K., and Whitley, T. E.: Export of nutrients from the Arctic Ocean, *Journal of Geophysical Research-Oceans*, 118, 1625-1644,
 10.1002/jgrc.20063, 2013.
- 640 Tuerena, R. E., Ganeshram, R. S., Geibert, W., Fallick, A. E., Dougans, J., Tait, A., Henley, S. F., and Woodward, E. M. S.: Nutrient cycling
 in the Atlantic basin: The evolution of nitrate isotope signatures in water masses, *Global Biogeochemical Cycles*, 29, 1830-1844,
 10.1002/2015gb005164, 2015.
- Vage, S., Basedow, S. L., Tande, K. S., and Zhou, M.: Physical structure of the Barents Sea Polar Front near Storbanken in August 2007,
Journal of Marine Systems, 130, 256-262, 10.1016/j.jmarsys.2011.11.019, 2014.
- 645 Van Oostende, N., Fawcett, S. E., Marconi, D., Lueders-Dumont, J., Sabadel, A. J. M., Woodward, E. M. S., Jonsson, B. F., Sigman, D. M.,
 and Ward, B. B.: Variation of summer phytoplankton community composition and its relationship to nitrate and regenerated nitrogen
 assimilation across the North Atlantic Ocean, *Deep-Sea Research Part I-Oceanographic Research Papers*, 121, 79-94,
 10.1016/j.dsr.2016.12.012, 2017.
- 650 Weigand, M. A., Foriel, J., Barnett, B., Oleynik, S., and Sigman, D. M.: Updates to instrumentation and protocols for isotopic analysis of
 nitrate by the denitrifier method, *Rapid Communications in Mass Spectrometry*, 30, 1365-1383, 10.1002/rcm.7570, 2016.
- 655 Woodgate, R. A.: Increases in the Pacific inflow to the Arctic from 1990 to 2015, and insights into seasonal trends and driving mechanisms
 from year-round Bering Strait mooring data, *Progress in Oceanography*, 160, 124-154, 10.1016/j.pcean.2017.12.007, 2018.




Figure 1. (a) Station locations within the Barents Sea from JR16006 during July 2017 and on repeat FerryBox transects in March, June, August and November 2018. Shading is the depth (in metres). The grey contours mark the 200 m, 300 m and 500 m isobaths. Key bathymetric features are marked: Bear Island Trough (BIT), Hopen Trench (HT), Spitzbergen Bank (SB), Central Bank (CB), Great Bank (GB), Storfjordrenna (SR), Kong Karls Land (KKL) and the Barents Sea Opening (BSO). Distances along the transects presented in Figure 2 and 3 are marked. (b) Schematic of the circulation of Atlantic Water (AW), Arctic Water (ArW) and the Norwegian Coastal Current (NCC). Shading is the July 2017 sea surface temperature gradient ($^{\circ}\text{C km}^{-1}$) calculated from the OSTIA SST product (Donlon et al., 2012) and shows the sea surface temperature expression of the Polar Front along the edges of Spitzbergen Bank and Grand Bank. The solid black contour marks the July 2017 sea ice edge (data from the National Snow and Ice Data Center). Grey bathymetric contours as in (a).



Table 1. Mean and standard deviation of nitrate concentration, N^* , $\delta^{15}N\text{-NO}_3$ and $\delta^{18}O\text{-NO}_3$ for all samples from the base of the mixed layer and 500m classified as being either Atlantic Water, Barents Sea Water or Arctic Water according to the temperature and salinity characteristics used by Oziel et al., 2016. To account for nitrate utilisation, only samples where nitrate was within 1SD of mean nitrate for a given water mass were used to estimate $\delta^{15}N\text{-NO}_3$ and $\delta^{18}O\text{-NO}_3$. For reference, the salinity thresholds are contoured on Figures 2 and 3.

675

Water mass	Temp. °C	Salinity	Density kg m ⁻³	Nitrate μM	N^* μM	$\delta^{15}N\text{-NO}_3$ ‰	$\delta^{18}O\text{-NO}_3$ ‰
Atlantic Water (AW) (78-500m)	>3	>34.8		11.8 ± 1.2 n=23	-0.2 ± 0.8 n=23	5.1 ± 0.1 n=22 5.0 ± 0.1 (>200m)	2.8 ± 0.3 n=20 2.6 ± 0.3 (>200m)
Arctic Water (ArW) (46-500m)	<0	<34.7		10 ± 1.1 n=13	-1.0 ± 0.7 n=13	5.1 ± 0.1 n=12	1.6 ± 0.3 n=12
Barents Sea Water (BSW) (60-500m)	<2	>34.8	>1027.8	10.4 ± 1.2 n=23	-1.1 ± 1.1 n=21	5.1 ± 0.4 n=20	2.2 ± 0.5 n=19




Figure 2. Full 2017 transect of (a) temperature, (b) salinity, (c) nitrate, and (d) NH_4^+ . Transect is displayed in Figure 1. Solid black and white lines are the 34.8 and 34.7 isohalines. Atlantic Water (AW), Arctic Water (ArW), Barents Sea Water (BSW) and the Norwegian Coastal Current (NCC) are indicated in (a). For reference, key bathymetry features are marked in (a): Spitzbergen Bank



(SB), Storfjordrenna (SR), Hopen Trench (HT), Bear Island Trough (BIT) and the Spitzbergen-Grand Bank Sill (SB-GB). PF marks the location of the surface expression of the Polar Front.






Figure 3. Full 2017 transect of (a) $\delta^{15}\text{N-NO}_3$, (b) $\delta^{18}\text{O-NO}_3$, (c) N^* , and (d) percentage of regenerated nitrate. Transect is displayed in Figure 1. Solid black and white lines are the 34.8 and 34.7 isohalines. Atlantic Water (AW), Arctic Water (ArW), Barents Sea Water (BSW) and the Norwegian Coastal Current (NCC) are indicated in (a). The percentage of regenerated nitrate is predicted using $\delta^{18}\text{O-NO}_3$ source values from the Atlantic (2.8‰) and a nitrified value calculated using the relationship between $\delta^{18}\text{O-H}_2\text{O}$ and temperature for the Barents Sea, and a nitrification value of 1.1‰. Circles with grey outline show values not plotted to colour scale as outside of the range used in the plot.





Figure 4. Depth profiles in upper 500 m of the Barents Sea shelf. Colour denotes temperature changes. (a) Salinity, (b) Nitrate (μM), (d) NH_4^+ (μM), (d) $\delta^{15}\text{N-NO}_3$ (‰ vs. AIR), (e) $\delta^{18}\text{O-NO}_3$ (‰ vs. VSMOW) and (f) $\Delta(15-18)$ (‰). Need to add (a), (b) to subplots.



695 **Figure 5.** (a) $\delta^{15}\text{N}-\text{NO}_3$ vs $\ln\text{NO}_3$, from JR16006 showing the isotope effect (ϵ) for individual stations and an average of 4.9‰. (b) $\delta^{18}\text{O}-\text{NO}_3$ vs $\ln\text{NO}_3$, from JR16006 showing ϵ for individual stations and an average of 6.0‰. In (a) and (b) only stations with 3+measurements in the upper 120m are used and ϵ is calculated from stations with less than a 0.2 unit change in salinity. (c) $\delta^{15}\text{N}-\text{NO}_3$ vs $\ln\text{NO}_3$ for all samples from JR16006. Atlantic Water $\epsilon = 4.7\text{‰}$ and Arctic Water $\epsilon = 1.8\text{‰}$. (d) $\delta^{15}\text{N}-\text{NO}_3$ vs $\ln\text{NO}_3$ for seasonal Norbjorn samples between 72 and 76°N from the Barents Sea Opening, $\epsilon = 4.1\text{‰}$.




Figure 6. (a) $\delta^{15}\text{N-NO}_3$ (circles) and $\delta^{15}\text{N-PN}$ (triangles) plotted against nitrate (JR16006), with the colour axis denoting changes in salinity. The lines show the fractionation models for a closed system at $\epsilon=4$ and 6‰ with an initial nitrate concentration of $11.8\mu\text{M}$ and $\delta^{15}\text{N-NO}_3$ of 5.1‰ . (b) $\delta^{15}\text{N-NO}_3$ (circles) and $\delta^{15}\text{N-PN}$ (triangles) plotted against nitrate from the March, June, August and November Norbjørn transects, with the colour denoting the month of sampling. The lines show the fractionation models for a closed system at $\epsilon=4$ and 6‰ . (c) Regression between measured and predicted $\delta^{15}\text{N-PN}$ in the upper 40m using a 5‰ fractionation for all samples. (d) Regression between measured and predicted $\delta^{15}\text{N-PN}$ in the upper 40m using a 5‰ fractionation for AW and a 1.8‰ fractionation for ArW.



710




Figure 7. Seasonal variability in (a) Nitrate (μM), (b) Nitrite (μM), (c) PON ($\mu\text{g L}^{-1}$), (d) $\delta^{15}\text{N-NO}_3$ (‰), (e) $\delta^{18}\text{O-NO}_3$ (‰) and (f) $\delta^{15}\text{N-PN}$ (‰), in surface waters along the Norbjørn Ferrybox transect of the Barents Sea Opening. The transect was completed in four different months of 2018, March = orange, June = blue, August = green, and November = pink.