

**Results from a
mesocosm
perturbation study**

J. Piontek et al.

Response of bacterioplankton activity in an Arctic fjord system to elevated $p\text{CO}_2$: results from a mesocosm perturbation study

J. Piontek^{1,2}, C. Borchard^{1,2}, M. Sperling^{1,2}, K. G. Schulz¹, U. Riebesell¹, and A. Engel^{1,2}

¹Helmholtz Centre for Ocean Research Kiel (GEOMAR), Germany

²Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany

Received: 10 July 2012 – Accepted: 30 July 2012 – Published: 8 August 2012

Correspondence to: J. Piontek (jpiontek@geomar.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The effect of elevated seawater carbon dioxide (CO_2) on the activity of a natural bacterioplankton community in an Arctic fjord system was investigated by a mesocosm perturbation study in the frame of the European Project on Ocean Acidification (EPOCA). A $p\text{CO}_2$ range of 175–1085 μatm was set up in nine mesocosms deployed in the Kongsfjorden (Svalbard). The bacterioplankton communities responded to rising chlorophyll *a* concentrations after a lag phase of only a few days with increasing protein production and extracellular enzyme activity and revealed a close coupling of heterotrophic bacterial activity to phytoplankton productivity in this experiment. The natural extracellular enzyme assemblages showed increased activity in response to moderate acidification. A decrease in seawater pH of 0.5 units roughly doubled rates of β -glucosidase and leucine-aminopeptidase. Activities of extracellular enzymes in the mesocosms were directly related to both seawater pH and primary production. Also primary production and bacterial protein production in the mesocosms at different $p\text{CO}_2$ were positively correlated. Therefore, it can be suggested that the efficient heterotrophic carbon utilization in this Arctic microbial food web had the potential to counteract increased phytoplankton production that was achieved under elevated $p\text{CO}_2$ in this study. However, our results also show that the transfer of beneficial $p\text{CO}_2$ -related effects on the cellular bacterial metabolism to the scale of community activity and organic matter degradation can be mitigated by the top-down control of bacterial abundances in natural microbial communities.

1 Introduction

After early investigations that suggested heterotrophic bacterial remineralization to be low in perennially cold environments (Pomeroy and Deibel, 1986), a multitude of field observations and experimental studies over the last two decades revealed a fully active microbial loop in the polar oceans (Rivkin et al., 1996; Rich et al., 1997; Yager et al.,

BGD

9, 10467–10511, 2012

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2001). Also in the Kongsfjorden (Svalbard, 79° N, 12° E), where the present study took place, heterotrophic bacteria contribute substantially to the biomass of the microbial community and play an important role in the structure and function of the microbial food web in all seasons (Iversen and Seuthe, 2011; Seuthe et al., 2011). During the vernal bloom in April bacteria are subject to intense grazing by heterotrophic dinoflagellates and ciliates, thereby acting as an important trophic link for organic carbon in microbial food webs. In the following nutrient-limited post-bloom period the microbial food web enters a regenerative state that is characterized by an efficient recycling of produced carbon by the microbial loop (Iversen and Seuthe, 2011; Seuthe et al., 2011). Hence, metabolic activity of heterotrophic bacterioplankton can be expected to be a major constraint on fluxes of labile organic carbon and nutrients in this high-latitude ecosystem.

The bottom-up regulation of bacterial activity in Arctic microbial food webs is mainly achieved by interactions with temperature and dissolved organic matter (DOM) (Pomeroy and Wiebe, 2001; Kirchman et al., 2005, 2009a). Psychrophilic and psychrotolerant heterotrophic bacteria in polar marine environments are physiologically adapted to low temperatures by biochemical properties of their cell constituents like cold-active enzymes of reduced activation energy and membrane lipids that sustain fluidity at subzero temperatures (Deming, 2002). Nevertheless, bacteria in polar oceans often act far below their temperature optimum, so that cold ambient temperature has a high potential to reduce their metabolic activity and growth that, in turn, determine the rate of bacterial DOM degradation (e.g. Simon et al., 1999; Huston et al., 2004). More recently, the availability of labile and semilabile DOM was proposed as a major constraint on bacterial activity in the Arctic Ocean (Kirchman et al., 2009a). Concentrations of dissolved organic carbon (DOC) determined in Arctic seawater are in the range of temperate oceans but comprise a large terrestrial fraction that is discharged by rivers and meltwater runoff (Anderson, 2002; Dittmar and Kattner, 2003). These terrestrial compounds do not provide labile carbon for bacterial utilization and consequently do not support much bacterial growth (Meon and Amon, 2004; Hansell et al., 2004). Thus, Arctic marine systems are not only characterized by low temperature but

Results from a mesocosm perturbation study

J. Piontek et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

also by a deficiency of carbon and energy resources for heterotrophic microbial growth (Kirchman et al., 2009a).

There is growing evidence that impacts of climate change are amplified by the Arctic environment, making the Arctic Ocean in particular sensitive to warming and acidification. A temperature increase of 0.4 °C in the Arctic over the past 150 yr corresponds to a rate two to three times higher than the global average, amplified by a decreasing surface albedo at enhanced ice and snow melting (Overpeck et al., 1997). The additional freshwater input from ice melting, in turn, reduces the alkalinity of seawater and hence the buffering capacity of the Arctic Ocean to acidification (Steinacher et al., 2009). Investigating the relevance of seawater pH for the structure and functioning of marine microbial communities gained little attention in past marine research, although substantial natural variability of pH occurs on short time scales and in marine microhabitats like particle aggregates and biofilms (Ploug et al., 1997; Dexter and Chandrasekaran, 2000). Biochemical studies demonstrated that fundamental processes of the marine microbial metabolism like numerous enzymatic reactions and the respiratory electron transfer are strongly dependent on pH conditions (e.g. Groudieva et al., 2004; Kolber, 2007). Ongoing systematic changes in seawater pH, however, strongly require a better mechanistic understanding of pH and $p\text{CO}_2$ effects on the regulation of both autotrophic and heterotrophic microbial activity in natural communities to assess consequences of ocean acidification for food web processes and organic matter turnover.

Our study was part of the joint EPOCA mesocosm experiment conducted in Kongsfjorden (Svalbard), 2010. The Kongsfjorden system in West Spitsbergen is mainly affected by the northbound transport of Atlantic water masses by the West Spitsbergen Current. It is an open fjord system without sill and, therefore, largely affected by mixing processes on the adjacent shelf (Hop et al., 2006). Nine mesocosms were deployed in the fjord during June and July, 2010, to investigate the CO_2 sensitivity of Arctic pelagic organisms and resulting consequences for the production and turnover of organic matter. More specifically, we tested in the framework of this experiment (i) whether bacterial

BGD

9, 10467–10511, 2012

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



protein production and rates of extracellular enzymes in the natural Kongsfjorden plankton community change under elevated $p\text{CO}_2$ and (ii) how changes in $p\text{CO}_2$ can induce systematic effects on heterotrophic bacterial activity in this perennially cold marine environment.

2 Material and methods

2.1 Mesocosm set up and sampling

A mesocosm study was conducted on Spitsbergen (Svalbard, 79° N, 12° E) between 30 May and 7 July 2010, in the framework of the European Project on Acidification (EPOCA). Kiel Off-Shore Mesocosms for Future Ocean Simulations (KOSMOS) were deployed in the Kongsfjorden, (78°56,2' N, 11°53,6' E), an Arctic glacial fjord system located on the west coast of Spitsbergen that is influenced by both Atlantic and Arctic water masses (Hop et al., 2002). A detailed description of the mesocosm set up is given by Riebesell et al. (2012) and Schulz et al. (2012). Briefly, nine mesocosms, each consisting of a 15 m long polyurethane bag, were deployed close to the shoreline. Each mesocosm enclosed about 45 m³ of seawater. A $p\text{CO}_2$ gradient that ranged initially from 250–1085 μatm was adjusted by the stepwise addition of CO_2 -enriched seawater during days –1–4. Two mesocosms at in situ $p\text{CO}_2$ of 175–180 μatm served as controls. The low in situ $p\text{CO}_2$ reflected the post-bloom stage of a *Phaeocystis*-dominated bloom event (F. Buchholz, personal communication, 2010). In the first phase of the experiment (days 4–13), the development of the mesocosm plankton communities was observed without any further manipulation under nutrient-deplete conditions. Nine days after the adjustment of target $p\text{CO}_2$ -levels (on day 13), 5 μmol nitrate l^{-1} , 0.3 μmol phosphate l^{-1} , and 2.5 μmol silicate l^{-1} were added to enhance phytoplankton production.

Depth-integrated samples of the water column enclosed by the mesocosm bags were taken by the use of a motor-operated water sampler (Hydrobios, Kiel, Germany) that

BGD

9, 10467–10511, 2012

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



permanently and evenly collected water while being lowered from the surface to 12 m depth (Riebesell et al., 2012). Depth-integrated samples of all mesocosms and the fjord were analysed for extracellular enzyme activity and bacterial protein production daily or every second day.

2.2 Rates of extracellular enzymes

Activity of extracellular enzymes was determined by the use of fluorogenic substrate analogues (Hoppe, 1983). Accordingly, rates of β -glucosidase, leucine (leu)-aminopeptidase, and alkaline phosphatase were assessed from the hydrolysis of 4-methylumbelliferyl- β -glucopyranoside, L-leucyl-4-methylcoumarinylamid-hydrochlorid, and 4-methylumbelliferyl-phosphate, respectively. The substrate analogues were added to whole seawater samples at final concentrations of 1, 5, 10, 20, 50, 80, 100, and 200 $\mu\text{mol l}^{-1}$ to achieve concentration kinetics. The fluorescence emitted by 4-methylumbelliferone (MUF) and 7-amino-4-methyl-coumarine (AMC) after enzymatic cleavage of the substrate analogues was detected at 355 nm excitation and 460 nm emission wavelength. Relative fluorescence units were converted into concentrations of MUF and AMC, respectively, after calibration with standard solutions of 2–100 nmol l^{-1} . Calibration factors were determined at five different pH values ranging from 7.6 to 8.4 to consider the pH dependence of MUF and AMC fluorescence.

Enzymatic rates were calculated from the increase in MUF and AMC concentration, respectively, over time. An initial fluorescence measurement was conducted immediately after the addition of the substrate analogue followed by 3–4 measurements within 24 h of incubation in the dark at 2 °C. On days 12, 14, 20, and 24 of the mesocosm study additional incubations at 5.5–6.5 °C were conducted. Instead of using single end point measurements, the slope of the linear regression between MUF and AMC concentrations, respectively, and time was applied for rate calculations. Two analytical replicates per sample revealed on average a standard deviation of 9%. Experimental data were fitted to a kinetic model using the Michaelis-Menten equation to determine the maximum velocity (V_{max}) and the half-saturation constant K_m of the enzymatic reactions.

BGD

9, 10467–10511, 2012

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3 Bacterial protein production

Bacterial protein production (BPP) was estimated from the uptake of ^{14}C -leucine that was added to depth-integrated samples of mesocosms and fjord at 40 nmol l^{-1} final concentration. Incubations were terminated by the addition of trichloroacetic acid (TCA) at a final concentration of 5% and processed by the microrcentrifuge method (Smith and Azam, 1992). Briefly, samples were centrifuged at $14\,000 \times g$ to gain a cell pellet that was washed twice with 5% TCA. Incorporation into the TCA-insoluble fraction was measured by liquid scintillation counting after resuspension of the cell pellet in scintillation cocktail. Tests revealed a linear increase of leucine uptake over at least 50 h, so that extended incubation times of up to 24 h could be applied at low bacterial activity. Two series of incubations were conducted in a temperature-controlled walk-in room at 2°C ($\text{BPP}_{2^\circ\text{C}}$) and in situ in 1 m water depth, respectively. Duplicate incubations revealed an analytical error $\leq 10\%$ for both incubation modes. Leucine incorporation was converted into BPP applying a conversion factor of 1.5 kg C mol^{-1} leucine, assuming no intracellular isotope dilution (Simon and Azam, 1989).

2.4 Cell-specific activity

Cell-specific rates of extracellular enzymes and BPP were determined by dividing the rate scaled to volume by total bacterial abundance. Specific growth rates (μ) were calculated according to

$$\mu = \text{BPP}/B \quad (1)$$

where BPP is bacterial protein production ($\text{ng C l}^{-1} \text{d}^{-1}$) and B bacterial biomass (ng C l^{-1}) (Kirchman, 2001).

In order to calculate cell-specific rates and specific growth rates as precisely as possible bacterial cell numbers were determined in the mesocosm subsamples that were used for the analysis of extracellular enzyme activity and BPP. Bacterial abundances were determined by flow cytometry (FACSCalibur, Becton Dickinson) after nucleic acid

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



staining with SybrGreen I (Invitrogen). Bacterial cell numbers were estimated after visual inspection and manual gating of the bacterial subpopulation in the cytogram of side scatter vs. green fluorescence. Yellow-green fluorescent latex beads (Polyscience) and TruCount beads (Becton Dickinson) were used to normalize the counted events to volume (Gasol and del Giorgio, 2000). Bacterial cell numbers were converted into bacterial biomass assuming a carbon content of $10 \text{ fg C cell}^{-1}$ (Caron et al., 1995).

2.5 ^{14}C primary production

Primary production was determined by the use of the radiotracer $\text{NaH}^{14}\text{CO}_3$ (Steeermann Nielsen, 1952; Gargas, 1975). Briefly, samples spiked with $8 \mu\text{mol l}^{-1}$ $\text{NaH}^{14}\text{CO}_3$ were incubated in situ at 1 m depth for 24 h. After incubation samples were filtered onto $0.4 \mu\text{m}$ polycarbonate filters. Particulate primary production was determined from the particulate matter collected on the filters. Subsamples of the filtrate were used to determine exudation. Both filters and filtrate were acidified to remove inorganic carbon prior to analysis by liquid scintillation counting. A detailed method description is given by Engel et al. (2012).

2.6 Supplemental experiments

2.6.1 Enrichment assay

An enrichment assay was conducted to investigate effects of labile carbon and inorganic nitrogen input on bacterial activity and growth. For this purpose, a sample integrated over 12 m depth was collected close to the mesocosm site in Kongsfjorden during the days of mesocosm deployment on 2 June 2010. Whole seawater samples of 20 ml each were supplemented in triplicate with glucose and ammonium separately and in combination at final concentrations of $20 \mu\text{mol l}^{-1}$ and $10 \mu\text{mol l}^{-1}$, respectively. Samples without any addition served as control. Rates of extracellular enzyme activity, BPP, and bacterial cell numbers were determined according to the methods described

BGD

9, 10467–10511, 2012

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



above in the initial fjord sample and in all incubations after 4 days in the dark at in situ temperature of 2 °C.

2.6.2 Acidification assay

The biochemical sensitivity of the natural extracellular enzyme assemblages to decreasing seawater pH was investigated by an acidification assay. Depth-integrated samples collected in the fjord and one non-acidified control mesocosm on day 25 of the study (2 July 2010) were acidified by the addition of 225–750 µl 0.1 N hydrochloric acid to 200 ml of sample. Samples were allowed to equilibrate at 2 °C in the dark for 3 h before pH was measured with a proton-sensitive combined pH-temperature electrode (WTW, Sentix 41; standard DIN/NBS buffers) and the initial sampling was conducted. Non-manipulated samples of the fjord and the mesocosm served as control incubations at in situ pH. Measurements for β -glucosidase and leu-aminopeptidase activity were accomplished at 4 time points within 24 h after the initial time point.

2.7 Data analysis

To test the effect of temperature on BPP and extracellular enzyme activity, Q_{10} factors were calculated as

$$Q_{10} = (r_2/r_1)^{10/(T_2-T_1)} \quad (2)$$

where r_2 and r_1 are the rates at high (T_2) and low (T_1) temperature, respectively.

The hydrolysis potential for polysaccharides and proteins ($\mu\text{mol l}^{-1}$) was calculated by integrating V_{max} of β -glucosidase and leu-aminopeptidase ($\mu\text{mol l}^{-1} \text{d}^{-1}$), respectively, over time according to

$$\sum_{t=m}^n r_t \quad (3)$$

where m and n are the initial and final time point, respectively, and r_t the rate of an individual day. Since rate measurements for extracellular enzymes were conducted

BGD

9, 10467–10511, 2012

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



every second day of the experiment rates for the missing days were interpolated by

$$r_t = \frac{(r_{t-1} + r_{t+1})}{2} \quad (4)$$

where r_{t-1} is the measured rate of the previous day and r_{t+1} the measured rate of the day after. To examine differences in bulk polysaccharide and protein hydrolysis the difference between time-integrated enzymatic rates of the 7 acidified mesocosms and the mean of the two control mesocosms could be calculated and is referred to as Δ hydrolysis potential.

Differences were tested for significance by means of t -test, paired t -test and, if tests for normality or equal variance failed, Wilcoxon signed rank test. Linear and non-linear regression analyses were conducted using the software Sigma Plot 12.1 (Systat). Significance was accepted for $p \leq 0.05$.

3 Results

3.1 Enrichment assays

The substrate supply to the bacterioplankton community in the Kongsfjorden was tested for limiting resources by an enrichment assay at the beginning of the mesocosm study. The amendment of glucose led to a stimulation of bacterial growth within 4 days of incubation. Adding glucose alone or in combination with ammonium increased cell numbers and BPP by 1.4 and 1.8 times the control incubation, respectively. Sole ammonium enrichment did not result in significant changes (Fig. 1). These results show that growth of the bacterioplankton community in the fjord was limited by the deficiency of labile carbon or energy when the mesocosm study started. Cell-specific V_{\max} of leu-aminopeptidase remained constant over incubation time and showed similar values of $32.7\text{--}36.9 \text{ amol cell}^{-1} \text{ h}^{-1}$ in non-amended controls and in amended incubations after four days (Fig. 1). This suggests that independent of carbon or energy limitation also

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the demand for organic nitrogen was high and could not be compensated by the supply of an inorganic nitrogen source. The initial β -glucosidase V_{\max} of $3.3 \text{ amol cell}^{-1} \text{ h}^{-1}$ decreased during incubation in all treatments. Nevertheless, rates determined after 4 days revealed significant differences between incubations with and without glucose (t -test, $p < 0.001$). Cell-specific V_{\max} of β -glucosidase in glucose-amended incubations were reduced by approximately 37 % to $2.0\text{--}2.2 \text{ amol cell}^{-1} \text{ h}^{-1}$, while rates in controls and ammonium-amended samples decreased strongly by approximately 83 % to $0.5\text{--}0.7 \text{ amol cell}^{-1} \text{ h}^{-1}$ (Fig. 1).

3.2 Bacterial protein production and specific growth rates

The synthesis of cellular proteins by heterotrophic bacterioplankton showed a similar temporal development in all mesocosms. Lowest $\text{BPP}_{20^\circ\text{C}}$ of $19\text{--}56 \text{ ng Cl}^{-1} \text{ h}^{-1}$ was determined shortly after CO_2 manipulation (days 2–7). Thereafter, $\text{BPP}_{20^\circ\text{C}}$ did not show a continuous increase but elevated values during days 12–16 and 22–29 after the first and the second maximum of chlorophyll *a* concentrations, respectively (Fig. 2). Maximum $\text{BPP}_{20^\circ\text{C}}$ of $168\text{--}268 \text{ ng Cl}^{-1} \text{ h}^{-1}$ was attained either on day 22 or day 28, representing on average an increase by a factor of 6.4 over 18–22 days of the experiment. Specific growth rates of bacteria showed an opposite temporal development. Maximum specific growth rates of $0.24\text{--}0.37 \text{ d}^{-1}$ were determined during the initial days of the experiment. Specific growth rates declined strongly after day 8 so that highest $\text{BPP}_{20^\circ\text{C}}$ at the end of the experiment coincided with lowest specific growth rates of $0.02\text{--}0.09 \text{ d}^{-1}$ (Fig. 2).

$\text{BPP}_{20^\circ\text{C}}$ in the fjord samples was in the range of the mesocosm samples on 11 of 20 sampling days. On 8 sampling days $\text{BPP}_{20^\circ\text{C}}$ in the fjord exceeded the maximum mesocosm value at the same day. Highest fjord $\text{BPP}_{20^\circ\text{C}}$ of $421 \text{ ng Cl}^{-1} \text{ h}^{-1}$ was determined on day 10, when the rate was almost twice the maximum mesocosm value (Figs. 2–3).

BGD

9, 10467–10511, 2012

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.3 Activity of hydrolytic extracellular enzymes

Rates of extracellular β -glucosidase, leu-aminopeptidase, and phosphatase were measured in mesocosm and fjord samples to assess the hydrolytic activity of the three major types of extracellular enzymes that drive the turnover of organic polymers in seawater. V_{\max} of β -glucosidase and leu-aminopeptidase started to increase in all mesocosms during days 6–8, shortly after CO_2 manipulation and in coincidence with the development of a first chlorophyll *a* maximum. Activities remained rather constant between days 12 and 20 followed by further increasing rates of both enzymes (Figs. 4–5). Maximum V_{\max} in the mesocosms ranged from 3.1–4.8 $\text{nmol l}^{-1} \text{h}^{-1}$ and 62.1–86.7 $\text{nmol l}^{-1} \text{h}^{-1}$ for β -glucosidase and leu-aminopeptidase, respectively, and were determined during the last days of the experiment.

The temporal development of cell-specific rates was different from that of bulk rates. Cell-specific V_{\max} in the mesocosms increased from days 4 to 12 up to 0.9–1.4 $\text{amol cell}^{-1} \text{h}^{-1}$ and 12.2–20.0 $\text{amol cell}^{-1} \text{h}^{-1}$ for β -glucosidase and leu-aminopeptidase, respectively, but showed only minor variation afterwards (Figs. 4–5). Hence, an increasing amount of bacterial enzymes released into seawater by a growing bacterioplankton community and not enhanced cellular metabolic activity increased the hydrolytic potential of the bacterioplankton community after day 12.

The temporal development of Δ *hydrolysis potential* reveals an elevated enzymatic potential for polysaccharide and protein degradation in the three mesocosms of highest $p\text{CO}_2$ during the first 20 days of the study (Fig. 6). After the second chlorophyll *a* maximum on day 21 bacterial abundances in the mesocosms diverged and the differences between the three high- CO_2 mesocosms and the controls were either reduced or compensated (Fig. 6).

Maximum rates of β -glucosidase and leu-aminopeptidase in the fjord were determined on day 8 and –3, respectively. For both enzymes elevated rates were measured during and shortly after the chlorophyll *a* maximum on days 10–14. After day 16 rates

BGD

9, 10467–10511, 2012

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of both enzymes did not show substantial variation in the fjord despite continuously increasing bacterial cell numbers (Fig. 3; Brussaard et al., 2012).

Mesocosms and fjord showed similar K_m values for leu-aminopeptidase activity with mean values of 54 and 50 $\mu\text{mol l}^{-1}$, respectively. In contrast, the K_m values of β -glucosidase were significantly higher in the mesocosms (average 94 $\mu\text{mol l}^{-1}$) (Mann-Whitney rank sum test, $p < 0.05$), revealing a lower affinity to substrates than in the fjord (average 52 $\mu\text{mol l}^{-1}$) (Table 1).

The ratio of leu-aminopeptidase: β -glucosidase activity is a measure for the protein relative to polysaccharide hydrolysis. This activity ratio changed in mesocosms and fjord over time. The temporal development in mesocosms and Kongsfjorden diverged after day 12 when the ratio increased in the mesocosms but remained lower in the fjord samples. Hence, the comparison of mesocosms and fjord reveals a higher potential for protein turnover at relatively lower polysaccharide degradation in the mesocosms after day 12. The divergence in the activity ratio of the two enzymes coincided with the divergence of the molar ratios of DON : DOC in mesocosms and fjord. The share of DON in the mesocosms was substantially lower than in the Kongsfjorden after day 12 of the experiment (Fig. 7).

Concentration kinetics of extracellular phosphatase showed that substrate concentrations of 200 $\mu\text{mol l}^{-1}$, the maximum applied in our incubations, were not saturating at several time points, so that V_{max} was not reached. Therefore, rates derived from a non-saturating substrate concentration of 10 $\mu\text{mol l}^{-1}$ are presented (Fig. 8). Hence, phosphatase rates are not directly comparable with rates of β -glucosidase and leu-aminopeptidase in this study, since rates shown for these two enzymes represent V_{max} . The temporal development of phosphatase activity shows a clearly different pattern than do β -glucosidase and leu-aminopeptidase. Rates show three activity maxima in all mesocosms during the experiment (Fig. 8). The first maximum in phosphatase activity during days 4–8 coincided with the first maximum in chlorophyll *a* concentrations. The second phosphatase maximum occurred at chlorophyll *a* minimum concentrations between the first and the second chlorophyll *a* peak during days 12–18. A third maximum

BGD

9, 10467–10511, 2012

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in phosphatase activity developed simultaneously with the third chlorophyll *a* maximum between days 22 and 28. However, this third phosphatase maximum had lower rates than the two previous maxima in most mesocosms (Fig. 8).

Phosphatase activity in the fjord increased from the beginning of the experiment until day 10 (Fig. 3). After a strong decrease in rates during days 12–14 phosphatase activity was below the detection limit in samples of the Kongsfjorden from day 16 until the end of the experiment. Extracellular phosphatase activity in the fjord was significantly related to chlorophyll *a* concentrations over the whole duration of the experiment (linear regression, $r^2 = 0.55$, $p < 0.001$, $n = 17$; not shown).

3.4 Temperature dependence of bacterial protein production and extracellular enzymes

The seawater temperature increased during the mesocosm study from 2.0 °C at the beginning of June to 5.2 °C one month later. There were no temperature differences between the nine mesocosms and also Q_{10} values determined for extracellular enzyme activity and BPP did not show significant variation between the mesocosms on the individual sampling days. However, Q_{10} values varied significantly over the duration of the study, revealing that the temperature sensitivity of the enzyme assemblages changed within two weeks. Q_{10} values for β -glucosidase and leu-aminopeptidase determined at four time points between day 12 and 24 ranged from 2.0–11.4 and 1.1–4.4 (mean values of all mesocosms), respectively. Activity of β -glucosidase was consistently more temperature sensitive than leu-aminopeptidase activity. Also Q_{10} values determined for BPP at nine sampling days showed high temporal variability and ranged from 0.1–11.7 (Fig. 9). The Q_{10} values for both enzymes and for BPP were related to phytoplankton exudation that can be considered as a measure for the input of labile organic matter. The Q_{10} values determined for the two extracellular enzymes at lower exudation levels of 0.9 and 0.7 $\mu\text{molCl}^{-1} \text{d}^{-1}$, respectively, were significantly higher than those at increased exudation of 1.5 $\mu\text{molCl}^{-1} \text{d}^{-1}$ (Mann-Whitney rank sum test, $p < 0.001$). Also

BGD

9, 10467–10511, 2012

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Q_{10} values for BPP decreased significantly with increasing exudation (linear regression, $r^2 = 0.51$, $p = 0.03$) (Fig. 9).

Q_{10} values for BPP are derived from a comparison of rates determined at 2 °C and at in situ temperature. Since 2 °C represents the initial fjord temperature during mesocosm deployment and CO₂-manipulation, the difference between bPP at the two temperatures represent the surplus of bacterial biomass production that can be attributed to seawater warming during the experiment. Summing up this surplus over the experiment, it can be estimated that rising in situ temperature alone increased bPP in the mesocosms by a factor of 1.3–1.7, corresponding to 27–61 $\mu\text{gC l}^{-1}$.

3.5 Testing for effects of different mesocosm $p\text{CO}_2$ levels on bacterial activity

The design of the mesocosm experiment with 8 different $p\text{CO}_2$ levels suggests the use of a regression model to test data for significant relationships between seawater $p\text{CO}_2$ and bacterial activity. Physiological performance and metabolic activity of heterotrophic microbial organisms like bacteria may not be directly affected by changes in seawater $p\text{CO}_2$ but by co-occurring changes in pH. The range of mesocosm $p\text{CO}_2$ from 175 to 1085 μatm corresponded to pH values of 7.63–8.34. Regression analysis was carried out with differences in proton concentration that could be calculated between each acidified mesocosm and the non-acidified control mesocosms ($\Delta [\text{H}]^+$, nmol l^{-1}). The resulting mean for $\Delta [\text{H}]^+$ over time were correlated with time-integrated rates of BPP_{2°C}, β -glucosidase and leu-aminopeptidase (Eq. 3) (Figs. 10–11). The temporal development of Δ hydrolysis potential reveals an elevated enzymatic potential for polysaccharide and protein degradation in the three mesocosms of highest $p\text{CO}_2$ during the first 20 days of the study when similar bacterial abundances were determined in all mesocosms (Fig. 6). Therefore, regression analyses of time-integrated β -glucosidase and leu-aminopeptidase rates were accomplished for the whole duration of the experiment (days 4–29) and also separately for the period of similar bacterial cell numbers (days 4–20). Cell-specific leu-aminopeptidase activity was directly related to differences in proton concentration over the whole experiment ($r^2 = 0.57$, $p < 0.05$) (Fig. 10). Hence,

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the cellular potential for enzymatic protein hydrolysis increased with gradual acidification over the applied $p\text{CO}_2$ range. Correlations of $\Delta [\text{H}]^+$ with cell-specific enzymatic rates integrated over days 4–20 revealed significant linear relationships for both β -glucosidase ($r^2 = 0.53$, $p < 0.05$) and leu-aminopeptidase activity ($r^2 = 0.79$, $p < 0.05$) (Fig. 10), and confirmed the trend of rising hydrolytic activity mediated by bacterial extracellular enzymes at lowered seawater pH. A significant relationship between $\Delta [\text{H}]^+$ and $\text{BPP}_{2\cdot\text{C}}$ could not be determined.

The pH manipulation was the primary source of variation in this experiment but a comparison of bacterial activity in the mesocosms at different $p\text{CO}_2$ should also consider potential impacts of changing phytoplankton productivity under different $p\text{CO}_2$. Although not linearly related to $p\text{CO}_2$, the dependence of primary production on the mesocosm $p\text{CO}_2$ level was evident. The Arctic phytoplankton communities responded with higher primary production to elevated CO_2 in this mesocosms experiment (Engel et al., 2012). Regression analysis revealed direct linear correlations of time-integrated primary production with BPP ($r^2 = 0.77$, $p < 0.05$) and cell-specific leu-aminopeptidase activity ($r^2 = 0.57$, $p < 0.05$) (Fig. 11). Hence, bacterioplankton responded with rising biomass production and protein degradation to elevated photosynthetic production of organic substrates. Leu-aminopeptidase activity was significantly related to both seawater pH and primary production.

3.6 Acidification assay

To exclusively test the direct influence of seawater pH on β -glucosidase and leu-aminopeptidase activity samples of the Kongsfjorden and one control mesocosm (M3) were acidified with dilute hydrochloric acid on day 25 of the experiment. The pH manipulation was the only source of variation in this experiment, while a comparison of the mesocosms at different $p\text{CO}_2$ includes the divergence of other environmental and biological parameters like phytoplankton community composition (Brussaard et al., 2012), primary production (Engel et al., 2012), net community production (Silyakowa et al., 2012) and nutrient utilization (Schulz et al., 2012). Rates of extracellular β -glucosidase

BGD

9, 10467–10511, 2012

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and leu-aminopeptidase in non-manipulated subsamples with in situ pH were compared with subsamples adjusted to three lower pH-levels ranging from 7.79–8.17. Both enzymes showed highest rates at lowest pH, revealing that the in situ pH of the Kongsfjorden and the control mesocosm (M3) did not provide optimum conditions for these enzymatic reactions. Cell-specific V_{\max} of β -glucosidase and leu-aminopeptidase in acidified Kongsfjorden samples increased by a factor of 2.1 and 1.8, respectively, when the in situ pH of 8.29 was reduced to 7.79. The effect of acidification was less pronounced in mesocosm samples. Here, cell-specific V_{\max} of both enzymes increased by a factor of 1.2 at pH 7.79.

4 Discussion

In the present mesocosm study, the microbial community of an open Arctic fjord system was exposed to different $p\text{CO}_2$ ranging from in situ values of $175 \mu\text{atm}$ to $1085 \mu\text{atm}$ according to projections for the future ocean. Hence, changing $p\text{CO}_2$ was the primary manipulation in this experiment. The parameters of the seawater carbonate system must be considered as environmental factors that contribute among others to the bottom-up control of the microbial community. In the present study, also temperature that increased substantially during the experiment and phytoplankton production that changed over time and in response to different $p\text{CO}_2$ had a high potential to affect growth and metabolic activity of bacterioplankton. In the following the influence of seawater pH, temperature and phytoplankton production on heterotrophic bacterial activity and their role in the bottom-up regulation of bacterioplankton activity will be discussed.

4.1 Phytoplankton-bacterioplankton coupling

The coupling of phytoplankton and bacterioplankton becomes evident from a phasing of biomass and production of the two groups (Ducklow and Yager, 2006; Kirchman et al., 2009b). This co-variation of phytoplankton and bacterioplankton biomass and

BGD

9, 10467–10511, 2012

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



production over time and space is characterized by a succession of phytoplankton followed by bacterioplankton after a temporal lag of days to weeks. The duration of this lag period in response to increasing phytoplankton biomass and production is decisive for the degradation efficiency and the export of organic matter produced in bloom situations (Kirchman, 2009b; Bird and Karl, 1999). The bacterioplankton community in the Kongsfjorden and in the mesocosms responded after a lag of only some days with increasing activities to rising chlorophyll *a* concentrations, revealing that there was neither a substrate threshold that needed to be exceeded nor any community intrinsic factor impeding a fast bacterial response (Figs. 2–5). Previous studies have shown that bacterial communities in polar marine systems can lag behind the development of vernal phytoplankton blooms by weeks (Billen and Becquevort, 1991; Bird and Karl, 1999). Our study, however, was conducted in early summer after the main *Phaeocystis* spring bloom in the Kongsfjorden. Therefore, it seems likely that bacterial cells did not return from a starvation induced dormancy that allows maximum conservation of carbon and energy during winter. A study accomplished in the Chuckchi Sea during summer has shown that the fraction of actively respiring bacterial cells and the cells' substrate affinity increased at elevated chlorophyll levels within some days, suggesting a substantial physiological and metabolic flexibility of bacterioplankton communities at this time of the year (Yager et al., 2001). Hence, the physiological state of the bacterial cells and the community composition after the vernal bloom might facilitate a faster response to subsequent bloom events during summer and thus promote a faster onset of organic matter recycling in cold marine systems.

In addition to the lag phase, also the dimension of bacterial activity achieved in response to elevated phytoplankton production determines the rate at which freshly produced organic matter is recycled. Absolute values for growth rates and extracellular enzyme activities are a measure to assess the overall magnitude of bacterial activity, although information is limited due to broad ranges observed in marine environments and overlapping limits for cold and temperate marine ecosystems. The specific growth rates of bacteria in polar marine systems are not necessarily lower than in oligotrophic

BGD

9, 10467–10511, 2012

**Results from a
mesocosm
perturbation study**

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



temperate regions (Rivkin et al., 1996; Rich et al., 1997; Ducklow et al., 1999). Specific growth rates determined in Kongsfjorden and the mesocosms are in the range of values published for both the Arctic Ocean (Kirchman et al., 2009b) and temperate marine ecosystems (Ducklow et al., 1993; Ducklow 1999) (Fig. 2). Similarly, high variability of cellular β -glucosidase and leu-aminopeptidase rates was found. Maximum and mean cell-specific rates of β -glucosidase and leu-aminopeptidase in the mesocosms and in Kongsfjorden are in good accordance with values determined off the coast of Spitsbergen during summer (Sala et al., 2010) (Figs. 3–5). The activities are at the lower limit of ranges published for temperate marine systems like the Central Atlantic (Baltar et al., 2009) and the Baltic Sea (Nausch et al., 1998). Overall, the magnitude of bacterial activity in mesocosms and Kongsfjorden represents the overlap of higher levels in cold marine environments and lower levels in the temperate ocean.

4.2 Microbial acquisition of organic matter by the use of extracellular enzymes

Organic substrates for bacterial utilization are mainly derived from complex high-molecular-weight compounds that are enzymatically hydrolysed outside the microbial cells into molecules sufficiently small for uptake. In the marine environment, heterotrophic bacteria must be considered as the main producers of β -glucosidase and leu-aminopeptidase, the most important polysaccharide- and protein-degrading extracellular enzymes, respectively. Culture studies have shown that some species of marine phagotrophic protozoa, mainly flagellates, produce and release glucosidases and proteases but to the best of our knowledge there is no evidence to date that non-bacterial producers are able to significantly influence the bulk rate of these two enzymes in natural marine communities (Karner et al., 1994). The temporal development of cell-specific β -glucosidase and leu-aminopeptidase activity reveals information on the modulation of enzyme activity in the mesocosms. Cell-specific enzymatic rates increased during the first chlorophyll peak, demonstrating a response of bacterial activity to changing environmental conditions on the cellular level in this first phase of the experiment. Afterwards increasing bulk activity was driven by the growing community and

BGD

9, 10467–10511, 2012

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



thereby increasing numbers of bacterial enzyme producers but not by the adjustments of cellular rates. Accordingly, the bulk activity of extracellular enzymes was highest towards the end of the mesocosm experiment when highest bacterial cell numbers were determined, revealing that the enzymatic turnover of organic matter was strongly affected by growth characteristics of the bacterioplankton community in this study.

A variety of heterotrophic and autotrophic microbial organisms contributes substantially to the enzymatic hydrolysis of organic phosphorus compounds. In addition to bacteria, also phytoplankton species and protozoa can be strong producers of extracellular phosphatase in natural marine communities (Hoppe, 2003). In the present study, two maxima of phosphatase activity coincided with chlorophyll *a* maxima, while only the phosphatase peak during days 12–18 developed at low chlorophyll *a* concentrations (Fig. 8). Therefore, it seems likely that in particular before nutrient addition on day 12 autotrophic organisms recycled a significant share of organic phosphorus. As a consequence, high chlorophyll *a* concentrations could be achieved without the availability of inorganic phosphorus (Schulz et al., 2012).

4.3 Growth limitation of bacterioplankton by organic matter resources

The availability of labile and semilabile carbon is considered to be a major factor regulating bacterial activity in the marine pelagic environment, including subpolar and polar oceans (Kirchman et al., 2005, 2009a; Granéli et al., 2004; Cuevas et al., 2011). Results of the enrichment assay conducted with Kongsfjorden samples at the beginning of the mesocosm study are in accordance with this conclusion, since the amendment of labile carbon stimulated bacterial biomass production and growth (Fig. 1). A recent study revealed high spatial variability of the limiting resource for bacterial growth at coastal and fjord stations located in the west of Spitsbergen. Enhanced carbon consumption was evident in incubations of coastal samples, while mineral nutrient deficiency limited bacterioplankton production in fjord samples of high phytoplankton biomass (Vadstein, 2011). Our incubations were accomplished during the initial days of the mesocosm experiment when low levels of phytoplankton biomass and primary

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



production prevailed (Engel et al., 2012; Schulz et al., 2012). Therefore, it can be assumed that regeneration of nutrients was high and prevented mineral nutrient limitation, while the input of fresh and labile organic compounds was low and thus limiting at that time. High levels of leu-aminopeptidase activity in glucose-amended Kongsfjorden samples show that also the demand for amino acids released from protein hydrolysis was high. Hence, the limitation of bacterial production and growth by the availability of labile carbon was evident but also a deficiency of labile organic nitrogen compounds must be assumed. The enclosure of the bacterioplankton community into the mesocosms likely intensified the shortage in organic nitrogen for bacterial consumption as suggested by higher activity ratios of leu-aminopeptidase: β -glucosidase in the mesocosms than in the fjord (Fig. 7). Higher activity ratios in the mesocosms coincided with a lower share of DON and imply that the high organic nitrogen demand of the strongly growing bacterioplankton populations in the mesocosms enhanced the need for enzymatic protein hydrolysis.

Overall, the enrichment assay and the activity ratios of leu-aminopeptidase: β -glucosidase in mesocosms and fjord demonstrate that organic matter resources had a high potential to affect bacterial growth characteristics in this study. Furthermore, results reveal an efficient adjustment of hydrolytic extracellular enzyme production to changes in substrate availability and to potentially changing requirements.

4.4 Effects of temperature on bacterial activity

Temperature is an extensively investigated constraint on bacterial activity in polar marine systems that has a high potential to change metabolic rates (e.g. Rivkin et al., 1996; Pomeroy and Wiebe, 2001; Kirchman et al., 2009a). In the present study, in situ temperature increased over time and enhanced the daily production of bacterial biomass in the mesocosms substantially. The daily values for Q_{10} , the most common measure for rate acceleration induced by temperature increase, varied significantly from values ≤ 1 up to 12 over the duration of the experiment (Fig. 9). Q_{10} values ≤ 2 are characteristic for psychrophilic bacteria in polar waters that show low-temperature

BGD

9, 10467–10511, 2012

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



optima for metabolic processes, while higher Q_{10} values were observed for psychro-
tolerant bacterial strains that can cope with low seawater temperatures but show tem-
perature optima around 20 °C (Morita, 1975). Hence, maximum Q_{10} values of 12 sug-
gests that the community metabolism in the mesocosms was mainly driven by the
activity of psychrotolerant instead of psychrophilic bacterial strains. Q_{10} values for
 β -glucosidase, leu-aminopeptidase, and BPP were related to rates of phytoplankton
exudation, strongly suggesting interactions of bacterial temperature sensitivity and sub-
strate availability. Low Q_{10} values for bacterial activity at elevated rates of phytoplankton
exudation reveal lower temperature sensitivity at high substrate availability. A similar re-
lationship between ambient substrate concentration and temperature sensitivity of bac-
terial permeases was determined in an Arctic polynya. Here, Q_{10} values up to 11 were
determined for bacterial amino acid uptake at amino acid concentrations $\leq 50 \text{ nmol l}^{-1}$,
while Q_{10} decreased at higher amino acid concentrations (Yager et al., 1999). The dif-
ferences in Q_{10} determined in our study cannot be induced by direct substrate–enzyme
interactions, since the fluorescent and radioactive markers for enzyme and BPP mea-
surements, respectively, were added at saturating concentrations. It is likely that the
variability in Q_{10} is related to metabolic adaptations of the bacterioplankton community
to different levels of substrate input by phytoplankton production. These adaptations to
different substrate concentrations can be achieved by the use of different isoenzymes.
Isoenzymes catalyze the same reaction but can differ with respect to other kinetic
and biochemical characteristics (Ferenci, 1996; Wick et al., 2001). The co-existence
of isoenzymes in natural marine assemblages was shown by biphasic and multiphasic
kinetics of extracellular enzyme activity and bacterial substrate uptake and by the iso-
lation of diverse isoenzymes for extracellular polysaccharide hydrolysis during a bloom
situation (Unanue et al., 1999; Tholosan et al., 1999; Arrieta and Herndl, 2002). Fur-
thermore, laboratory studies revealed that extracellular isoenzymes produced by polar
marine bacteria can show different optimum temperatures for their catalytic reaction
(Nichols et al., 1999). Even isoenzymes produced by a single bacterial strain can differ
with regard to their temperature characteristics (Maki et al., 2006). Hence, a modulation

**Results from a
mesocosm
perturbation study**

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of the community metabolism by isoenzymes that are produced in response to different levels of substrate availability and show different temperature sensitivities could explain short-term variability in Q_{10} values for bacterial activity in natural communities.

4.5 Bacterial activity under elevated $p\text{CO}_2$

5 The general dependence of enzymatic reactions on pH is well-known for more than one hundred years and was confirmed for extracellular enzymes of aquatic habitats about 20 yr ago (Arrhenius, 1889; Münster, 1991). Also extracellular enzyme activity of cold-adapted marine isolates was shown to respond with characteristic optimum curves to a pH range from 2 to 12 (Groudieva et al., 2004). Recent studies revealed
10 that also moderate decreases in seawater pH as projected for the near future can significantly impact bulk rates of hydrolytic extracellular enzymes produced from natural marine bacterioplankton communities (Grossart et al., 2006; Tanaka et al., 2008, Piontek et al., 2010; Yamada and Suzumura, 2010). Most of these studies show elevated activity of different extracellular enzymes in response to seawater acidification. A meta-
15 analysis of ocean acidification effects on microbially driven biogeochemical processes has shown that effects on extracellular enzyme activity are among the most sensitive ones (Liu et al., 2011). During our mesocosm study, the direct response of the natural enzyme assemblages to lowered seawater pH was tested by an acidification assay with dilute hydrochloric acid. A decrease of 0.5 pH units roughly doubled rates
20 of β -glucosidase and leu-aminopeptidase in the Kongsfjorden samples and strongly suggests direct biochemical effects of lowered seawater pH on the enzymatic reaction velocities (Fig. 12). A comparison with the mean Q_{10} values for β -glucosidase and leu-aminopeptidase activity in the Kongsfjorden reveals that the increase in rates after moderate acidification equals the rate increase at a temperature elevated by 3°C.
25 Therefore, our results indicate that future changes in seawater temperature and pH have a similar potential to increase the hydrolytic activity of extracellular enzymes in this fjord system. Also extracellular enzyme activities in the high- CO_2 mesocosms were enhanced and directly related to the rising hydrogen ion concentrations over the

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



applied $p\text{CO}_2$ gradient (Fig. 10). Hence, results of both acidification assay and mesocosm study confirm an acceleration of organic matter-degrading extracellular enzyme activity at seawater pH projected for the near future. Enhanced extracellular enzyme activity in the mesocosms, however, cannot be exclusively attributed to direct biochemical pH effects. Instead, bacterial activity in the mesocosms was additionally stimulated by increased phytoplankton production under elevated $p\text{CO}_2$. Primary production determines the input of labile organic substrates for bacterial utilization. A positive linear correlation of primary production with rates of leu-aminopeptidase strongly suggests that an increased supply of organic compounds to bacterioplankton under elevated $p\text{CO}_2$ stimulated the cellular production of protein-degrading extracellular enzymes. Furthermore, a direct relationship between primary production and BPP shows that enhanced primary production under elevated $p\text{CO}_2$ enhanced not only the hydrolysis of organic matter but also its heterotrophic turnover. Thus, our results reveal that the bacterioplankton community of the Kongsfjorden had the potential to efficiently degrade surplus organic substrates derived from increased autotrophic carbon fixation under elevated $p\text{CO}_2$.

In the present mesocosm study lower bacterial abundances developed in the high- CO_2 mesocosms after day 20 due to higher viral lysis rates (Brussaard et al., 2012). This top-down regulation of bacterial cell numbers and thus enzyme producers strongly affected the bulk hydrolytic potential of β -glucosidase and leu-aminopeptidase in the mesocosms (Fig. 6). In accordance with the bottom-up regulation by seawater pH and primary production, high- CO_2 mesocosms showed a higher hydrolytic potential until day 20. This pattern changed and the trend of enhanced bulk hydrolysis under high $p\text{CO}_2$ was attenuated or even inverted when bacterial abundances diverged during the last phase of the experiment and top-down regulation led to lower bacterial cell numbers at high $p\text{CO}_2$ (Fig. 6; Brussaard et al., 2012). Also the community composition of bacterioplankton changed after day 20, showing a higher diversity of particle-attached bacteria in high- CO_2 mesocosms (Sperling et al., 2012). Therefore, also increased competitive relationships may have reduced community performance (Becker et al.,

BGD

9, 10467–10511, 2012

**Results from a
mesocosm
perturbation study**

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2012). The complex regulation of extracellular enzyme activity within natural plankton communities can partly explain why bulk enzymatic rates were affected to a varying degree under high $p\text{CO}_2$ in recent perturbation studies elsewhere (Liu et al., 2011; Grossart et al., 2006; Tanaka et al., 2008).

5 Concluding remarks

The present study investigates how elevated seawater $p\text{CO}_2$ can impact the metabolic activity of a natural bacterioplankton community in an Arctic fjord system. A short lag phase of extracellular enzyme activities and BPP after increasing phytoplankton production revealed high flexibility and pulse-responsiveness of two bacterial processes that substantially contribute to the heterotrophic carbon recycling and nutrient remineralization. Changes in seawater $p\text{CO}_2$ induced beneficial effects on bacterial activity by two principal mechanisms. Firstly, acidification increased rates of extracellular β -glucosidase and leu-aminopeptidase, strongly suggesting direct pH effects to enhance extracellular hydrolytic activity under elevated $p\text{CO}_2$. Secondly, BPP and leu-aminopeptidase activity were directly related to primary production at changing $p\text{CO}_2$. This suggests that elevated phytoplankton production under high CO_2 , as observed in the present mesocosm experiment and in previous CO_2 -perturbation studies (e.g. Egge et al., 2009), is efficiently channelled into the microbial food web in natural plankton communities. Therefore, enhanced heterotrophic carbon utilization may counterbalance increased autotrophic carbon fixation under high CO_2 and prevent enhanced carbon export. However, the top-down control of bacterial abundances led to higher cell losses at high CO_2 in the present mesocosm study and partly counteracted effects of lowered pH and enhanced primary production on the bulk hydrolysis of polysaccharides and proteins. Our results strongly suggest further investigation of acidification effects on both the biochemical sensitivity of bacterial isoenzymes and the regulation of bacterial activity in natural communities to better evaluate impacts of climate change on heterotrophic microorganisms and their feedback potential to ocean acidification.

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. This work is a contribution to the “European Project on Ocean Acidification” (EPOCA) which received funding from the European Community’s Seventh Framework Programme (FP7/2007-2013) under grant agreement no. 211384. We gratefully acknowledge the logistical support of Greenpeace International for its assistance with the transport of the mesocosm facility from Kiel to Ny-Ålesund and back to Kiel. We also thank the captains and crews of M/V ESPERANZA of Greenpeace and R/V Viking Explorer of the University Centre in Svalbard (UNIS) for assistance during mesocosm transport and during deployment and recovery in Kongsfjorden. We thank the staff of the French-German Arctic Research Base at Ny-Ålesund, in particular Marcus Schuhmacher, for on-site logistical support. Sebastian Krug and Nicole Händel are gratefully acknowledged for their support during the experiment.

The service charges for this open access publication have been covered by a Research Centre of the Helmholtz Association.

References

- Anderson, L. G.: DOC in the Arctic ocean, in: Biogeochemistry of Marine Dissolved Organic Matter, edited by: Hansell, D. A. and Carlson, C. A., Elsevier, 665–683, 2002.
- Arrhenius, S.: Über die Reaktionsgeschwindigkeit bei der Inversion von Rohrzucker durch Säuren, *Z. Phys. Chem.*, 4, 226–248, 1889.
- Arrieta, J. M. and Herndl, G. J.: Changes in bacterial β -glucosidase diversity during a coastal phytoplankton bloom, *Limnol. Oceanogr.*, 47, 549–599, 2002.
- Baltar, F., Arístegui, J., Sintes, E., van Aken, H. M., Gasol, J. M., and Herndl, G. J.: Prokaryotic extracellular enzymatic activity in relation to biomass production and respiration in the mesopelagic and bathypelagic waters of the (sub)tropical Atlantic, *Environ. Microbiol.*, 11, 1998–2014, 2009.
- Becker, J., Eisenhauer, N., Scheu, S., and Jousset, A.: Increasing antagonistic interactions cause bacterial communities to collapse at high diversity, *Ecol. Lett.*, 15, 468–474, 2012.
- Billen, G. and Becquevorts, S.: Phytoplankton-bacterial relationship in the Antarctic Marine Ecosystem, *Polar Res.*, 10, 245–253, 1991.

Results from a mesocosm perturbation study

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Results from a
mesocosm
perturbation study**

J. Piontek et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Bird, D. F. and Karl, D. M.: Uncoupling of bacteria and phytoplankton during the austral spring bloom in Gerlache Strait, Antarctic Peninsula, *Aquat. Microb. Ecol.*, 19, 13–27, 1999.
- Brussaard, C. P. D., Noordeloos, A. A. M., Witte, H., Collenteur, M., Schulz, K. G., Bellerby, R. G. J., Ludwig, A., Czerny, J., and Riebesell, U.: Arctic microbial community dynamics influenced by elevated CO₂ levels, *Biogeosciences*, under review, 2012.
- Cuevas, L. A., Egge, J., Thingstad, T. F., and Töpper, B.: Organic carbon and mineral nutrient limitation of oxygen consumption, bacterial growth and efficiency in the Norwegian Sea, *Polar Biol.*, 34, 871–882, 2011.
- Deming, J. W.: Psychrophiles and polar regions, *Curr. Opin. Microbiol.*, 5, 301–309, 2002.
- Ducklow, H. W.: The bacterial component of the oceanic euphotic zone, *FEMS Microbiol. Ecol.*, 30, 1–10, 1999.
- Ducklow, H. W. and Yager, P. L.: Pelagic bacterial processes in polynyas, in: *Polynyas: Windows into Polar Oceans*, edited by: Smith Jr., W. O. and Barber, D. G., Elsevier, 1–33, 2006.
- Ducklow, H. W., Kirchman, D. L., Quinby, H. L., Carlson, C. A., and Dam, H. G.: Stocks and dynamics of bacterioplankton carbon during the spring bloom in the Eastern North Atlantic Ocean, *Deep-Sea Res. Pt. II*, 40, 245–263, 1993.
- Dexter, S. C. and Chandrasekaran, P.: Direct measurement of pH within marine biofilms on passive metals, *Biofouling*, 4, 313–325, 2000.
- Dittmar, T. and Kattner, G.: The biogeochemistry of the river and shelf ecosystem of the Arctic ocean: a review, *Mar. Chem.*, 83, 103–120, 2003.
- Egge, J. K., Thingstad, T. F., Larsen, A., Engel, A., Wohlers, J., Bellerby, R. G. J., and Riebesell, U.: Primary production during nutrient-induced blooms at elevated CO₂ concentrations, *Biogeosciences*, 6, 877–885, doi:10.5194/bg-6-877-2009, 2009.
- Engel, A., Borchard, C., Piontek, J., Riebesell, U., and Schulz, K.: CO₂ increases ¹⁴C-primary production in an arctic plankton community, *Biogeosciences*, under review, 2012.
- Ferenci, T.: Adaptation to life at micromolar nutrient levels: the regulation of *Escherichia coli* glucose transport by endoinduction and cAMP, *FEMS Microbiol. Rev.*, 18, 301–317, 1996.
- Gargas, E.: A manual for phytoplankton primary production studies in the Baltic, *Balt. Mar. Biolog. Publ.*, 2, 1–88, 1975.
- Granéli, W., Carlsson, P., and Bertilsson, S.: Bacterial abundance, production and organic carbon limitation in the Southern Ocean (39–62° S, 4–14° E) during the austral summer 1997/1998, *Deep-Sea Res. Pt. II*, 51, 2569–2582, 2004.

- Groudieva, T., Kambourova, M., Yusef, H., Royter, M., Grote, R., Trinks, H., and Antranikian, G.: Diversity and cold-active hydrolytic enzymes of culturable bacteria associated with Arctic sea ice, *Spitzbergen, Extremophiles*, 8, 475–488, 2004.
- Grossart, H. P., Allgaier, M., Passow, U., and Riebesell, U.: Testing the effect of CO₂ concentration on the dynamics of marine heterotrophic bacterioplankton, *Limnol. Oceanogr.*, 51, 1–11, 2006.
- Hansell, D. A., Kadko, D., and Bates, N. B.: Degradation of terrigenous dissolved organic carbon in the Western Arctic Ocean, *Science*, 304, 858–861, 2004.
- Hop, H., Pearson, T., Hegseth, E. N., Kovacs, K. M., Wiencke, C., Kwasniewski, S., Eiane, K., Mehlum, F., Gulliksen, B., Wlodarska-Kowalczyk, M., Lydersen, C., Weslawski, J. M., Cochrane, S., Gabrielsen, G. W., Leaky, R. J. G., Lønne, O. J., Zajaczkowski, M., Falk-Petersen, S., Kendall, M., Wängberg, S.-Å., Bischof, K., Voronkov, A. Y., Kovaltchouk, N. A., Wiktor, J., Poltermann, M., di Prisco, G., Papucci, C., and Gerland, S.: The marine ecosystem of Kongsfjorden, Svalbard, *Polar Res.*, 21, 167–208, 2002.
- Hop, H., Falk-Petersen, S., Svendsen, H., Kwasniewski, S., Pavlov, V., Pavlova, O., and Søreide, J. E.: Physical and biological characteristics of the pelagic system across Fram Strait to Kongsfjorden, *Prog. Oceanogr.*, 71, 182–231, 2006.
- Hoppe, H.-G.: Significance of exoenzymatic activities in the ecology of brackish water – measurements by means of methylumbelliferyl-substrates, *Mar. Ecol.-Prog. Ser.*, 11, 299–308, 1983.
- Hoppe, H.-G.: Phosphatase activity in the sea, *Hydrobiologia*, 493, 187–200, 2003.
- Huston, A. L., Methe, B., and Deming, J. W.: Purification, characterization, and sequencing of an extracellular cold-active aminopeptidase produced by marine psychrophile *Colwellia psychrerythraea* strain 34H, *Appl. Environ. Microbiol.*, 70, 3321–3328, 2004.
- Iversen, K. R. and Seuthe, L.: Seasonal microbial processes in a high-latitude fjord (Kongsfjorden, Svalbard): I. heterotrophic bacteria, picoplankton and nanoflagellates, *Polar Biol.*, 34, 731–749, 2011.
- Karner, M., Ferrier-Pagès, C., and Rassoulzadegan, F.: Phagotrophic nanoflagellates contribute to occurrence of α -glucosidase and aminopeptidase in marine environments, *Mar. Ecol.-Prog. Ser.*, 114, 237–244, 1994.
- Kirchman, D. L.: Measuring bacterial biomass production and growth rates from leucine incorporation in natural aquatic environments, in: *Methods in Microbiology*, vol. 30, edited by: Paul, J. H., Academic Press, London, 227–238, 2001.

**Results from a
mesocosm
perturbation study**

J. Piontek et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Results from a
mesocosm
perturbation study**J. Piontek et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Kirchman, D. L., Malmstrom, R. R., and Cottrell, M. T.: Control of bacterial growth by temperature and organic matter in the Western Arctic, *Deep-Sea Res. Pt. II*, 52, 3386–3395, 2005.
- Kirchman, D. L., Morán, X. A. G., and Ducklow, H.: Microbial growth in the polar oceans – role of temperature and potential impact of climate change, *Nat. Rev. Microbiol.*, 7, 451–459, 2009a.
- 5 Kirchman, D. L., Hill, V., Cottrell, M. T., Gradinger, R., Malmstrom, R. R., and Parker, A.: Standing stocks, production, and respiration of phytoplankton and heterotrophic bacteria in the Western Arctic Ocean, *Deep-Sea Res. Pt. II*, 56, 1237–1248, 2009b.
- Kolber, Z.: Energy cycle in the ocean: powering the microbial world, *Oceanography*, 20, 79–88, 2007.
- 10 Liu, J., Weinbauer, M. G., Maier, C., Dai, M., and Gattuso, J.-P.: Effect of ocean acidification on microbial diversity and on microbe-driven biogeochemistry and ecosystem functioning, *Aquat. Microb. Ecol.*, 61, 291–305, 2010.
- Maki, S., Yoneta, M., and Takada, Y.: Two isocitrate dehydrogenases from a psychrophilic bacterium, *Colwellia psychrerythaea*, *Extremophiles*, 10, 237–249, 2006.
- 15 Meon, B. and Amon, R. M. W.: Heterotrophic bacterial activity and fluxes of dissolved free amino acids and glucose in the Arctic rivers Ob, Yenisei and the adjacent Kara Sea, *Aquat. Microb. Ecol.*, 37, 121–135, 2004.
- Nausch, M., Pollehne, F., and Kerstan, E.: Extracellular enzyme activities in relation to hydrodynamics in the Pomeranian Bight (Southern Baltic Sea), *Microb. Ecol.*, 36, 251–258, 1998.
- 20 Nichols, D. S., Greenhill, A. R., Shadbolt, C. T., Ross, T., and McMeekin, T. A.: Physicochemical parameters for growth of the sea ice bacteria *Glaciecola punicea* ACAM 611^T and *Gelidibacter* sp. Strain IC158, *Appl. Environ. Microbiol.*, 65, 3757–3760, 1999.
- Overpeck, J., Hughen, K., Hardy, D., Bradley, R., Case, R., Douglas, M., Finney, B., Gajewski, K., Jacoby, G., Jennings, A., Lamoureux, S., Lasca, A., MacDonald, G., Moore, J., Retelle, M., Smith, S., Wolfe, A., and Zielinski, G.: Arctic Environmental Change of the Last Four Centuries, *Science*, 1251–1256, 1997.
- Piontek, J., Lunau, M., Händel, N., Borchard, C., Wurst, M., and Engel, A.: Acidification increases microbial polysaccharide degradation in the ocean, *Biogeosciences*, 7, 1615–1624, doi:10.5194/bg-7-1615-2010, 2010.
- 30 Ploug, H., Kühl, M., Buchholz-Cleven, B., and Jørgensen, B. B.: Anoxic aggregates – an ephemeral phenomenon in the pelagic environment?, *Aquat. Microb. Ecol.*, 13, 285–294, 1997.

**Results from a
mesocosm
perturbation study**J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Pomeroy, L. R. and Deibel, D.: Temperature regulation of bacterial activity during the spring bloom in Newfoundland coastal waters, *Science*, 233, 359–361, 1986.
- Pomeroy, L. R. and Wiebe, W. J.: Temperature and substrates as interactive limiting factors for marine heterotrophic bacteria, *Aquat. Microb. Ecol.*, 23, 187–204, 2001.
- 5 Rich, J., Gosselin, M., Sherr, E., Sherr, B., and Kirchman, D. L.: High bacterial production, uptake and concentration of dissolved organic matter in the Central Arctic Ocean, *Deep-Sea Res. Pt. II*, 44, 1645–1663, 1997.
- Riebesell, U., v. Bröckel, K., Büdenbender, J., Czerny, J., Deckelnick, M., Hoffmann, D., Lenz, U., Ludwig, A., and Schulz, K. G.: A mobile, free-floating mesocosm platform for in situ experimentation, *Biogeosciences*, under review, 2012.
- 10 Rivkin, R. B., Anderson, M. R., and Lajzerowicz, C.: Microbial processes in cold oceans. I. Relationship between temperature and bacterial growth rate, *Aquat. Microb. Ecol.*, 10, 243–254, 1996.
- Sala, M. M., Arrieta, J. M., Boras, J. A., Duarte, C. M., and Vaqué, D.: The impact of ice melting on bacterioplankton in the Arctic Ocean, *Polar Biol.*, 33, 1683–1694, 2010.
- 15 Schulz, K. G., Bellerby, R. G. J., Brussaard, C., Büdenbender, J., Czerny, J., Fischer, M., Koch-Klavsen, S., Krug, S., Lischka, S., Ludwig, A., Meyerhöfer, M., Nondal, G., Silyakova, A., Stuhr, A. and Riebesell, U.: Temporal biomass dynamics of an Arctic plankton bloom in response to increasing levels of atmospheric carbon dioxide, *Biogeosciences*, under review, 2012.
- 20 Seuthe, L., Iversen, R. K., and Narcy, F.: Microbial processes in a high-latitude fjord (Kongsfjorden, Svalbard): II. Ciliates and dinoflagellates, *Polar Biol.*, 34, 751–766, 2011.
- Silyakowa, A., Bellerby, R. G. J., Nondal, G., Czerny, J., Schulz, K. G., Engel, A., Tanaka, T., De Lange, T., and Riebesell, U.: Net community production and stoichiometry of nutrient consumption in a pelagic ecosystem of a northern high latitude fjord: mesocosm CO₂ perturbation study, *Biogeosciences*, under review, 2012.
- 25 Simon, M. and Azam, F.: Protein content and protein synthesis rates of planktonic marine bacteria, *Mar. Ecol.-Prog. Ser.*, 51, 201–213, 1989.
- Simon, M., Glöckner, F. O., and Amann, R.: Different community structure and temperature optima of heterotrophic picoplankton in various regions of the Southern Ocean, *Aquat. Microb. Ecol.*, 18, 275–284, 1999.
- 30 Smith, D. C. and Azam, F.: A simple, economical method for measuring bacterial protein synthesis rates in seawater using ³H-leucine, *Mar. Microb. Food Webs*, 6, 107–114, 1992.

**Results from a
mesocosm
perturbation study**

J. Piontek et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sperling, M., Piontek, J., Gerdts, G., Wichels, A., Schunck, H., Roy, A.-S., La Roche, J., Gilbert, J., Bittner, L., Romac, S., Riebesell, U., and Engel, A.: Effect of elevated CO₂ on the dynamics of particle attached and free living bacteria in a natural bacterioplankton community in an arctic fjord, *Biogeosciences*, under review, 2012.

5 Steemann Nielsen, E.: The use of radioactive carbon (¹⁴C) for measuring primary production in the sea, *J. Cons. Perm. Int. Explor. Mer*, 18, 117–140, 1952.

Steinacher, M., Joos, F., Frölicher, T. L., Plattner, G.-K., and Doney, S. C.: Imminent ocean acidification in the Arctic projected with the NCAR global coupled carbon cycle-climate model, *Biogeosciences*, 6, 515–533, doi:10.5194/bg-6-515-2009, 2009.

10 Tanaka, T., Thingstad, T. F., Løvdal, T., Grossart, H.-P., Larsen, A., Allgaier, M., Meyerhöfer, M., Schulz, K. G., Wohlers, J., Zöllner, E., and Riebesell, U.: Availability of phosphate for phytoplankton and bacteria and of glucose for bacteria at different pCO₂ levels in a mesocosm study, *Biogeosciences*, 5, 669–678, doi:10.5194/bg-5-669-2008, 2008.

Tholosan, O., Lamy, F., Garcin, J., Polychronaki, T., and Bianchi, A.: Biphase extracellular proteolytic enzyme activity in benthic water and sediment in the Northwestern Mediterranean Sea, *Appl. Environ. Microbiol.*, 65, 1619–1626, 1999.

Unanue, M., Ayo, B., Agis, M., Slezak, D., Herndl, G. J., and Iriberry, J.: Ecto enzymatic activity and uptake of monomers in marine bacterioplankton described by a biphase kinetic model, *Microb. Ecol.*, 37, 36–48, 1999.

20 Vadstein, O.: Large variation in growth-limiting factors for marine heterotrophic bacteria in the Arctic waters of Spitsbergen (78° N), *Aquat. Microb. Ecol.*, 63, 289–297, 2011.

Wick, L. M., Quadroni, M., and Egli, T.: Short- and long-term changes in proteome composition and kinetic properties in a culture of *Escherichia coli* during transition from glucose-excess to glucose-limited growth conditions in continuous culture and vice versa, *Environ. Microbiol.*, 3, 588–599, 2001.

25 Yager, P. L. and Deming, J. W.: Pelagic microbial activity in an arctic polynya: testing for temperature and substrate interactions using a kinetic approach, *Limnol. Oceanogr.*, 44, 1882–1893, 1999.

Yager, P. L., Connely, T. L., Mortazavi, B., Wommack, K. E., Bano, N., Bauer, J. E., Opsahl, S., and Hollibaugh, J. T.: Dynamic bacterial and viral response to an algal bloom at subzero temperature, *Limnol. Oceanogr.*, 46, 790–801, 2001.

30 Yamada, N. and Suzumura, M.: Effects of seawater acidification on hydrolytic enzyme activities, *J. Oceanogr.*, 66, 233–241, 2010.

Results from a mesocosm perturbation study

J. Piontek et al.

Table 1. Half-saturation constant K_m for β -glucosidase and leucine-aminopeptidase activity in mesocosms and Kongsfjorden samples.

	β -glucosidase		leu-aminopeptidase	
	mesocosms	fjord	mesocosms	fjord
mean K_m (pmol l^{-1})	94	52	54	50
median K_m (pmol l^{-1})	50	34	51	40
range (pmol l^{-1})	3–696	3–252	2–188	19–126

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Results from a mesocosm perturbation study

J. Piontek et al.




Fig. 1. Bacterial growth characteristics (upper panel; BPP: bacterial protein production) and extracellular enzyme activities (lower panel) in samples of the Kongsfjorden with and without amendments of ammonium (NH₄) and glucose (glu) (t₀: initial sampling, t₄: sampling after 4 days of incubation).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Results from a mesocosm perturbation study

J. Piontek et al.




Fig. 2. Temporal development of bacterial protein production ($BPP_{2^{\circ}C}$) and specific growth rates (d^{-1}) at $2^{\circ}C$ in mesocosms of low ($175\text{--}250\ \mu atm$; blue symbols), medium ($340\text{--}600\ \mu atm$; grey symbols), and high pCO_2 ($675\text{--}1085\ \mu atm$; red symbols). $BPP_{2^{\circ}C}$ is normalized to volume ($ng\ C\ l^{-1}\ h^{-1}$; upper panel) and bacterial abundance ($fmol\ C\ cell^{-1}\ d^{-1}$; lower panel). Green lines represent chlorophyll *a* (Chl *a*) concentrations in the mesocosms.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Results from a mesocosm perturbation study

J. Piontek et al.




Fig. 3. Temporal development of bacterial production and growth (BPP_{2°C}, specific growth rate; upper panel) and extracellular enzyme activities (leu-aminopeptidase, β-glucosidase, phosphatase; lower panel) in the Kongsfjorden.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Results from a mesocosm perturbation study

J. Piontek et al.




Fig. 4. Temporal development of extracellular β -glucosidase activity in mesocosms of low (175–250 μatm ; blue symbols), medium (340–600 μatm ; grey symbols), and high $p\text{CO}_2$ (675–1085 μatm ; red symbols). Maximum reaction velocities (V_{\max}) are normalized to volume ($\text{nmol l}^{-1} \text{h}^{-1}$; upper panel) and bacterial abundance ($\text{amol cell}^{-1} \text{h}^{-1}$; lower panel). Green lines represent chlorophyll *a* (Chl *a*) concentrations in the mesocosms.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Results from a mesocosm perturbation study

J. Piontek et al.




Fig. 5. Temporal development of extracellular leucine (leu)-aminopeptidase activity in mesocosms of low (175–250 μatm ; blue symbols), medium (340–600 μatm ; grey symbols), and high $p\text{CO}_2$ (675–1085 μatm ; red symbols). Maximum reaction velocities (V_{max}) are normalized to volume ($\text{nmol l}^{-1} \text{h}^{-1}$; upper panel) and bacterial abundance ($\text{amol cell}^{-1} \text{h}^{-1}$; lower panel). Green lines represent chlorophyll *a* (Chl *a*) concentrations in the mesocosms.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 6. Temporal development of Δ hydrolysis potential for polysaccharide (upper panel) and protein degradation (lower panel) in acidified mesocosms. The shaded area represents the time period after the second chlorophyll *a* maximum when bacterial abundances in the mesocosms diverged (see text for further explanation).

**Results from a
mesocosm
perturbation study**J. Piontek et al.




Fig. 7. Activity ratios of leucine (leu)-aminopeptidase: β -glucosidase (upper panel) and molar ratios of DON : DOC (lower panel) in mesocosms (grey line) and Kongsfjorden (black circles). Mean values (\pm SD) of all mesocosms are shown.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Results from a mesocosm perturbation study

J. Piontek et al.




Fig. 8. Temporal development of extracellular phosphatase activity in mesocosms of low (175–250 µatm; blue symbols), medium (340–600 µatm; grey symbols), and high $p\text{CO}_2$ (675–1085 µatm; red symbols). Reaction velocities at 10 µmol l⁻¹ substrate concentration ($V_{10\mu\text{M}}$) are normalized to volume (nmol l⁻¹ h⁻¹). Green lines represent chlorophyll *a* (Chl *a*) concentrations in the mesocosms.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Results from a mesocosm perturbation study

J. Piontek et al.




Fig. 9. Q_{10} values for extracellular enzyme activities (β -glucosidase, leu-aminopeptidase) and bacterial protein production (BPP) in relation to phytoplankton exudation at different sampling days (e.g. d_{14} : day 14 of the experiment).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 10. Relationships between seawater $p\text{CO}_2$ /delta proton concentration ($\Delta [\text{H}^+]$) and extracellular enzyme activities. Separate correlations are shown for rates integrated over days 4–20 (open symbols) the whole experiment (filled symbols). Lines represent significant linear regressions.




Fig. 11. Relationships between primary production and bacterial protein production (BPP) (upper panel) and primary production and extracellular enzyme activities (diamonds: leu-aminopeptidase, squares: β -glucosidase) (lower panel). Black lines represent significant linear regressions.




Fig. 12. Activity of extracellular β -glucosidase and leu-aminopeptidase in samples of Kongsfjorden and one control mesocosm (M3) at in situ pH (fjord: 8.29, M3: 8.31) and after acidification with dilute hydrochloric acid (pH 7.79–8.17).

Results from a mesocosm perturbation study

J. Piontek et al.




Fig. 13. Pathways of direct pH effects and indirect $p\text{CO}_2$ effects on extracellular enzyme activity in the mesocosm plankton community. (1) Biochemical pH effect on enzymatic reaction velocity, (2) Effect of primary production (PP) on bacterial enzyme production, (3) Effect of PP on bacterial protein production (BPP), (4) Effect of bacterial abundance on bulk enzymatic activity. See text for further explanation.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
