Supplement of Clim. Past, 10, 1165–1182, 2014
<http://www.clim-past.net/10/1165/2014/>
doi:10.5194/cp-10-1165-2014-supplement
© Author(s) 2014. CC Attribution 3.0 License.

Supplement of

Impact of precession on the climate, vegetation and fire activity in southern Africa during MIS4

M.-N. Woillez et al.

Correspondence to: M.-N. Woillez (marienoelle.woillez@gmail.com)

FIG. 1 – Earth's orbital configuration for a) present-day, b) MIS4_max, 72 kyr BP, c) MIS4_min, 60 kyr BP, where e is the eccentricity, ϵ is the obliquity and ω is the longitude of the perihelion. WS stands for winter solstice, VE for vernal equinox, SS for summer solstice and AE for autumnal equinox (for the northern hemisphere).

FIG. 2 – Top : histogram of monthly temperature values over southern Africa for the present-day IPSL simulation (red) and the MIS4_min simulation (blue) (values interpolated to 0.16°). Bottom : spline depending on temperature used for the downscaling.

FIG. 3 – Observed annual burned area, average over 1997-2011 (as percentage of the area of a grid-cell). Data from the Global Fire Emissions Database (GFED, <http://globalfire-data.org>).

FIG. 4 – a) Mean air temperature anomaly and b) mean monthly precipitation anomaly simulated by IPSL_CM5A in MIS4_max compared to a present day simulation (MIS4_max - present).