An investigation of dust storms with some climatic factors in Iraq using ground and remote sensing measurements

Ali. A. Attiya^{1,2*}, Brian G. Jones²

¹*Atmospheric Science Department, Science Faculty, Mustansiriyah University, Iraq*

²*School of Earth and Environmental Sciences, University of Wollongong, NSW 2522, Australia*

*Corresponding author: Ali Attiya aaa800@uowmail.edu.au; ali_atmo78@yahoo.com

Orcid numbers: Ali Attiya - 0000-0001-9109-082X; Brian Jones - 0000-0003-0835-7197

Abstract

Dust storms are frequent in the desert and semi-desert areas including Iraq. This paper aims to examine the types of dust activity and its relationship with some meteorological factors using ground and remote sensing measurements. Dust types are classified into three types: floating/ suspended dust (visibility range > 10 km), blowing/ rising dust (visibility range between 1 and 10 km), and dust storms (visibility range below 1 km). The outcomes of this study revealed three periods for dust activity. The study showed three durations according to the recurrence of all dust activity: 1983-1992, 1993-2000 and 2001-2013. A negative correlation between annual mean precipitation and dust activity was detected. The relationship between precipitation with floating dust and blowing dust were increased in the south area of Iraq. In this area, the relationship coefficients between the annual mean of suspended and rising dust activity with precipitation were -0.13 and -0.39. Moreover, the relationship between the annual total suspended dust event and precipitation over Iraq were -0.298. Furthermore, a reduced direction in precipitation amounts and an elevated direction in air temperature averages through the study period. Any changes in temperature averages and precipitation mean during the long term are represented to be an indicator of climate change in a specific region. Thus, atmospheric factors examined consider necessary to get information on the past, present and future weather patterns of this given study region.

Keywords: Dust types, temperature, precipitation, satellite remote sensing, and atmospheric factors.

1. Introduction

Dust storms play a significant role in various fields at the global scale such as the economy, human health, environment and climate (Goudie, 2014; Kok *et al.*, 2018). Alluvial plains and dry areas (annual mean of precipitation at less than 200 mm and low vegetation) situated in low territories in considering major dust sources at global scale (Ginoux *et al.*, 2012).

Online International Conference on Atmospheric and Earth Sciences
November 05-06, 2021

Aeolian erosion includes the complex process of reactions between local or regional climate conditions, the atmospheric circulation and soil type characteristics (Wang *et al.*, 2015). The southwestern and central of Asia consist from deserts regions and these regions have a large probability to produce dust events (Ginoux *et al.*, 2012). The long aridity seasons lead to decrease vegetation, thus causing the Aeolian erosion and leading mobilization dust motivation and generating sever dust storms (Notaro *et al.*, 2015). Dust events can transport for long distance and they affect climates and human societies in the dry regions and remote regions from dust sources. There are areas in the southwestern and middle Asia faced increased in severe dust storms frequency during the recent years (Zoljoodi *et al.*, 2013). Dry and semi-dry lands in the central and southwestern Asia produce huge quantities of airborne mineral dust particles during the spring and summer seasons and these particles travelled for many thousands of kilometres per year by wind across various areas in the world such as central Asia and the Arabian Sea (Orlovsky *et al.*, 2013), China (Guan *et al.*, 2015) and Japan Sea (Irino and Tada, 2000). Climate observations from ground stations are applied to study and analysis dust events. There are studies employed the statistical methods to assess and analysis of dust events according to frequency and the seasonal variability in dry and semi-dry lands over Asia (Furman, 2003; Notaro *et al.*, 2015; Pi *et al.*, 2017). Furman (2003) suggested the Middle East region represents as one of important dust source areas after Africa desert (Sahara) and this study investigated in the temporal characteristics of dust phenomenon and its source areas in this region. The findings of this study revealed dust activities are high in the summer season over south-west parts of the Middle East while much recurrent during winter and spring above the north-west parts in this region as well as dust events occur through different times of the year. The study by (Attiya *et al.*, 2018) indicated that the southern and central regions Iraq (lower Mesopotamia) are considered as an important region for airborne dust plumes in the Middle East based on the analysis results of TOMS sensor data. Dust types classified according to horizontal visibility data based on the World Meteorology Organization (WMO) classification into 4 dust events, (a) Suspended Dust event (SD) with a visibility range of exceeding 10000 m, (b) Rising Dust event (RD) when visibility range below 10000 m, (c) Dust Storms event (DS) with visibility range below 1000 m and (d) Severe Dust Storms event (SDs) with visibility rang less than 200 m (Shao, 2008). There are little researches about different dust events analysis over the Middle East, particularly in Iraq, although the Middle East area represents one of the important dust source areas in the world. In recent years, Iraq suffered from several dust events yearly according to the Ministry of Environment in Iraq that indicated Iraq faced 122 intensity dust storms and 283 days of dust during the year as shown by (Varoujan *et al.*, 2013). The present study aims to analyse the dust events variability and its relationship with precipitation over Iraq, particularly above the south part of Iraq (Nasiriya), and its correlation to the annual average for air temperatures trends and precipitation rates between 1980 and 2015 utilizing the weather measurements by the Iraqi Meteorological Organization (IMO). This study aims to provide good information and better understanding for spatiotemporal variability about dust events in Iraq during this study period (35-years).

2. Methodology and datasets

The observations utilized for the analysis in this study contain climate data (number of days of dust storms, rising dust and suspended dust) from the Iraqi Meteorological Organization for seven climatic stations covering the period from 1980 to 2015 in Iraq. Monthly mean precipitation observations and a total number of dust events in Iraq and the south region (Nasiriya) were also acquired from the Iraqi Meteorological Organization for the period between 1980 and 2015. The correlation coefficients were created by a statistical program (SPSS) for the total annual number of dust types and the total annual precipitation, and for every dust type (overall Iraq and the south (Nasiriya) area separately). Some studies utilized horizontal visibility observations from climate stations to assess dust phenomenon in the Middle Eastern and Asia (Shao and Dong, 2006; Orlovsky *et al.*, 2013; Mashat *et al.*, 2017). The World Meteorological Organization (WMO) recognised weather events to 99 weather codes. Dust events have specific codes including suspended dust code (SD=06), rising dust code (RD=07), dust storms codes (DS=09 or 30, 31 and 32), and severe dust storms codes (SDS=33, 34 and 35; Shao, 2008). Table 1 shows dust events codes utilized in this research according to the Iraqi Meteorological Organisation (IMO).

Table 1: Showing synoptic weather codes of dust events applied in the present study.

No	Synoptic weather codes	Weather phenomenon characterization
1	6	Suspended Dust (SD)
2	7	Rising Dust (RD)
3	09 or 30, 31,32,33,34,35 and 98	Dust Storms and Severe Dust Storms (DS)

3. Results and discussion

3.1 Temporal changes in dust events

Figure 1 displays line charts (time series) of the decadal and yearly changes of total dust events during the period from 1980 to 2015. The horizontal line (orange) represents the mean (1028) computed for the full studied duration in the study region. Based on the change of the total number of dust days from the horizontal line (orange), the line charts in Figure 1 could be divided into three durations: 1983 to 1992 (D1) in which dust events were higher than the mean, 1993 to 2000 (D2) in which dust events were lower than the mean, and 2001 to 2012 (D3) in which dust event return to higher than the mean.

Figure 1: Annual total of all dust types. The solid orange line represents the mean (1028) over the long-period for the study region

War can lead to more dust events. Between 1983 and 1992, the total dust events were more than the mean. The increased dust events may be related to different wars which occurred during this period, such as the Iraqi-Iranian war in 1980-1988 and the Iraq - Kuwait war in 1991. At these times various heavy army equipment and vehicles were stationed in the dry and desert border areas between Iraq and the surrounding countries, as part of the planning and beginning of the war. Military operations and vehicle movements have led to terrestrial degradation of the surface thus increasing soil vulnerability that lead to a remarkable increase in dust mobilisation and transport in the air (El-Shobokshy and Al-Saedi, 1993). Between 1993 and 2000, the total frequency of airborne dust activities decreased to less than mean. From 2001, the total number of dust events then increased to more than the mean. This increase may have been caused by climate change, drought and desertification phenomena (Notaro *et al.*, 2015).

Figure 2 displays the time trend of suspended dust events. It shows a similar distribution to that of all dust types displayed in Figure 1. The interpretation used for all dust types is fully appropriate for suspended dust activity. This dust kind is mostly produced from local dust sources.

The time trend of rising dust activity (Figure 3) displays an almost a similar distribution to suspended dust, with a lower dust frequency. Rising dust clearly increased between 2001 and 2004 and then dropped below the mean after that. This remarkable increase may be referred to anthropogenic impacts due to the second Gulf war in 2003 between the Iraqi and American military forces. The time trend for dust storms (Figure 4) has a similar pattern to rising dust, while there is another increase in 2008. The sources of dust storms and rising dust are not

necessarily the same as the sources of suspended dust. Dust storms and rising dust are more possibly generated from regional and remote dust source areas.

Figure 2: Long time-pattern of total suspended dust days from 1980 to 2015 at study site; the orange line represents the long-term of the mean (649.5 days). The black line represents the annual number of activity; the green line represents the trend direction.

Figure 3: Long time-pattern of total rising dust days between 1980 and 2015. The horizontal line (orange) represents the long-time mean (329 days) at the study site, 1980-2015. The black line represents the annual number of activity; the green line represents the trend direction.

Figure 4: Long time-pattern of total dust storms days during 1980 and 2015. The horizontal line (orange) represents the long-time mean (329 days) at study site, 1980-2015. The black line represents the annual number of activity; the green line represents the trend direction.

3.2 The relationship between precipitation distribution and various dust types in Iraq

Figure 5 displays the number of dust days for every dust type with the annual precipitation distribution above Iraq and the annual precipitation distribution above the southern (Nasiriyah city) part of Iraq. The correlations coefficients were computed for every dust type using the statistical program (SPSS). With the annual precipitation over Iraq, the correlation (R) was slightly negative between precipitation and suspended dust (-0.298), rising dust (0.035), dust storms (-0.109) and all dust types (-0.22). Whereas, when considering the annual precipitation over the southern region (Nasiriya), the correlation was higher between precipitation and suspended dust (-0.39), rising dust (-0.13), dust storms (-0.22) and all dust types (-0.35).

Figure 5: The relationship between precipitation and type of dust activity above Iraq and the southern (Nasiriyah city) area in Iraq between 1980 and 2015

Online International Conference on Atmospheric and Earth Sciences
November 05-06, 2021

Figure 6: The annual total precipitation and air temperature mean for the whole of Iraq between 1980 and 2015. Note the increasing trend in air temperatures and decreasing trend in precipitation.

Environment and climate changes are likely contributed elements in the occurrence of dust storms and soil erosion process by active winds in the future (IPCC *et al.*, 2012), particularly through the strong winds occurrence, increase the desiccation periods in several regions (Dai, 2011). Any changes in temperature and precipitation rates over the three and a half decades represent an indicator of climatic change in a specific area (Elagib and Mansell, 2000; Moonen *et al.*, 2002). Thus, the climatic factors examined, such as temperatures and precipitation amounts, are necessary to obtain a knowledge of the past, present and future climate of a given area. Figure 6 displays the distribution of annual total precipitation and annual average air temperatures from 1980 to 2015. Overall, note the downward trend in precipitation rates and increasing trend in air temperatures.

4. Conclusion

In this current study, annual analyses of dust events have applied to define if there were any large changes in dust activity for study region to provide a general view. The present work investigated the correlation between the three dust events and rainfall rates in Iraq through the study period (35 years). The findings found that three remarkable periods that there was variability in total dust activity through the study period over Iraq. The correlation between total dust events and annual rainfall rates are negative. Also, the correlation between suspended dust event and total annual rainfall rates were negative, and the correlation between both rising dust, suspended dust events with rainfall rates are higher in the southern region (Nasiriya) in Iraq.

Acknowledgements

The researchers are thankful to University of Wollongong, Australia, the Bureau of meteorology in Iraq for supplying weather data.

References

- Attiya, A. A., Jones, B. G. & Marx, S. 2018, "Spatial-temporal analysis of Aerosol Index (AI) distribution and some climatic factors: case study from Iraq, 1980–2015", *Modeling Earth Systems and Environment*, 1-14, <https://doi.org/10.1007/s40808-018-0528-8>.
- Dai, A. 2011, "Drought under global warming: a review", *Wiley Interdisciplinary Reviews: Climate Change*, 2, 45-65.

Online International Conference on Atmospheric and Earth Sciences
November 05-06, 2021

- El-Shobokshy, M. S. & Al-Saedi, Y. G. 1993, "The impact of the gulf war on the Arabian environment—I. Particulate pollution and reduction of solar irradiance", *Atmospheric Environment. Part A. General Topics*, 27, 95-108.
- Elagib, N. A. & Mansell, M. G. 2000, "Recent trends and anomalies in mean seasonal and annual temperatures over Sudan", *Journal of Arid Environments*, 45, 263-288.
- Furman, H. K. H. 2003, "Dust storms in the Middle East: sources of origin and their temporal characteristics", *Indoor and Built Environment*, 12, 419-426.
- Ginoux, P., Prospero, J. M., Gill, T. E., Hsu, N. C. & Zhao, M. 2012, "Global-scale attribution of anthropogenic and natural dust sources and their emission rates based on MODIS Deep Blue aerosol products", *Reviews of Geophysics*, 50,
- Goudie, A. S. 2014, "Desert dust and human health disorders", *Environment International*, 63, 101-113.
- Guan, Q., Yang, J., Zhao, S., Pan, B., Liu, C., Zhang, D. & Wu, T. 2015, "Climatological analysis of dust storms in the area surrounding the Tengger Desert during 1960–2007", *Climate dynamics*, 45, 903-913.
- Ipcc, Oppenheimer, M., Diop, C., Hess, J., Lempert, R., Li, J. & Myeong, S. 2012, "Managing the risks of extreme events and disasters to advance climate change adaptation", *A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC)*, 25-64.
- Irino, T. & Tada, R. 2000, "Quantification of aeolian dust (Kosa) contribution to the Japan Sea sediments and its variation during the last 200 ky", *Geochemical Journal*, 34, 59-93.
- Kok, J. F., Ward, D. S., Mahowald, N. M. & Evan, A. T. 2018, "Global and regional importance of the direct dust-climate feedback", *Nature communications*, 9, 241.
- Mashat, A.-W. S., Alamoudi, A. O., Awad, A. M. & Assiri, M. E. 2017, "Synoptic characteristics of dusty spring days over central and eastern Saudi Arabia", *Air Quality, Atmosphere & Health*, 10, 307-323.
- Moonen, A., Ercoli, L., Mariotti, M. & Masoni, A. 2002, "Climate change in Italy indicated by agrometeorological indices over 122 years", *Agricultural and Forest Meteorology*, 111, 13-27.
- Notaro, M., Yu, Y. & Kalashnikova, O. V. 2015, "Regime shift in Arabian dust activity, triggered by persistent Fertile Crescent drought", *Journal of Geophysical Research: Atmospheres*, 120,
- Orlovsky, N., Orlovsky, L. & Indoitu, R. 2013, "Severe dust storms in Central Asia", *Arid ecosystems*, 3, 227-234.
- Pi, H., Sharratt, B. & Lei, J. 2017, "Atmospheric dust events in central Asia: Relationship to wind, soil type, and land use", *Journal of Geophysical Research: Atmospheres*, 122, 6652-6671.
- Shao, Y. 2008, *Physics and Modelling of wind erosion*, Springer Science & Business Media, Germany.
- Shao, Y. & Dong, C. 2006, "A review on East Asian dust storm climate, modelling and monitoring", *Global and Planetary Change*, 52, 1-22.
- Varoujan, K. S., Nadhir, A.-A. & Sven, K. 2013, "Sand and dust storm events in Iraq", *Natural Science*, 5, 1084-1094.
- Wang, H., Jia, X., Li, K. & Li, Y. 2015, "Horizontal wind erosion flux and potential dust emission in arid and semiarid regions of China: A major source area for East Asia dust storms", *Catena*, 133, 373-384.
- Zoljoodi, M., Didevarasl, A. & Saadatabadi, A. R. 2013, "Dust events in the western parts of Iran and the relationship with drought expansion over the dust-source areas in Iraq and Syria", *Atmospheric Climate and sciences*, 3, 321-336.