HARMONICITY OF (φ, φ') -HOLOMORPHIC SECTIONS OF A (SEMI-RIEMANNIAN) ALMOST PARACONTACT FIBER BUNDLE OF TYPE II

ARISTIDE AYIBE

ABSTRACT. In this paper (φ, φ') -holomorphic maps from an almost para-Hermitian manifold to an almost paracontact metric manifold are studied and a criterion for the harmonicity of such (φ, φ') -holomorphic maps is obtained.

Also (φ, φ') -holomorphic sections of (*semi-Riemannian*) almost paracontact fiber bundles of type II are studied and a criterion for the harmonicity of such (φ, φ') -holomorphic sections is obtained.

1. INTRODUCTION

A semi-Riemannian (or pseudo-Riemannian) manifold M is a C^∞ -manifold endowed with a symmetric non-degenerate $(0, 2)$ tensor field g on M with constant indices of positivity and negativity $ind_+ M$ and $ind_- M$, respectively. The non-degeneracy means that $ind_+ M + ind_- M = \dim M$ (the dimension of M).

As in the Riemannian's case, a C^2 -map $u : (M, g) \rightarrow (N, h)$ between semi-Riemannian manifolds is called harmonic, when the tension field $\tau(u)$ of u vanishes, where $\tau(u)$ is defined by $\tau(u) = \text{Trace} \nabla^* u$.

Let $(N \xrightarrow{\pi} M)$ be a fiber bundle. A section of $(N \xrightarrow{\pi} M)$ is a continuous map $u : M \rightarrow N$ such that $\pi \circ u = Id$.

Let (N, h) and (M, g) be two semi-Riemannian manifolds of index $s' (0 \leq s' \leq \dim N)$ and $s (0 \leq s \leq \dim M)$ respectively with $s \leq s'$ and $(N \xrightarrow{\pi} M)$ a fiber bundle.

We say that $[(N, h) \xrightarrow{\pi} (M, g)]$ is a semi-Riemannian fiber bundle, when for all $y \in N$, $du(y)|_{H_y} : H_y \rightarrow T_{\pi(y)} M$ is an isometry for the metric $h(y)$ on H_y and $g(\pi(y))$ on $T_{\pi(y)} M$, where $H_y = (T_y \pi^{-1}(\pi(y)))^\perp$ is the orthogonal complement of $T_y \pi^{-1}(\pi(y))$ and the fibers, $\pi^{-1}(x)$, for all $x \in M$ are semi-Riemannian submanifolds of N .

In other words a semi-Riemannian fiber bundle $[(N, h) \xrightarrow{\pi} (M, g)]$ is a fiber bundle, where (N, h) and (M, g) are two semi-Riemannian manifolds and π a semi-Riemannian submersion.

2010 *Mathematics Subject Classification.* 53C15; 53C20; 53C25; 58E20; 58Exx.

Key words and phrases. (semi-Riemannian) almost paracontact fiber bundle of type II, (φ, φ') -holomorphic sections.

©2021 Ilirias Research Institute, Prishtinë, Kosovë.

Submitted March 3, 2020. Published May 24, 2021.

Communicated by Guest editor Siraj Uddin.

Let us recall that semi-Riemannian submersions have been studied by many authors. (See e.g. [1] and [2])

As in the Riemannian's case (see [3], p.160), a C^2 -section $u : M \rightarrow N$ of a semi-Riemannian fiber bundle is called harmonic, when the vertical tension field $\tau^v(u)$ of u vanishes, where $\tau^v(u) = \text{Trace} \nabla^{*v} d^v u$.

An almost para-complex manifold (M, J) is a differentiable manifold M with a tensor J satisfying: for each $x \in M$, $J_x : T_x M \rightarrow T_x M$ such that : $J_x^2 = Id_{T_x M}$.

An almost para-Hermitian manifold (M, g, J) is an almost para-complex manifold (M, J) with a semi-Riemannian metric g satisfying: for all $X, Y \in \chi(M)$, $g(JX, JY) = -g(X, Y)$, where $\chi(M)$ is the set of vector fields on M .

An almost para-Hermitian manifold (M, J, g) is said to be:

- ★ para-Kähler if $\nabla J = 0$;
- ★ nearly para-Kähler if $(\nabla_X J)X = 0$, for all $X \in \chi(M)$;
- ★ almost para-Kähler if $d\Phi = 0$, where Φ is the para-Kähler 2-form defined by : for all $X, Y \in \chi(M)$, $\Phi(X, Y) = g(X, JY)$;
- ★ quasi-para-Kähler if $(\nabla_X J)Y - (\nabla_{JX} J)JY = 0$, for all $X, Y \in \chi(M)$;
- ★ almost semi-para-Kähler if $\delta J = 0$, where δ denotes the codifferential in (M, g) .

Let M be a differentiable manifold of dimension $2m+1$. An almost paracontact structure on M is a triple (φ, ξ, η) , where

ξ is a characteristic vector field,

η is a 1-form such that $\eta(\xi) = 1$. and

φ a tensor field of type $(1, 1)$ satisfying $\varphi^2 = I - \eta \otimes \xi$, where I stands for the identity endomorphism of the fiber bundle TM :

Then (M, φ, ξ, η) is called almost paracontact manifold. Furthermore $\varphi(\xi) = 0$ and $\eta \circ \varphi = 0$.

Let (M, φ, ξ, η) be almost paracontact manifold.

If M is equipped with a pseudo-Riemannian metric g satisfying :

$g(\varphi X, \varphi Y) = -g(X, Y) + \eta(X)\eta(Y)$, for all $X, Y \in \chi(M)$, then $(M, \varphi, \xi, \eta, g)$ is called almost para-contact metric manifold.

As in the case of almost contact metric manifolds, any almost paracontact metric manifold admits a fundamental 2-form, Φ defined by $\Phi(X, Y) = g(X, \varphi Y)$, for all $X, Y \in \chi(M)$.

An almost paracontact metric manifold $(M, \varphi, \xi, \eta, g)$ is said to be:

- ★ paracontact if $\Phi = d\eta$,
- ★ para-cosymplectic if $\nabla \Phi = 0$ and $\nabla \eta = 0$,
- ★ almost para-cosymplectic if $d\Phi = 0$ and $d\eta = 0$
- ★ nearly para-K-cosymplectic if $(\nabla_X \varphi)X = 0, \nabla_X \xi = 0$, for all $X \in \chi(M)$,
- ★ quasi-para-K-cosymplectic if $(\nabla_X \varphi)Y - (\nabla_{\varphi X} \varphi)\varphi Y + \eta(Y)\nabla_{\varphi X} \xi = 0$, for all $X, Y \in \chi(M)$,
- ★ semi para-cosymplectic if $\delta \Phi = 0 (\Rightarrow \delta \varphi = 0)$ and $\delta \eta = 0$.
- ★ semi para-Sasakian if $\eta = \frac{1}{2m} \delta \Phi$.
- ★ quasi-para-Sasakian if $d\Phi = 0$ and M is normal.
- ★ para-Sasakian if $(\nabla_X \varphi)Y = -g(X, Y)\xi + \eta(Y)X$, for all $X, Y \in \chi(M)$.

A C^2 -map $u : (M, \varphi) \rightarrow (N, \varphi', \xi', \eta')$ from an almost para-Hermitian manifold to an almost paracontact metric manifold. is said to be (φ, φ') -holomorphic, when $du\varphi = \varphi' du$.

A (semi - Riemannian) almost paracontact fiber bundle of type II,

$[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, g)]$ is a semi-Riemannian fiber bundle where, $(N, \varphi', \xi', \eta', h)$

is a paracontact metric manifold and (M, φ, g) is an almost para-Hermitian manifold and π is a (φ, φ') -holomorphic map.

In other words a *(semi - Riemannian)* almost paracontact fiber bundle of type II,

$[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, g)]$ is a fiber bundle where, $(N, \varphi', \xi', \eta', h)$ is a paracontact metric manifold and (M, φ, g) is an almost para-Hermitian manifold and π is a paracontact semi-Riemannian submersion of type II.

We recall that almost paracontact semi-Riemannian submersions of type II have been studied by Tshikuna-Matamba Tshikunguila, in [4] and by Ylmaz Gunduzalp and Bayram Sahin (in [5]), who call them para-contact para-complex semi-Riemannian submersions.

A section $u : M \rightarrow N$ of a *(semi - Riemannian)* almost paracontact fiber bundle of type II, $[(N, h, \varphi, \xi, \eta) \xrightarrow{\pi} (M, \varphi, g)]$ is said to be (φ, φ') -holomorphic when $d^v u \circ \varphi = \varphi' \circ d^v u$.

S. Ianus and A.M.Pastore in [6] proved that a (φ, φ') -holomorphic map between contact manifolds is harmonic.

In [7], Domingo China gives an expression of the tension field $\tau(u)$ of a (φ, φ') -holomorphic map $u : (M, \varphi, \xi, \eta, g) \rightarrow (N, \varphi', \xi', \eta', h)$ between almost contact metric manifolds. Namely he shows:

Theorem 1

Let $(M, \varphi, \xi, \eta, g)$ and $(N, \varphi', \xi', \eta', h)$ be two almost contact metric manifolds.

If $u : M \rightarrow N$ is a (φ, φ') -holomorphic map, then we have:

$$\begin{aligned} \tau(u) &= \varphi'(trac_g(du)^*(\nabla' \varphi')) - du(\varphi \delta \varphi + \delta \eta \xi) + [\nabla'_{du(\xi)}(\eta'(du(\xi)))] \\ &- trac_g(du)^*(\nabla' \eta') \xi, \end{aligned}$$

where $trac_g(du)^*(\nabla' \varphi') = \sum_{k=1}^{2m+1} (\nabla'_{du(e_k)} \varphi') du(e_k)$, and

$$trac_g u^*(\nabla' \eta') = \sum_{k=1}^{2m+1} (\nabla'_{du(e_k)} \eta') du(e_k),$$

with $\{e_1, \dots, e_{2m+1}\}$ an orthonormal basis of TM .

This allows him to generalize the above result of S. Ianus and A.M.Pastore in [5].

A *(Riemannian)* almost contact fiber bundle of type I, $[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, \xi, \eta, g)]$ is a Riemannian fiber bundle, where $(N, \varphi', \xi', \eta', h)$ and $(M, \varphi, \xi, \eta, g)$ are almost contact metric manifolds and π is a (φ, φ') -holomorphic map satisfying $d\pi(\xi) = \xi'$.

In [8] we extend the result of Domingo China to section of a *(Riemannian)* almost contact fiber bundle of type I.

Namely we show:

Proposition 2

Let $[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, \xi, \eta, g)]$ be a *(Riemannian)* almost contact fiber bundle of type I and $u : M \rightarrow N$ be a holomorphic section of $[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, \xi, \eta, g)]$. Then

$$\begin{aligned} \tau^v(u) &= \varphi'[trac_g(d^v u)^*(\nabla^v \varphi') + trac_g(\nabla_{d^h u}^v \varphi')(d^v u) + trac_g(A_{d^h u} d^h u \circ \varphi)] \\ &+ trac_g(T_{d^v u} d^h u \circ \varphi) + d^v u(\delta \varphi) - d^v u(\delta \eta \xi), \end{aligned}$$

where T and A are the fundamental tensors of a Riemannian fiber bundle, du is the differential of u and the superscripts h and v mean respectively the projection onto the horizontal, respectively the vertical bundle.

The aim of this work is to:

1) Give an equivalent of the result of Domingo Chinaea when the domain is an almost para-Hermitian manifold and the target, an almost paracontact metric manifold.

2) Extend (1) to (φ, φ') -holomorphic sections of a (*semi – Riemannian*) almost paracontact fiber bundle of type II.

We reach our aim by giving an expression of :

First the tension field $\tau(u)$ of a (φ, φ') -holomorphic map $u : (M, \varphi) \rightarrow (N, \varphi', \xi', \eta')$ from an almost para-Hermitian manifold to an almost paracontact metric manifold.

And secondly an expression of the vertical tension field $\tau^v(u)$ of a (φ, φ') -holomorphic section of a (*semi – Riemannian*) almost paracontact fiber bundle of type II.

The paper is organized as follows. In section II, we recall definitions and properties which are useful for the other sections. In section III, we give the results of the maps case. Section IV is devoted to proofs of the results of the maps case. In section V, we give the results of the sections case. Section VI is devoted to proofs of the results of sections case.

2. DEFINITIONS

2.1. Vertical and horizontal fiber bundles. Let $[(N, h) \xrightarrow{\pi} (M, g)]$ be a semi-Riemannian fiber bundle.

We define the vertical fiber bundle W to be the vector bundle over N which fibers are the vertical tangent spaces $T_y \pi^{-1}(\pi(y))$, for all $y \in N$.

Also we define the horizontal fiber bundle H to be the vector bundle over N which fibers are the horizontal tangent spaces H_y , for all $y \in N$.

2.2. Pull back bundle. Let $(B \xrightarrow{\pi} N)$ be a vector bundle, and $u : M \rightarrow N$ a smooth map, then there exists a unique vector bundle $(u^{-1}(B) \xrightarrow{\pi'} M)$ called pull-back bundle such that for all $x \in M$, $\pi'^{-1}(\{x\}) = B_{u(x)}$.

Let's assume that $(B \xrightarrow{\pi} N)$ is equipped with a semi-Riemannian metric h and a compatible covariant derivative ∇ . Then $(u^{-1}(B) \xrightarrow{\pi'} M)$ is equipped with a semi-Riemannian metric a and a compatible covariant derivative ∇^* defined respectively by:

$$a : M \rightarrow (u^{-1}(B))^* \otimes (u^{-1}(B))^*,$$

$$a_x(Y, Z) = h_{u(x)}(Y, Z), \forall x \in M, \forall Y, Z \in (u^{-1}(B))_x.$$

And

$$\nabla_X^*(wou) = \nabla_{du(X)} w, \forall X \in \chi(M), \forall w \in \Gamma(B).$$

2.3. Harmonic section. Let $u : M \rightarrow N$ be a C^2 section of a semi-Riemannian fiber bundle $[(N, h) \xrightarrow{\pi} (M, g)]$.

The vertical tension field of u is given by $\tau^v(u) = \text{Trace} \nabla^{*v} d^v u$.

If $\{e_1, \dots, e_{2m}\}$ is a local orthonormal frame in TM such that

$$\begin{aligned} g(e_k, e_k) &= 1, \forall 1 \leq k \leq m, \\ g(e_k, e_k) &= -1, \forall m+1 \leq k \leq 2m, \\ g(e_i, e_j) &= 0, \forall i \neq j, \end{aligned}$$

then

$$\tau^v(u) = \sum_{k=1}^m [\nabla_{e_k}^{*v} (d^v u(e_k)) - d^v u(\nabla_{e_k} e_k)] - \sum_{k=m+1}^{2m} [\nabla_{e_k}^{*v} (d^v u(e_k)) - d^v u(\nabla_{e_k} e_k)],$$

where ∇ denotes the Levi-Civita covariant derivative on M , ∇^v denotes the Levi-Civita covariant derivative on W induced by the metric h and ∇^{*v} the pull-back of the Levi-Civita covariant derivative ∇^v on W to the pull-back bundle $(u^{-1}(W) \rightarrow M)$.

We recall that the section u is called harmonic when $\tau^v(u) = 0$.

2.4. Fundamental tensors of a semi Riemannian fiber bundle.(See [2]).

Let $[(N, h) \xrightarrow{\pi} (M, g)]$ be a semi-Riemannian fiber bundle. The two fundamental tensors of π are defined by :

$$\begin{aligned} T_E F &= \nabla_{E^v}^h F^v + \nabla_{E^v}^v F^h \\ A_E F &= \nabla_{E^h}^h F^v + \nabla_{E^h}^v F^h \end{aligned}$$

for all vectors fields E and F on N , where the superscripts h and v mean respectively the projection onto the horizontal, respectively the vertical bundle.

Remark:

The tensor $A = 0$ if the horizontal distribution is integrable.

The tensor $T = 0$ if the fibers are totally geodesic.

2.5. Link between the covariant derivative of $Z \in \Gamma(u^{-1}(TN))$ and it's extension. Let $(N \xrightarrow{\pi} M)$ be a fiber bundle. $\Gamma(N)$ denotes the set of the sections of $(N \xrightarrow{\pi} M)$.

Let $Z \in \Gamma(u^{-1}(TN))$ and $(E_j)_{1 \leq \alpha \leq m}$ be an orthonormal local frame in TN defined on a open subset V of N , then for all $x_0 \in u^{-1}(V)$, $Z(x_0) = Z^j(x_0)E_j(u(x_0))$.

To Z we associate the vector field $\tilde{Z}$ of N defined on V by : for all $y \in V$, $\tilde{Z}(y) = Z^j(\pi(y))E_j(y)$.

On the one hand :

$$\begin{aligned} \forall X \in \chi(M), \forall x \in u^{-1}(V), \nabla_X^{*'} Z(x) &= \nabla_X^{*'} [Z^j(x)E_j(u(x))] \\ &= D_X(Z^j)(x)E_j(u(x)) + Z^j(x)\nabla_X^{*'} E_j(u(x)) \\ &= D_X(Z^j)(x)E_j(u(x)) + Z^j(x)\nabla'_{du(X)(x)} E_j. \end{aligned}$$

On the other hand with the fact that u is a section i.e. $\pi \circ u = Id_M$, one has :
 $\forall X \in \chi(M), \forall x \in u^{-1}(V)$,

$$\begin{aligned} (\nabla_X^{*'} \tilde{Z}ou)(x) &= \nabla'_X [(Z^j \circ \pi \circ u) E_j ou](x) \\ &= D_X (Z^j \circ \pi \circ u)(x) E_j ou(x) + Z^j \circ \pi \circ u(x) (\nabla_X^{*'} E_j ou)(x) \\ &= D_X (Z^j)(x) E_j ou(x) + Z^j(x) \nabla'_{du(X)(x)} E_j. \end{aligned}$$

It follows that :

$$\forall X \in \chi(M), \nabla_X^{*'} Z = \nabla_X^{*'} (\tilde{Z}ou).$$

Since :

for all $X \in \chi(M)$, $\nabla_X^{*'} (\tilde{Z}ou) = \nabla'_{du(X)} \tilde{Z}$ we get:

$$\forall X \in \chi(M), \nabla_X^{*'} Z = \nabla'_{du(X)} \tilde{Z}, \quad (2.1)$$

where ∇' denotes the Levi-Civita covariant derivative on N and $\nabla^{*'}$ the pull-back of ∇' to the pull-back bundle $(u^{-1}(TN) \rightarrow M)$.

2.6. Fundamental properties of a (semi-Riemannian) almost paracontact fiber bundle of type II. See Prop.3.2 and prop.3.3 in [4]). Let $[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, g)]$,

be a (semi – Riemannian) almost paracontact fiber bundle of type II. Then

$$\pi^* \Phi' = \Phi,$$

$$U \in \Gamma(W) \implies \varphi' U \in \Gamma(W),$$

$$Y \in \Gamma(H) \implies \varphi' Y \in \Gamma(H),$$

$$\xi' \in \Gamma(W),$$

$$\eta'(X) = 0, \text{ for all } X \in \Gamma(H),$$

$\varphi' X'$ is basic associated to φX , when X' is basic associated to X .

$(\nabla_X^h \varphi') Y'$ is basic associated to $(\nabla_X \varphi) Y$, when X' and Y' are basic associated respectively to X and Y .

2.7. Trace of some 2-covariant tensors. Let $[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, g)]$

be a (semi – Riemannian) almost paracontact fiber bundle of type II.

Let θ be a $u^{-1}(TN)$ -valued 2-covariant tensor on M .

As g is non-degenerate, there exists an ordered basis $(e_1, e_2, \dots, e_{2m})$ in TM such that:

$$g(e_i, e_i) = 1, \forall 1 \leq i \leq m,$$

$$g(e_i, e_i) = -1, \forall m+1 \leq i \leq 2m,$$

$$g(e_i, e_j) = 0, \forall i \neq j.$$

Let $\{(e_1)^*, \dots, (e_{2m})^*\}$ be its dual basis.

We call trace of θ , the section $trac_g(\theta)$ of $u^{-1}(TN)$ defined by:

$$trac_g(\theta) = \sum_{i,j=1}^{2m} g^{ij} \theta(e_i, e_j).$$

If $\{e_1, \dots, e_{2m}\}$ is orthonormal, then $trac_g(\theta) = \sum_{i=1}^m \theta(e_i, e_i) - \sum_{i=m+1}^{2m} \theta(e_i, e_i)$.

Like above let us assume that $\{X_1, \dots, X_m, \varphi X_1, \dots, \varphi X_m\}$ is an orthonormal local frame in TM .

The trace of the tensor $(d^v u)^*(\nabla^v \varphi')$ is given by:

$$\text{trac}_g(d^v u)^*(\nabla^v \varphi') = \sum_{k=1}^m (\nabla_{d^v u(X_k)}^v \varphi')(d^v u(\tilde{X}_k)) - \sum_{k=1}^m (\nabla_{d^v u(\varphi X_k)}^v \varphi')(d^v u(\tilde{\varphi X}_k)).$$

The trace of the tensor $(d^v u)^*(\nabla' \eta')$ is given by:

$$\text{trac}_g(d^v u)^*(\nabla' \eta') = \sum_{k=1}^m (\nabla_{d^v u(X_k)}' \eta')(d^v u(\tilde{X}_k)) - \sum_{k=1}^m (\nabla_{d^v u(\varphi X_k)}' \eta')(d^v u(\tilde{\varphi X}_k)).$$

The trace of the tensor $A_{d^h u} d^h u \circ \varphi$ is given by:

$$\text{trac}_g(A_{d^h u} d^h u \circ \varphi) = \sum_{k=1}^m [A_{d^h u(X_k)}(d^h u \circ \tilde{\varphi}(X_k)) - A_{d^h u(JX_k)}(d^h u \circ \tilde{\varphi}(X_k))].$$

The trace of the tensor $T_{d^v u} d^h u \circ \varphi$ is given by:

$$\text{trac}_g(T_{d^v u} d^h u \circ \varphi) = \sum_{k=1}^m [T_{d^v u(X_k)}(d^h u \circ \tilde{\varphi}(X_k)) - T_{d^v u(JX_k)}(d^h u \circ \tilde{\varphi}(X_k))].$$

The trace of the tensor field $(\nabla_{d^h u}^v \varphi')(d^v u)$ is given by:

$$\text{trac}_g(\nabla_{d^h u}^v \varphi')(d^v u) = \sum_{k=1}^m (\nabla_{d^h u(X_k)}^v \varphi')(d^v u(\tilde{X}_k)) - \sum_{k=1}^m (\nabla_{d^h u(\varphi X_k)}^v \varphi')(d^v u(\tilde{\varphi X}_k)).$$

The trace of the tensor field $(\nabla_{d^h u}^v \eta')(d^v u)$ is given by:

$$\text{trac}_g(\nabla_{d^h u}^v \eta')(d^v u) = \sum_{k=1}^m (\nabla_{d^h u(X_k)}^v \eta')(d^v u(\tilde{X}_k)) - \sum_{k=1}^m (\nabla_{d^h u(\varphi X_k)}^v \eta')(d^v u(\tilde{\varphi X}_k)).$$

3. RESULTS FOR MAPS

Proposition 3.1. : *Let $u : (M, \varphi, g) \rightarrow (N, \varphi', \xi', \eta', h)$ be a (φ, φ') -holomorphic map from an almost para-Hermitian manifold to almost paracontact metric manifold.*

Then the tension field of u is given by:

$$\tau(u) = -\varphi'[\text{trac}_g(du)^*(\nabla' \varphi') + du(\delta \varphi)] - \text{trac}_g(du)^*(\nabla' \eta') \xi'.$$

We have the characterisation theorem :

Theorem 3.2. : *Let $u : (M, \varphi, g) \rightarrow (N, \varphi', \xi', \eta', h)$ be a (φ, φ') -holomorphic map from an almost para-Hermitian manifold to an almost paracontact metric manifold. Then u is a harmonic map if only if*

$$\begin{aligned} \text{trac}_g(du)^*(\nabla' \varphi') &= -du(\delta \varphi) \\ \text{and} \\ \text{trac}_g(du)^*(\nabla' \eta') &= 0. \end{aligned}$$

Lemma 3.3. : *Let $u : (M, \varphi, g) \rightarrow (N, \varphi', \xi', \eta', h)$ be a (φ, φ') -holomorphic map from an almost para-Hermitian manifold to an almost paracontact metric manifold.*

1) *If $(N, \varphi', \xi', \eta', h)$ is quasi para-K-cosymplectic, then*

$$\begin{aligned} \text{trac}_g(du)^*(\nabla' \varphi') &= 0, \text{ and} \\ \text{trac}_g(du)^*(\nabla' \eta') &= 0. \end{aligned}$$

2) *If $(N, \varphi', \xi', \eta', h)$ is para-Sasakian, then*

$$\begin{aligned} \text{trac}_g(du)^*(\nabla'\varphi) &= -2 \sum_{k=1}^m h(du(X_k), du(X_k))\xi', \text{ and} \\ \text{trac}_g(du)^*(\nabla'\eta') &= 0. \end{aligned}$$

The above lemma allows us to obtain the expression of $\tau(u)$ when $(N, \varphi', \xi', \eta', h)$ is quasi para-K-cosymplectic or para-Sasakian.

Corollary 3.4. *Let $u : (M, \varphi, g) \rightarrow (N, \varphi', \xi', \eta', h)$ be a (φ, φ') -holomorphic map from an almost para-Hermitian manifold to an almost paracontact metric manifold.*

1) *If $(N, \varphi', \xi', \eta', h)$ is quasi para-K-cosymplectic or para-Sasakian, then*

$$\tau(u) = -\varphi' du(\delta\varphi).$$

From the previous result, we obtain some criteria for harmononicity of (φ, φ') -holomorphic maps between some classes of almost paracontact metric manifolds.

Corollary 3.5. *Let $u : (M, \varphi, g) \rightarrow (N, \varphi', \xi', \eta', h)$ be a (φ, φ') -holomorphic map from an almost semi-para-Kähler manifold to a quasi para-K-cosymplectic or para-Sasakian manifold.*

Then u is an harmonic map.

4. PROOFS OF THE RESULTS

4.1. Proof of proposition 3.1.

Proof. Let $(X_1, \dots, X_m, \varphi X_1, \dots, \varphi X_m)$ be a local φ -basis of M . We have :

$$\begin{aligned} \tau(u) &= \sum_{k=1}^m [(\nabla'^* du)(X_k, X_k) - (\nabla'^* du)(\varphi X_k, \varphi X_k)] \\ &= \sum_{k=1}^m [\nabla'^*_{X_k}(du(X_k)) - \nabla'^*_{\varphi X_k}(du(\varphi X_k))] - du\left[\sum_{k=1}^m (\nabla_{X_k} X_k - \nabla_{\varphi X_k} \varphi X_k)\right]. \\ -\delta\varphi &= \sum_{k=1}^m (\nabla\varphi)(X_k, X_k) - \sum_{k=1}^m (\nabla\varphi)(\varphi X_k, \varphi X_k) \\ &= \sum_{k=1}^m [\nabla_{X_k} \varphi X_k - \varphi(\nabla_{X_k} X_k)] - \sum_{k=1}^m \nabla_{\varphi X_k} \varphi(\varphi X_k) - \varphi(\nabla_{\varphi X_k} \varphi X_k) \\ &= \sum_{k=1}^m [\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} X_k] - \sum_{k=1}^m \varphi(\nabla_{X_k} X_k - \nabla_{\varphi X_k} \varphi X_k), \end{aligned}$$

since : $\varphi^2(X_k) = X_k$

So:

$$\varphi(-\delta\varphi) = \varphi\left(\sum_{k=1}^m [\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} (X_k)]\right) - \varphi^2\left(\sum_{k=1}^m [\nabla_{X_k} X_k - \nabla_{\varphi X_k} \varphi X_k]\right).$$

As $\varphi^2 = I$, we have:

$$\varphi(-\delta\varphi) = \varphi\left(\sum_{k=1}^m [\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} (X_k)]\right) - \sum_{k=1}^m [\nabla_{X_k} X_k - \nabla_{\varphi X_k} \varphi X_k]..$$

We deduce that

$$\sum_{k=1}^m [\nabla_{X_k} X_k - \nabla_{\varphi X_k} \varphi X_k] = \varphi(\delta\varphi) + \varphi\left(\sum_{k=1}^m [\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} (X_k)]\right).$$

By applying du , we get

$$du\left(\sum_{k=1}^m [\nabla_{X_k} X_k + \nabla_{\varphi X_k} \varphi X_k]\right) = duo\varphi(\delta\varphi) + duo\varphi\left(\sum_{k=1}^m [\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} (X_k)]\right).$$

$\tau(u)$ becomes including the fact that $duo\varphi = \varphi'odu$,

$$\begin{aligned} \tau(u) &= \sum_{k=1}^m [\nabla_{X_k}^{*'}(du(X_k)) - \nabla_{\varphi X_k}^{*'}(du(\varphi X_k))] \\ &\quad - \{duo\varphi(\delta\varphi) + duo\varphi\left(\sum_{k=1}^m [\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} (X_k)]\right)\} \\ &= \sum_{k=1}^m [\nabla_{X_k}^{*'}(du(X_k)) - \nabla_{\varphi X_k}^{*'}(du(\varphi X_k))] \\ &\quad - duo\varphi(\delta\varphi) - \varphi'odu\left(\sum_{k=1}^m [\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} (X_k)]\right). \end{aligned}$$

One knows that, for any C^2 -map $f : M \rightarrow N$ and any vectors fields X and Y , we have

$$\begin{aligned} 0 &= d^2 f(X, Y) = \nabla_X^{*'}(df(Y)) - \nabla_Y^{*'}(df(X)) - df[X, Y] \\ df(\nabla_X Y - \nabla_Y X) &= df[X, Y] = \nabla_X^{*'}(df(Y)) - \nabla_Y^{*'}(df(X)). \end{aligned}$$

Then

$$du(\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} X_k) = \nabla_{X_k}^{*'}(du(\varphi X_k)) - \nabla_{\varphi X_k}^{*'}(du(X_k)).$$

Since u is (φ, φ') -holomorphic map, we have:

$$du(\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} X_k) = \nabla_{X_k}^{*'}(\varphi'odu(X_k)) - \nabla_{\varphi X_k}^{*'}(du(X_k)),$$

This leads to

$$\begin{aligned} \tau(u) &= \sum_{k=1}^m [\nabla_{X_k}^{*'}(du(X_k)) - \nabla_{\varphi X_k}^{*'}(du(\varphi X_k))] - duo\varphi(\delta\varphi) \\ &\quad - \varphi'\{\nabla_{X_k}^{*'}(\varphi'odu(X_k)) - \nabla_{\varphi X_k}^{*'}(du(X_k))\}. \end{aligned}$$

Let us set Z , the sum of the two first elements of $\tau(u)$ ie

$$Z = \sum_{k=1}^m [\nabla_{X_k}^{*'}(du(X_k)) - \nabla_{\varphi X_k}^{*'}(du(\varphi X_k))]$$

With $\varphi'^2(Z) := Z - \eta'(Z)\xi \Rightarrow Z = \varphi'^2(Z) + \eta'(Z)\xi'$, we have :

$$\begin{aligned} Z &= \varphi'\{\varphi'\{\sum_{k=1}^m [\nabla_{X_k}^{*'}(du(X_k)) - \nabla_{\varphi X_k}^{*'}(du(\varphi X_k))]\}\} \\ &\quad + \eta'\{\sum_{k=1}^m [\nabla_{X_k}^{*'}(du(X_k)) - \nabla_{\varphi X_k}^{*'}(du(\varphi X_k))]\}\xi', \end{aligned}$$

thus

$$\begin{aligned} Z &= \varphi'\{\sum_{k=1}^m [\varphi'(\nabla_{X_k}^{*'}(du(X_k))) - \varphi'(\nabla_{\varphi X_k}^{*'}(du(\varphi X_k)))]\} \\ &\quad + \eta'\{\sum_{k=1}^m [\nabla_{X_k}^{*'}(du(X_k)) - \nabla_{\varphi X_k}^{*'}(du(\varphi X_k))]\}\xi', \end{aligned}$$

So $\tau(u)$ turns to :

$$\begin{aligned}
\tau(u) &= \varphi' \left\{ \sum_{k=1}^m [\varphi'(\nabla_{X_k}^{*'}(du(X_k))) - \varphi'(\nabla_{\varphi X_k}^{*'}(du(\varphi X_k)))] \right\} \\
&- du \circ \varphi(\delta\varphi) - \varphi' \left\{ \sum_{k=1}^m [\nabla_{X_k}^{*'}(\varphi' \circ du(X_k)) - \nabla_{\varphi X_k}^{*'}(du(X_k))] \right\} \\
&+ \eta' \left\{ \sum_{k=1}^m [\nabla_{X_k}^{*'}(du(X_k)) - \nabla_{\varphi X_k}^{*'}(du(\varphi X_k))] \right\} \xi'.
\end{aligned}$$

By gathering we have :

$$\begin{aligned}
\tau(u) &= \varphi' \left\{ \sum_{k=1}^m [-\nabla_{X_k}^{*'}(\varphi' \circ du(X_k)) + \nabla_{\varphi X_k}^{*'}(du(X_k)) \right. \\
&+ \left. \varphi'(\nabla_{X_k}^{*'}(du(X_k))) - \varphi'(\nabla_{\varphi X_k}^{*'}(du(\varphi X_k)))] \right\} \\
&- du \circ \varphi(\delta\varphi) + \eta' \left\{ \sum_{k=1}^m [\nabla_{X_k}^{*'}(du(X_k)) - \nabla_{\varphi X_k}^{*'}(du(\varphi X_k))] \right\} \xi'.
\end{aligned}$$

Since $\nabla_X^{*'} V = \nabla'_{du(X)} V$, where V is a C^∞ -section of the induced bundle $u^{-1}(TN)$,

$$\begin{aligned}
\tau(u) &= \varphi' \left\{ \sum_{k=1}^m [-\nabla'_{du(X_k)}(\varphi' \circ du(X_k)) + \nabla'_{du(\varphi X_k)}(du(X_k)) \right. \\
&+ \left. \varphi'(\nabla'_{du(X_k)}(du(X_k))) - \varphi'(\nabla'_{du(\varphi X_k)}(du(\varphi X_k)))] \right\} \\
&- du \circ \varphi(\delta\varphi) + \eta' \left\{ \sum_{k=1}^m [\nabla'_{du(X_k)}(du(X_k)) - \nabla'_{du(\varphi X_k)}(du(\varphi X_k))] \right\} \xi'.
\end{aligned}$$

Let us call the elements in the brackets $E_i (i \in \{1, \dots, 4\})$.

$$E_1 + E_3 = -(\nabla'_{du(X_k)} \varphi')(du(X_k)).$$

Since $\varphi^2(X_k) = X_k$, then

$$\nabla'_{du(\varphi X_k)}(du(X_k)) = \nabla'_{du(\varphi X_k)}(du(\varphi^2 X_k)) = \nabla'_{du(\varphi X_k)} \varphi'(du(\varphi X_k)).$$

$$\begin{aligned}
E_2 + E_4 &= \nabla'_{du(\varphi X_k)} \varphi'(du(\varphi X_k)) - \varphi'(\nabla'_{du(\varphi X_k)}(du(\varphi X_k))) \\
&= (\nabla'_{du(\varphi X_k)} \varphi') du(\varphi X_k).
\end{aligned}$$

Thus

$$\begin{aligned}
E_1 + E_2 + E_3 + E_4 &= -(\nabla'_{du(X_k)} \varphi')(du(X_k)) + (\nabla'_{du(\varphi X_k)} \varphi') du(\varphi X_k) \\
&=: -trac_g(du)^*(\nabla' \varphi').
\end{aligned}$$

So

$$\begin{aligned}
\tau(u) &= -\varphi'(trac_g(du)^*(\nabla' \varphi')) - du \circ \varphi(\delta\varphi) \\
&+ \eta' \left\{ \sum_{k=1}^m [\nabla'_{du(X_k)}(du(X_k)) - \nabla'_{du(\varphi X_k)}(du(\varphi X_k))] \right\} \xi'.
\end{aligned}$$

We have:

$$\begin{aligned}
trac_g(du)^*(\nabla' \eta') &:= \sum_{k=1}^m [(\nabla'_{du(X_k)} \eta')(du(X_k)) - (\nabla'_{du(\varphi X_k)} \eta') du(\varphi X_k)] \\
&= \sum_{k=1}^m [\nabla'_{du(X_k)} (\eta' o du(X_k)) - \eta'(\nabla'_{du(X_k)} (du(X_k))) \\
&\quad - \nabla'_{du(\varphi X_k)} (\eta' o du(\varphi X_k)) + \eta'(\nabla'_{du(\varphi X_k)} (du(\varphi X_k)))].
\end{aligned}$$

Since u is (φ, φ') -holomorphic map and $\eta' o \varphi' = 0$, we have:

$$\begin{aligned}
\eta' o du(\varphi X_k) &= \eta' o \varphi' du(X_k) = 0 \\
\eta' o du(X_k) &= \eta' o du(\varphi^2 X_k) = \eta' o \varphi' du(\varphi X_k) = 0,
\end{aligned}$$

then the first and the third elements vanish so

$$\begin{aligned}
trac_g(du)^*(\nabla' \eta') &= \sum_{k=1}^m [-\eta'(\nabla'_{du(X_k)} (du(X_k))) + \eta'(\nabla'_{du(\varphi X_k)} (du(\varphi X_k)))] \\
&= \eta' \left\{ \sum_{k=1}^m [-\nabla'_{du(X_k)} (du(X_k)) + \nabla'_{du(\varphi X_k)} (du(\varphi X_k))] \right\} \\
&= -\eta' \left\{ \sum_{k=1}^m [\nabla'_{du(X_k)} (du(X_k)) - \nabla'_{du(\varphi X_k)} (du(\varphi X_k))] \right\}.
\end{aligned}$$

And $\tau(u)$ becomes

$$\tau(u) = -\varphi'(trac_g(du)^*(\nabla' \varphi')) - du o \varphi(\delta \varphi) - trac_g(du)^*(\nabla' \eta') \xi'.$$

Finally:

$$\tau(u) = -\varphi'[trac_g(du)^*(\nabla' \varphi') + du(\delta \varphi)] - trac_g(du)^*(\nabla' \eta') \xi'.$$

□

4.2. Proof of theorem 3.2.

Proof. : A C^2 -map u is a harmonic map if only if $\tau(u) = 0$, which is equivalent in virtue of Proposition 3.1 to the equalities

$$\begin{aligned}
trac_g(du)^*(\nabla' \varphi') &= -du(\delta \varphi) \\
&\text{and} \\
trac_g(du)^*(\nabla' \eta') &= 0.
\end{aligned}$$

□

4.3. Proof of lemma 3.3.

Proof. Let $\{X_1, \dots, X_m, \varphi X_1, \dots, \varphi X_m\}$ be a local φ -basis of M .

First assume that $(N, \varphi', \xi', \eta', h)$ is quasi para-K-cosymplectic ie $(\nabla'_X \varphi')Y - (\nabla'_{\varphi' X} \varphi')\varphi Y + \eta'(Y)\nabla'_{\varphi' X} \xi' = 0$, for all $X, Y \in \chi(N)$.

Let us recall that

$$trac_g(du)^*(\nabla' \varphi') = \sum_{k=1}^m (\nabla'_{du(X_k)} \varphi')(du(X_k)) - (\nabla'_{du(\varphi X_k)} \varphi') du(\varphi X_k)$$

We shew that $\eta(du(X_k)) = 0$, then

$$\begin{aligned}
(\nabla'_{du(X_k)} \varphi') du(X_k) - (\nabla'_{du(\varphi X_k)} \varphi') du(\varphi X_k) &= (\nabla'_{du(X_k)} \varphi') du(X_k) - (\nabla'_{\varphi' du(X_k)} \varphi') \varphi' du(X_k) \\
&\quad + \eta(du(X_k)) \nabla'_{\varphi'(du(X_k))} \xi' \\
&= 0.
\end{aligned}$$

Let us now look at the case of $\text{trac}_g(du)^*(\nabla'\eta')$.

$$\text{trac}_g(du)^*(\nabla'\eta') = \sum_{k=1}^m [(\nabla'_{du(X_k)}\eta')(du(X_k)) - (\nabla'_{du(\varphi X_k)}\eta')du(\varphi X_k)].$$

Since $(\nabla'_X\eta')Y = h((\nabla'_X\varphi')\varphi'Y, \xi'), \forall X, Y \in \chi(N)$,

$$\begin{aligned} \text{trac}_g(du)^*(\nabla'\eta') &= h\left\{\sum_{k=1}^m [(\nabla'_{du(X_k)}\varphi')\varphi'(du(X_k)) - (\nabla'_{du(\varphi X_k)}\varphi')\varphi'(du(\varphi X_k))], \xi'\right\} \\ &= \sum_{k=1}^m h\{(\nabla'_{du(X_k)}\varphi')\varphi'(du(X_k)) - (\nabla'_{\varphi'(du(X_k))}\varphi')\varphi'(\varphi'(du(X_k))), \xi'\}. \end{aligned}$$

$$\begin{aligned} \text{trac}_g(du)^*(\nabla'\eta') &= h\left\{\sum_{k=1}^m (\nabla'_{du(X_k)}\varphi')\varphi'(du(X_k)) - (\nabla'_{\varphi'(du(X_k))}\varphi')\varphi'(\varphi'(du(X_k))) \right. \\ &\quad \left. + \eta'(\varphi'(du(X_k)))\nabla'_{\varphi'(du(X_k))}\xi', \xi'\right\}, \end{aligned}$$

because $\eta' \circ \varphi' = 0 \Rightarrow \eta'(\varphi'(du(X_k)))\nabla'_{\varphi'(du(X_k))}\xi = 0$

This also shows that if $(N, \varphi', \xi', \eta', h)$ is quasi para-K-cosymplectic, then $\text{trac}_g(du)^*(\nabla'\eta') = 0$

Secondly assume that $(N, \varphi', \xi', \eta', h)$ is para-Sasakian ie $(\nabla'_X\varphi')Y = -h(X, Y)\xi' + \eta'(Y)X, \forall X, Y \in \chi(N)$.

$$\begin{aligned} (\nabla'_{du(X_k)}\varphi')du(X_k) &= -h(du(X_k), du(X_k))\xi' + \eta'(du(X_k))du(X_k) \\ &= -h(du(X_k), du(X_k))\xi' \end{aligned}$$

$$\begin{aligned} -(\nabla'_{du(\varphi X_k)}\varphi')du(\varphi X_k) &= h(du(\varphi X_k), du(\varphi X_k))\xi' - \eta'(du(\varphi X_k))du(\varphi X_k) \\ &= h(\varphi'du(X_k), \varphi'du(X_k))\xi' \\ &= -h(du(X_k), du(X_k))\xi', \end{aligned}$$

then

$$\text{trac}_g(du)^*(\nabla'\varphi) = -2 \sum_{k=1}^m h(du(X_k), du(X_k))\xi'$$

Let us now look at the case of $\text{trac}_g(du)^*(\nabla'\eta)$.

We just shew above that

$$\text{trac}_g(du)^*(\nabla'\eta') = \sum_{k=1}^m h\{(\nabla'_{du(X_k)}\varphi')\varphi'(du(X_k)) - (\nabla'_{\varphi'(du(X_k))}\varphi')\varphi'(\varphi'(du(X_k))), \xi'\}.$$

By the para-Sasakian structure (see above)

$$(\nabla'_{du(X_k)}\varphi')\varphi'(du(X_k)) = -h(du(X_k), \varphi'du(X_k))\xi'$$

$$\begin{aligned} (\nabla'_{\varphi'du(X_k)}\varphi')\varphi'(\varphi'(du(X_k))) &= -h(\varphi'(du(X_k)), \varphi'^2(du(X_k)))\xi' \\ &= -h(\varphi'(du(X_k)), du(X_k))\xi', \end{aligned}$$

because $\varphi'^2(du(X_k)) = du(X_k) - \eta'(du(X_k))\xi' = du(X_k)$ by the fact that $\eta'(du(X_k)) = 0$ (see above).

One can conclude that $\text{trac}_g(du)^*(\nabla'\eta') = 0$. $\square$

5. RESULTS FOR SECTIONS

Proposition 5.1. *Let $[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, g)]$ be a (semi – Riemannian) almost paracontact fiber bundle of type II and $u : M \rightarrow N$ be a holomorphic section of $[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, g)]$. Then we have:*

$$\begin{aligned} \tau^v(u) &= -\varphi' \{ \text{trac}_g(d^v u)^*(\nabla^v \varphi') + \text{trac}_g(\nabla_{d^h u}^v \varphi')(d^v u) + \text{trac}_g(A_{d^h u} d^h u o \varphi) \\ &\quad + \text{trac}_g(T_{d^v u} d^h u o \varphi) + d^v u(\delta \varphi) \} \\ &\quad - \{ \text{trac}_g(d^v u)^*(\nabla' \eta') + \text{trac}_g(\nabla_{d^h u}' \eta')(d^v u) \} \xi'. \end{aligned}$$

Corollary 5.2. *Let $u : M \rightarrow N$ be a holomorphic section of a (semi – Riemannian) almost paracontact fiber bundle of type II, $[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, g)]$. If the fibers are quasi-para-K-oosymplectic submanifolds, M is almost semi-para-Kähler and if*

$$\begin{aligned} \text{trac}_g(\nabla_{d^h u}^v \varphi')(d^v u) &= 0 \\ \text{trac}_g(A_{d^h u} d^h u o \varphi) &= 0 \\ \text{trac}_g(T_{d^v u} d^h u o \varphi) &= 0, \end{aligned}$$

then u is a harmonic section.

Corollary 5.3. :

Let $u : M \rightarrow N$ be a holomorphic section of a (semi – Riemannian) almost paracontact fiber bundle of type II, $[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, g)]$. If the fibers are totally geodesic submanifold and quasi-para-K-oosymplectic submanifolds, M is almost semi-para-Kähler manifold, the horizontal distribution is integrable and $(\nabla_Y^v \varphi')U = 0$ for all Y horizontal and U vertical, then u is a harmonic section.

6. PROOFS OF THE RESULTS:

6.1. Proof of proposition 4.1.

Proof. Let $\{X_1, \dots, X_m, \varphi X_1, \dots, \varphi X_m\}$ be a local φ -basis of M . Then we have :

$$\begin{aligned} \tau^v(u) &= \sum_{k=1}^m (\nabla^{*v} d^v u)(X_k, X_k) - \sum_{k=1}^m (\nabla^{*v} d^v u)(\varphi X_k, \varphi X_k) \\ &= \sum_{k=1}^m [\nabla_{X_k}^{*v} (d^v u(X_k)) - \nabla_{\varphi X_k}^{*v} (d^v u(\varphi X_k))] - d^v u \left[\sum_{k=1}^m (\nabla_{X_k} X_k - \nabla_{\varphi X_k} \varphi X_k) \right]. \end{aligned}$$

In the maps case we shew that

$$\sum_{k=1}^m [\nabla_{X_k} X_k - \nabla_{\varphi X_k} \varphi X_k] = \varphi(\delta \varphi) + \varphi \left(\sum_{k=1}^m [\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} (X_k)] \right).$$

By applying $d^v u$, we get

$$d^v u \left(\sum_{k=1}^m [\nabla_{X_k} X_k + \nabla_{\varphi X_k} \varphi X_k] \right) = d^v u o \varphi(\delta \varphi) + d^v u o \varphi \left(\sum_{k=1}^m [\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} (X_k)] \right).$$

$\tau^v(u)$ becomes including the fact that $d^v u o \varphi = \varphi' o d^v u$,

$$\begin{aligned}
\tau^v(u) &= \sum_{k=1}^m [\nabla_{X_k}^{*v}(d^v u(X_k)) - \nabla_{\varphi X_k}^{*v}(d^v u(\varphi X_k))] \\
&- \{d^v u \circ \varphi(\delta\varphi) + d^v u \circ \varphi(\sum_{k=1}^m [\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k}(X_k)])\} \\
&= \sum_{k=1}^m [\nabla_{X_k}^{*v}(d^v u(X_k)) - \nabla_{\varphi X_k}^{*v}(d^v u(\varphi X_k))] \\
&- d^v u \circ \varphi(\delta\varphi) - \varphi' \circ d^v u(\sum_{k=1}^m [\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k}(X_k)]).
\end{aligned}$$

We shew in the maps case that

$$du(\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} X_k) = \nabla_{X_k}'(du(\varphi X_k)) - \nabla_{\varphi X_k}'(du(X_k)).$$

Taking the vertical part we get:

$\forall k = 1, \dots, m,$

$$\begin{aligned}
d^v u(\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} X_k) &= \nabla_{X_k}^{*v}(du(\varphi X_k)) - \nabla_{\varphi X_k}^{*v}(du(X_k)) \\
&= \nabla_{X_k}^{*v}(d^v u(\varphi X_k) + d^h u(\varphi X_k)) - \nabla_{\varphi X_k}^{*v}(d^v u(X_k) + d^h u(X_k)).
\end{aligned}$$

So

$$\begin{aligned}
d^v u(\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} X_k) &= \nabla_{X_k}^{*v}(d^v u(\varphi X_k)) - \nabla_{\varphi X_k}^{*v}(d^v u(X_k)) + \nabla_{X_k}^{*v}(d^h u(\varphi X_k)) \\
&- \nabla_{\varphi X_k}^{*v}(d^h u(X_k)).
\end{aligned}$$

Since u is (φ, φ') -holomorphic section, we have:

$$\begin{aligned}
d^v u(\nabla_{X_k} \varphi X_k - \nabla_{\varphi X_k} X_k) &= \nabla_{X_k}^{*v}(\varphi' \circ d^v u(X_k)) - \nabla_{\varphi X_k}^{*v}(d^v u(X_k)) \\
&+ \nabla_{X_k}^{*v}(d^h u(\varphi X_k)) - \nabla_{\varphi X_k}^{*v}(d^h u(X_k)),
\end{aligned}$$

This leads to

$$\begin{aligned}
\tau^v(u) &= \sum_{k=1}^m [\nabla_{X_k}^{*v}(d^v u(X_k)) - \nabla_{\varphi X_k}^{*v}(d^v u(\varphi X_k))] - d^v u \circ \varphi(\delta\varphi) \\
&- \varphi' \{ \nabla_{X_k}^{*v}(\varphi' \circ d^v u(X_k)) - \nabla_{\varphi X_k}^{*v}(d^v u(X_k)) \\
&+ \nabla_{X_k}^{*v}(d^h u(\varphi X_k)) - \nabla_{\varphi X_k}^{*v}(d^h u(X_k)) \}.
\end{aligned}$$

Let us set Z , the sum of the two first elements of $\tau^v(u)$ ie

$$Z = \sum_{k=1}^m [\nabla_{X_k}^{*v}(d^v u(X_k)) - \nabla_{\varphi X_k}^{*v}(d^v u(\varphi X_k))].$$

With $\varphi'^2(Z) := Z - \eta(Z)\xi'$, we have :

$$\begin{aligned}
Z &= \varphi' \{ \varphi' \{ \sum_{k=1}^m [\nabla_{X_k}^{*v}(d^v u(X_k)) - \nabla_{\varphi X_k}^{*v}(d^v u(\varphi X_k))] \} \} \\
&+ \eta' \{ \sum_{k=1}^m [\nabla_{X_k}^{*v}(d^v u(X_k)) - \nabla_{\varphi X_k}^{*v}(d^v u(\varphi X_k))] \} \xi'.
\end{aligned}$$

So $\tau^v(u)$ turns to :

$$\begin{aligned}
\tau^v(u) &= \varphi' \left\{ \sum_{k=1}^m [\varphi'(\nabla_{X_k}^{*v}(d^v u(X_k))) - \varphi'(\nabla_{\varphi X_k}^{*v}(d^v u(\varphi X_k)))] \right\} \\
&- d^v u o \varphi(\delta \varphi) - \varphi' \left\{ \sum_{k=1}^m [\nabla_{X_k}^{*v}(\varphi' o d^v u(X_k)) - \nabla_{\varphi X_k}^{*v}(d^v u(X_k)) \right. \\
&+ \left. \nabla_{X_k}^{*v}(d^h u(\varphi X_k)) - \nabla_{\varphi X_k}^{*v}(d^h u(X_k))] \right\} \\
&+ \eta' \left\{ \sum_{k=1}^m [\nabla_{X_k}^{*v}(d^v u(X_k)) - \nabla_{\varphi X_k}^{*v}(d^v u(\varphi X_k))] \right\} \xi'.
\end{aligned}$$

By gathering we have :

$$\begin{aligned}
\tau^v(u) &= \varphi' \left\{ \sum_{k=1}^m [-\nabla_{X_k}^{*v}(\varphi' o d^v u(X_k)) + \nabla_{\varphi X_k}^{*v}(d^v u(X_k)) - \nabla_{X_k}^{*v}(d^h u(\varphi X_k)) \right. \\
&+ \left. \nabla_{\varphi X_k}^{*v}(d^h u(X_k)) + \varphi'(\nabla_{X_k}^{*v}(d^v u(X_k))) - \varphi'(\nabla_{\varphi X_k}^{*v}(d^v u(\varphi X_k)))] \right\} \\
&- d^v u o \varphi(\delta \varphi) + \eta' \left\{ \sum_{k=1}^m [\nabla_{X_k}^{*v}(d^v u(X_k)) - \nabla_{\varphi X_k}^{*v}(d^v u(\varphi X_k))] \right\} \xi'.
\end{aligned}$$

From relation (2.1), we get

$$\begin{aligned}
\tau^v(u) &= \varphi' \left\{ \sum_{k=1}^m [-\nabla_{du(X_k)}^v(\varphi' o d^v \tilde{u}(X_k)) + \nabla_{du(\varphi X_k)}^v(d^v \tilde{u}(X_k)) \right. \\
&- \left. \nabla_{du(X_k)}^v(d^h \tilde{u}(\varphi X_k)) + \nabla_{du(\varphi X_k)}^v(d^h \tilde{u}(X_k)) \right. \\
&+ \left. \varphi'(\nabla_{du(X_k)}^v(d^v \tilde{u}(X_k))) - \varphi'(\nabla_{du(\varphi X_k)}^v(d^v \tilde{u}(\varphi X_k))) \right\} \\
&- d^v u o \varphi(\delta \varphi) + \eta' \left\{ \sum_{k=1}^m [\nabla_{du(X_k)}^v(d^v \tilde{u}(X_k)) - \nabla_{du(\varphi X_k)}^v(d^v \tilde{u}(\varphi X_k))] \right\} \xi'.
\end{aligned}$$

As $du(Y) = d^v u(Y) + d^h u(Y)$,

$$\begin{aligned}
\tau^v(u) &= \varphi' \left\{ \sum_{k=1}^m [-\nabla_{d^v u(X_k)}^v(\varphi' o d^v \tilde{u}(X_k)) - \nabla_{d^h u(X_k)}^v(\varphi' o d^v \tilde{u}(X_k)) \right. \\
&+ \left. \nabla_{d^v u(\varphi X_k)}^v(d^v \tilde{u}(X_k)) + \nabla_{d^h u(\varphi X_k)}^v(d^v \tilde{u}(X_k)) \right. \\
&- \left. \nabla_{d^v u(X_k)}^v(d^h \tilde{u}(\varphi X_k)) - \nabla_{d^h u(X_k)}^v(d^h \tilde{u}(\varphi X_k)) \right. \\
&+ \left. \nabla_{d^v u(\varphi X_k)}^v(d^h \tilde{u}(X_k)) + \nabla_{d^h u(\varphi X_k)}^v(d^h \tilde{u}(X_k)) \right. \\
&+ \left. \varphi'(\nabla_{d^v u(X_k)}^v(d^v \tilde{u}(X_k))) + \varphi'(\nabla_{d^h u(X_k)}^v(d^v \tilde{u}(X_k))) \right. \\
&- \left. \varphi'(\nabla_{d^v u(\varphi X_k)}^v(d^v \tilde{u}(\varphi X_k))) - \varphi'(\nabla_{d^h u(\varphi X_k)}^v(d^v \tilde{u}(\varphi X_k))) \right\} \\
&- d^v u o \varphi(\delta \varphi) + \eta' \left\{ \sum_{k=1}^m [\nabla_{d^v u(X_k)}^v(d^v \tilde{u}(X_k)) - \nabla_{d^v u(\varphi X_k)}^v(d^v \tilde{u}(\varphi X_k))] \right\} \xi' \\
&+ \eta' \left\{ \sum_{k=1}^m [\nabla_{d^h u(X_k)}^v(d^v \tilde{u}(X_k)) - \nabla_{d^h u(\varphi X_k)}^v(d^v \tilde{u}(\varphi X_k))] \right\} \xi'.
\end{aligned}$$

Let us call the elements in the brackets $E_i (i \in \{1, \dots, 12\})$.

$E_1 + E_9 = -(\nabla_{d^v u(X_k)}^v \varphi')(d^v \tilde{u}(X_k))$,
since φ is invariant on the vertical distribution also on the horizontal one.

$$\begin{aligned} E_2 + E_{10} &= -(\nabla_{d^h u(X_k)}^v \varphi')(d^v \tilde{u}(X_k)). \\ \text{As } \varphi^2(X_k) &= X_k \text{ then,} \\ \nabla_{d^v u(\varphi X_k)}^v(d^v \tilde{u}(X_k)) &= \nabla_{d^v u(\varphi X_k)}^v(d^v \tilde{u}(\varphi^2 X_k)) = \nabla_{d^v u(\varphi X_k)}^v(\varphi'(d^v \tilde{u}(\varphi X_k))). \end{aligned}$$

$$\begin{aligned} E_3 + E_{11} &= \nabla_{d^v u(\varphi X_k)}^v(\varphi'(d^v \tilde{u}(\varphi X_k))) - \varphi'(\nabla_{d^v u(\varphi X_k)}^v(d^v \tilde{u}(\varphi X_k))) \\ &= (\nabla_{d^v u(\varphi X_k)}^v \varphi')(d^v \tilde{u}(\varphi X_k)). \end{aligned}$$

In the same way:

$$\nabla_{d^h u(\varphi X_k)}^v(d^v \tilde{u}(X_k)) = \nabla_{d^h u(\varphi X_k)}^v(\varphi'(d^v \tilde{u}(\varphi X_k))).$$

$$\begin{aligned} E_4 + E_{12} &= \nabla_{d^h u(\varphi X_k)}^v(\varphi'(d^v \tilde{u}(\varphi X_k))) - \varphi'(\nabla_{d^h u(\varphi X_k)}^v(d^v \tilde{u}(\varphi X_k))) \\ &= (\nabla_{d^h u(\varphi X_k)}^v \varphi')(d^v \tilde{u}(\varphi X_k)). \end{aligned}$$

$$\begin{aligned} E_5 + E_6 + E_7 + E_8 &= -T_{d^v u(X_k)}(d^h \tilde{u}(\varphi X_k)) - A_{d^h u(X_k)}(d^h \tilde{u}(\varphi X_k)) \\ &\quad + T_{d^v u(\varphi X_k)}(d^h \tilde{u}(X_k)) + A_{d^h u(\varphi X_k)}(d^h \tilde{u}(X_k)). \end{aligned}$$

Since $\varphi^2(X_k) = X_k \Rightarrow d^h u(X_k) = d^h u \circ \varphi(X_k)$,

$$\begin{aligned} E_5 + E_6 + E_7 + E_8 &= -T_{d^v u(X_k)}(d^h u \circ \varphi(X_k)) + T_{d^v u(\varphi X_k)}(d^h u \circ \varphi(X_k)) \\ &\quad - A_{d^h u(X_k)}(d^h u \circ \varphi(X_k)) + A_{d^h u(\varphi X_k)}(d^h u \circ \varphi(X_k)). \end{aligned}$$

Putting all this together,

$$\begin{aligned} \tau^v(u) &= \varphi' \left\{ \sum_{k=1}^m [-(\nabla_{d^v u(X_k)}^v \varphi')(d^v \tilde{u}(X_k)) + (\nabla_{d^v u(\varphi X_k)}^v \varphi')(d^v \tilde{u}(\varphi X_k)) \right. \\ &\quad - (\nabla_{d^h u(X_k)}^v \varphi')(d^v \tilde{u}(X_k)) + (\nabla_{d^h u(\varphi X_k)}^v \varphi')(d^v \tilde{u}(\varphi X_k)) \\ &\quad - T_{d^v u(X_k)}(d^h u \circ \varphi(X_k)) + T_{d^v u(\varphi X_k)}(d^h u \circ \varphi(X_k)) \\ &\quad - A_{d^h u(X_k)}(d^h u \circ \varphi(X_k)) + A_{d^h u(\varphi X_k)}(d^h u \circ \varphi(X_k))] \} \\ &\quad - d^v u \circ \varphi(\delta \varphi) + \eta' \left\{ \sum_{k=1}^m [\nabla_{d^v u(X_k)}^v(d^v \tilde{u}(X_k)) - \nabla_{d^v u(\varphi X_k)}^v(d^v \tilde{u}(\varphi X_k))] \right\} \xi' \\ &\quad + \eta' \left\{ \sum_{k=1}^m [\nabla_{d^h u(X_k)}^v(d^v \tilde{u}(X_k)) - \nabla_{d^h u(\varphi X_k)}^v(d^v \tilde{u}(\varphi X_k))] \right\} \xi'. \end{aligned}$$

One gets :

$$\begin{aligned}
\tau^v(u) &= \varphi' \left\{ \sum_{k=1}^m [-(\nabla_{d^v u(X_k)}^v \varphi')(d^v u(\tilde{X}_k)) + (\nabla_{d^v u(\varphi X_k)}^v \varphi')(d^v u(\tilde{\varphi X}_k))] \right. \\
&+ \sum_{k=1}^m [-(\nabla_{d^h u(X_k)}^v \varphi')(d^v u(\tilde{X}_k)) + (\nabla_{d^h u(\varphi X_k)}^v \varphi')(d^v u(\tilde{\varphi X}_k))] \\
&+ \sum_{k=1}^m [-T_{d^v u(X_k)}(d^h u o \varphi(X_k)) + T_{d^v u(\varphi X_k)}(d^h u o \varphi(X_k))] \\
&+ \sum_{k=1}^m [-A_{d^h u(X_k)}(d^h u o \varphi(X_k)) + A_{d^h u(\varphi X_k)}(d^h u o \varphi(X_k))] \Big\} \\
&- d^v u o \varphi(\delta \varphi) + \eta' \left\{ \sum_{k=1}^m [\nabla_{d^v u(X_k)}^v (d^v u(\tilde{X}_k)) - \nabla_{d^v u(\varphi X_k)}^v (d^v u(\tilde{\varphi X}_k))] \right\} \xi' \\
&+ \eta' \left\{ \sum_{k=1}^m [\nabla_{d^h u(X_k)}^v (d^v u(\tilde{X}_k)) - \nabla_{d^h u(\varphi X_k)}^v (d^v u(\tilde{\varphi X}_k))] \right\} \xi'.
\end{aligned}$$

Then

$$\begin{aligned}
\tau^v(u) &= \varphi' [-trac_g(d^v u)^*(\nabla^v \varphi') - trac_g(\nabla_{d^h u}^v \varphi')(d^v u) - trac_g(A_{d^h u} d^h u o \varphi) \\
&- trac_g(T_{d^v u} d^h u o \varphi) - d^v u(\delta \varphi)] \\
&+ \eta' \left\{ \sum_{k=1}^m [\nabla_{d^v u(X_k)}^v (d^v u(\tilde{X}_k)) - \nabla_{d^v u(\varphi X_k)}^v (d^v u(\tilde{\varphi X}_k))] \right\} \xi' \\
&+ \eta' \left\{ \sum_{k=1}^m [\nabla_{d^h u(X_k)}^v (d^v u(\tilde{X}_k)) - \nabla_{d^h u(\varphi X_k)}^v (d^v u(\tilde{\varphi X}_k))] \right\} \xi'.
\end{aligned}$$

On the one hand :

$$\begin{aligned}
&\eta' \left\{ \sum_{k=1}^m [\nabla_{d^v u(X_k)}^v (d^v u(\tilde{X}_k)) - \nabla_{d^v u(\varphi X_k)}^v (d^v u(\tilde{\varphi X}_k))] \right\} \\
&= \eta' \left\{ \sum_{k=1}^m [\nabla'_{d^v u(X_k)} (d^v u(\tilde{X}_k)) - \nabla'_{d^v u(\varphi X_k)} (d^v u(\tilde{\varphi X}_k))] \right\},
\end{aligned}$$

since $\eta'(\text{"horizontal"}) = 0$.

We have :

$$\begin{aligned}
trac_g(d^v u)^*(\nabla' \eta') &= \sum_{k=1}^m [(\nabla'_{d^v u(X_k)} \eta')(d^v u(\tilde{X}_k)) - (\nabla'_{d^v u(\varphi X_k)} \eta')(d^v u(\tilde{\varphi X}_k))] \\
&= \sum_{k=1}^m [D_{d^v u(X_k)}(\eta'(d^v u(\tilde{X}_k))) - \eta'(\nabla'_{d^v u(X_k)}(d^v u(\tilde{X}_k))) \\
&- D_{d^v u(\varphi X_k)}(\eta'(d^v u(\tilde{\varphi X}_k))) + \eta'(\nabla'_{d^v u(\varphi X_k)}(d^v u(\tilde{\varphi X}_k)))].
\end{aligned}$$

As $X_k = \varphi^2(X_k) \Rightarrow \eta'(d^v u(X_k)) = \eta'(d^v u(\varphi^2(X_k))) = \eta' o \varphi'(d^v u(\varphi X_k)) = 0$.

And

$$\eta'(d^v u(\varphi X_k)) = \eta o \varphi'(d^v u(X_k)) = 0,$$

then

$$trac_g(d^v u)^*(\nabla' \eta') = -\eta' \left\{ \sum_{k=1}^m [\nabla'_{d^v u(X_k)}(d^v u(\tilde{X}_k)) + \nabla'_{d^v u(\varphi X_k)}(d^v u(\tilde{\varphi X}_k))] \right\}.$$

On the other hand

$$\text{trac}_g(\nabla'_{d^h u} \eta')(d^v u) = \sum_{k=1}^m [(\nabla'_{d^h u(X_k)} \eta')(d^v u(\tilde{X}_k)) - (\nabla'_{d^h u(\varphi X_k)} \eta')(d^v u(\tilde{\varphi} X_k))],$$

in the same way as above

$$\text{trac}_g(\nabla'_{d^h u} \eta')(d^v u) = -\eta' \left\{ \sum_{k=1}^m [\nabla'_{d^h u(X_k)}(d^v u(\tilde{X}_k)) - \nabla'_{d^h u(\varphi X_k)}(d^v u(\tilde{\varphi} X_k))] \right\}.$$

We can conclude that

$$\begin{aligned} \tau^v(u) &= -\varphi' \{ \text{trac}_g(d^v u)^*(\nabla^v \varphi') + \text{trac}_g(\nabla^v_{d^h u} \varphi')(d^v u) + \text{trac}_g(A_{d^h u} d^h u \circ \varphi) \\ &\quad + \text{trac}_g(T_{d^v u} d^h u \circ \varphi) + d^v u(\delta \varphi) \} \\ &\quad - \{ \text{trac}_g(d^v u)^*(\nabla' \eta') + \text{trac}_g(\nabla'_{d^h u} \eta')(d^v u) \} \xi'. \end{aligned}$$

□

Remark. 1) The fibers of a (semi – Riemannian) almost paracontact fiber bundle of type II are almost para-contact metric manifolds. (See Prop.3.1 in [4])

2) Let us justify the fact that

if the fibers are quasi-K-cosymplectic submanifolds, then

$$\text{trac}_g(d^v u)^*(\nabla^v \varphi') = 0 \text{ and } \text{trac}_g(d^v u)^*(\nabla' \eta') = 0$$

Assume the fibers are quasi-K-cosymplectic.

$$\text{Then } (\nabla^v_U \varphi')V - (\nabla^v_{\varphi' U} \varphi')\varphi'V + \eta'(V)\nabla^v_{\varphi' U} \xi' = 0, \forall U, V \in \Gamma(W).$$

We have :

$$\begin{aligned} \text{trac}_g(d^v u)^*(\nabla^v \varphi') &= \sum_{k=1}^m (\nabla^v_{d^v u(X_k)} \varphi')(d^v u(\tilde{X}_k)) - \sum_{k=1}^m (\nabla^v_{d^v u(\varphi X_k)} \varphi')(d^v u(\tilde{\varphi} X_k)) \\ &= \sum_{k=1}^m (\nabla^v_{d^v u(X_k)} \varphi')(d^v u(\tilde{X}_k)) - \sum_{k=1}^m (\nabla^v_{\varphi'(d^v u(X_k))} \varphi')(\varphi'(d^v u(\tilde{X}_k))) \end{aligned}$$

since u is a (φ, φ') -holomorphic section.

$$\text{But } \eta'(d^v u(\tilde{X}_k)) = \eta'(d^v u(\tilde{\varphi}^2 X_k)) = \eta'(\varphi'(d^v u(\tilde{\varphi} X_k))) = \eta' \circ \varphi'(d^v u(\tilde{X}_k)) = 0,$$

$$\text{then } \eta'(d^v u(\tilde{X}_k))\nabla^v_{\varphi'(d^v u(X_k))} \xi' = 0.$$

So

$$\begin{aligned} \text{trac}_g(d^v u)^*(\nabla^v \varphi') &= \sum_{k=1}^m \{ (\nabla^v_{d^v u(X_k)} \varphi')(d^v u(\tilde{X}_k)) - (\nabla^v_{\varphi'(d^v u(X_k))} \varphi')(\varphi'(d^v u(\tilde{X}_k))) \} \\ &\quad + \eta'(d^v u(\tilde{X}_k))\nabla^v_{\varphi'(d^v u(X_k))} \xi'. \end{aligned}$$

This shows that if the fibers are para-quasi-K-cosymplectic, $\text{trac}_g(d^v u)^*(\nabla^v \varphi') = 0$.

Let us now look at the case of $\text{trac}_g(d^v u)^*(\nabla' \eta')$.

$$\text{trac}_g(d^v u)^*(\nabla' \eta') = \sum_{k=1}^m [(\nabla'_{d^v u(X_k)} \eta')(d^v u(\tilde{X}_k)) - (\nabla'_{d^v u(\varphi X_k)} \eta')(d^v u(\tilde{\varphi} X_k))].$$

Since $(\nabla'_X \eta')Y = h((\nabla'_X \varphi')\varphi'Y, \xi')$, for all $X, Y \in \chi(N)$,

$$\begin{aligned}
\text{trac}_g(d^v u)^*(\nabla' \eta') &= \sum_{k=1}^m h\{(\nabla'_{d^v u(X_k)} \varphi') \varphi'(d^v \tilde{u}(X_k)) - (\nabla'_{d^v u(\varphi X_k)} \varphi')(\varphi(d^v \tilde{u}(\varphi X_k))), \xi'\} \\
&= \sum_{k=1}^m h\{(\nabla^v_{d^v u(X_k)} \varphi') \varphi'(d^v \tilde{u}(X_k)) - (\nabla^v_{\varphi'(d^v u(X_k))} \varphi')(\varphi'(\varphi'(d^v \tilde{u}(X_k)))) \\
&\quad + \eta'(\varphi'(d^v \tilde{u}(X_k))) \nabla^v_{\varphi'(d^v u(X_k))} \xi', \xi'\},
\end{aligned}$$

because ξ' is vertical and $\eta'(\varphi'(d^v \tilde{u}(X_k))) \nabla^v_{\varphi'(d^v u(X_k))} \xi' = 0$ since $\eta' \circ \varphi' = 0$.

This also shows that if the fibers are quasi-K-cosymplectic, $\text{trac}_g(d^v u)^*(\nabla' \eta') = 0$.

Remark. If $(\nabla_Y^v \varphi')U = 0$ for all Y horizontal and U vertical, then $(\nabla_Y' \eta')(U) = 0$

In fact

$$\begin{aligned}
(\nabla_X' \eta')U &= h((\nabla_X' \varphi') \varphi' U, \xi') \\
&= h((\nabla_X^v \varphi') \varphi' U, \xi'),
\end{aligned}$$

since ξ is vertical.

This equality testifies the assuming.

6.2. Proof of corollary 5.2.

Proof. Let $[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, g)]$ be a (Riemannian) almost contact fiber bundle of type II.

Assume $\text{trac}_g(\nabla_{d^h u}^v \varphi')(d^v u) = 0$, $\text{trac}_g(A_{d^h u} d^h u \circ \varphi) = 0$, $\text{trac}_g(T_{d^v u} d^h u \circ \varphi) = 0$ and $\text{trac}_g(\nabla_{d^h u}^v \eta')(d^v u) = 0$, then the formula in Proposition 5.1 gives:

$$\tau^v(u) = -\varphi'\{\text{trac}_g(d^v u)^*(\nabla^v \varphi') + d^v u(\delta \varphi)\} - [\text{trac}_g(d^v u)^*(\nabla' \eta')]\xi'.$$

Moreover if M is almost semi-para-Kähler ie $\delta \varphi = 0$ and if the fibers are quasi-para-K-cosymplectic which implies $\text{trac}_g(d^v u)^*(\nabla^v \varphi') = 0$ and $\text{trac}_g(d^v u)^*(\nabla' \eta') = 0$, then $\tau^v(u) = 0$, i.e. u is a harmonic section. $\square$

6.3. Proof of corollary 5.3.

Proof. Let $[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, g)]$ be a (Riemannian) almost contact fiber bundle of type II.

Assume the fibers are quasi-para-K-cosymplectic submanifolds and M is almost semi-para-Kähler, then the formula in Proposition 5.1 gives:

$$\begin{aligned}
\tau^v(u) &= -\varphi'\{\text{trac}_g(\nabla_{d^h u}^v \varphi')(d^v u) + \text{trac}_g(A_{d^h u} d^h u \circ \varphi) + \text{trac}_g(T_{d^v u} d^h u \circ \varphi)\} \\
&\quad - [\text{trac}_g(\nabla_{d^h u}^v \eta')(d^v u)]\xi'.
\end{aligned}$$

Moreover if the fibers are totally geodesic ($ieT = 0$), the horizontal distribution is integrable ($ieA = 0$) and $(\nabla_Y^v \varphi')U = 0$ for all Y horizontal and U vertical, which implies $(\nabla_Y' \eta')(U) = 0$, then one gets $\tau^v(u) = 0$, i.e. u is a harmonic section. $\square$

Before giving examples we should talk a bit about fibers.

Proposition 6.1. Let $[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, g)]$ be a (Riemannian) almost contact fiber bundle of type II.

- 1) If the total space is para-cosymplectic, then the fibers are para-cosymplectic.
- 2) If the total space is almost para-cosymplectic, then the fibers are almost para-cosymplectic.

3) If the total space is quasi-para-K-cosymplectic, then the fibers are quasi-para-K-cosymplectic.

4) If the total space is para-Sasakian, then the fibers are para-Sasakian.

Proof. Let $(\varphi', \xi', \eta', h)$ be the almost paracontact metric structure of the total space and $(\varphi^\Delta, \xi^\Delta, \eta^\Delta, h^\Delta)$, its restriction on the fibers.

1) Assume that the total space is para-cosymplectic ie $\nabla' \varphi' = 0$ and $\nabla' \eta' = 0$

$$\begin{aligned} \forall U, V \in \Gamma(W), (\nabla'_U \varphi') V &= 0 \\ \nabla'_U (\varphi' V) - \varphi' (\nabla'_U V) &= 0 \\ \nabla'_U (\varphi' V) - \varphi' (\nabla_U^v V) - \varphi' (\nabla_U^h V) &= 0 \end{aligned}$$

Since $\varphi'(\text{horizontal})$ is *horizontal* and $\varphi'(\text{vertical})$ is *vertical*, taking the vertical part of the last equality one has

$$\begin{aligned} \nabla_U^v (\varphi' V) - \varphi' (\nabla_U^v V) &= 0 \\ (\nabla_U^\Delta \varphi^\Delta) V &= 0 \end{aligned}$$

$$\begin{aligned} \forall U, V \in \Gamma(W), (\nabla'_U \eta') V &= 0 \\ U. \eta'(V) - \eta' (\nabla'_U V) &= 0 \\ U. \eta'(V) - \eta' (\nabla_U^v V) - \eta' (\nabla_U^h V) &= 0 \\ U. \eta'(V) - \eta' (\nabla_U^v V) &= 0 \\ (\nabla_U^\Delta \eta^\Delta) V &= 0, \end{aligned}$$

since $\eta'(\text{horizontal}) = 0$.

Then we conclude that the fibers are para-cosymplectic.

In the same way, one shows 2), 3) and 4). □

6.4. Example. An example of (*semi-Riemannian*) almost paracontact fiber bundle of type II,

$$[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, g)]$$

for which the Corollary 4.3 applies is a fiber bundle which projection is an almost paracontact (semi-Riemannian) submersion of type II, whose total space is an almost para-cosymplectic or quasi-para-Sasakian manifold with totally geodesic fibers and satisfies :

$$(\nabla_Y^v \varphi') U = 0 \text{ for all } Y \text{ horizontal and } U \text{ vertical.}$$

In fact:

For almost paracontact submersion whose total space is almost para-cosymplectic or quasi-para-Sasakian the horizontal distribution is integrable (ie $A = 0$), see theorem 3.1 and corollary 3.2 in [5].

The base almost *semi-Kähler* (ie $\delta\varphi = 0$). See Prop.3.4 and 3.5 in [4] or Coro. 3.2 and Prop.3.7 in [5].

The fibers are quasi-para-K-cosymplectic submanifolds. (see Prop. 6.1 above).

6.5. Example. Or simply take a fiber bundle which projection is an almost paracontact (semi-Riemannian) submersion of type II whose total space is para-cosymplectic with totally geodesic fibers.

In fact:

If total space is para-cosymplectic, it is almost para-cosymplectic (it satisfies example 1) and moreover

$$\nabla' \varphi' = 0 \Rightarrow (\nabla_Y^v \varphi')U = 0 \text{ for all } Y \text{ horizontal and } U \text{ vertical.}$$

6.6. Example. Let $[(N, \varphi', \xi', \eta', h) \xrightarrow{\pi} (M, \varphi, g)]$ be a (*Riemannian*) almost paracontact fiber bundle of type II such that N is quasi-para-K-cosymplectic or para-Sasakian and $u : M \rightarrow N$ be a section.

If u is a (φ, φ') -holomorphic map, then u is a harmonic section.

In fact :

On the one hand, a section that is a (φ, φ') -holomorphic map is obvious a (φ, φ') -holomorphic section.

In fact

$$\begin{aligned} du\varphi &= \varphi'odu \\ d^v u\varphi &= \varphi'^v du \\ d^v u\varphi &= \varphi'od^v u \end{aligned}$$

because for all $E \in \chi(N)$, $\varphi'(E^v) = \varphi'^v(E)$ since π is an almost paracontact metric submersion.

On the other hand, since u is a (φ, φ') -holomorphic map, then with Corollary 3.5 above, u is a harmonic map; moreover one knows that a section that is harmonic map is a harmonic section.

Acknowledment:

I'm very grateful to my father St Paul for his tremendous help.

REFERENCES

- [1] S. Ianus, A.M.Pastore and M. Falcitelly 2004., Riemannian submersions and related topics, World Scientific Publ. C.
- [2] Ylmaz Gunduzalp, Almost para-Hermitian submersion, Matematik Bechnik 68, 4 (2016), 241253.
- [3] C. M. Wood, The gauss section of a Riemannian immersion, J. London Math. Soc (2) 33 (1986) 157-168.
- [4] Tshikuna-Matamba Tshikunguila, On the base space of an almost paracontact submersion, Facta Universitatis, Vol.31 n°5 (2016) p. 1041-1049.
- [5] Ylmaz Gunduzalp and Bayram Sahin, Para-contact Para-complex semi-Riemannian submersion, Bull. Malays. Math. Sci. Soc. (2) 37 (1) (2014), p.139-152.
- [6] S. Ianus and A.M.Pastore, harmonic maps on contact metric manifolds, Ann. Math. Blaise Pascal, 2 (1995), 43-53.
- [7] Domingo Chinea, harmonicity on maps between almost contact manifolds, Acta

Math. Hungar., in Acta Mathematica Hungarica 126(4) (2009) 352-365

[8] Aristide Ayibe, Harmonicity of (φ, φ') -holomorphic sections of a (*Riemannian*) almost contact fiber bundle of type I, Far East Journal of Mathematical sciences 121(2) (2019), p.169-197.

ARISTIDE AYIBE

UNIVERSITE FELIX HOUPHOUËT BOIGNY D'ABIDJAN, COTE D'IVOIRE., U.F.R: MATHEMATIQUES
INFORMATIQUE., LABORATOIRE: MATHEMATIQUES FONDAMENTALES

E-mail address: e-mail : aristideayibe@gmail.com