***Pseudomonas oleovorans* Endocarditis in a Child: The First Reported Case**

Pseudomonas oleovorans Endokarditi Olan Çocuk Hasta:
Literatürdeki İlk Olgu

Kamile Arıkan¹, Hayrettin Hakan Aykan², Ateş Kara¹, Ali Bülent Cengiz¹

¹ Department of Pediatrics, Hacettepe University School of Medicine, Ankara, Turkey

² Department of Pediatric Cardiology, Hacettepe University School of Medicine, Ankara, Turkey

Cite this article as: Arıkan K, Aykan HH, Kara A, Cengiz AB. *Pseudomonas oleovorans* endocarditis in a child: the first reported case. J Pediatr Inf 2018;12(3):e115-e117

Abstract

Infectious endocarditis (IE) is a rare, but potentially fatal disease in pediatric patients. Most patients have underlying hemodynamic problems that predispose them to the risks of IE. Modified Duke criteria are commonly used for clinical diagnosis of IE. Though *Streptococcus viridans* and *Staphylococcus aureus* are the main pathogens of the disease, other pathogens have also been isolated. Herein we describe the first reported case of infective endocarditis caused by *Pseudomonas oleovorans* in a girl with operated ventricular septal defect (VSD).

Keywords: *Pseudomonas oleovorans*, congenital heart defect, *Pseudomonas* spp., infective endocarditis

Öz

Enfektif endokardit (EE) çocuklarda nadir görülen ancak mortalitesi yüksek bir hastalıktır. Çoğu hastanın EE için risk oluşturan hemodinamik bozuklukları bulunmaktadır. EE klinik tanısı için çoğunlukla modifiye Duke kriterleri kullanılmaktadır. *Streptococcus viridans* ve *Staphylococcus aureus* en sık izole edilen etkenler olmakla birlikte, etyolojide farklı mikroorganizmalar da saptanabilmektedir. Burada ventriküler septal defekt nedeniyle opere olan bir kız hastada etken olarak *Pseudomonas oleovorans*'ın saptandığı literatürdeki ilk EE'li olgu sunulmaktadır.

Anahtar Terimler: *Pseudomonas oleovorans*, konjenital kalp hastalığı, *Pseudomonas* spp., enfektif endokardit

Introduction

Pseudomonas and related genera are aerobic, motile, non-spore-forming, nonfermentative, gram-negative bacilli that live in soil, in water and on plants and animals. Most organisms of these genera are ubiquitous and rarely pathogenic in humans. Although the *Pseudomonas* spp. may produce disease in any individual, they are usually opportunists that more commonly cause disease in patients with burns, cystic fibrosis, malignant diseases and immunodeficiency. While most infections

are caused by *Pseudomonas aeruginosa*, other *Pseudomonas* spp. may also give rise to severe infections less frequently. Infective endocarditis (IE) is a bacterial or fungal infection of the endocardium and/or heart valves that damages the endocardial tissue and/or valves. *Pseudomonas oleovorans* has been reported as a pathogen in sepsis, peritonitis, meningitis but it has not been reported as a pathogen in infective endocarditis (IE) (1-3). Herein we describe a case of infective endocarditis caused by *P. oleovorans* in a girl operated for ventricular septal defect (VSD).

Received: 21.02.2018

Accepted: 25.07.2018

Correspondence Address / Yazışma Adresi

Kamile Arıkan

Hacettepe Üniversitesi Tıp Fakültesi,
Çocuk Enfeksiyon Hastalıkları Bilim Dalı
Ankara-Türkiye

E-mail: kamile.arikan5@gmail.com

©Copyright 2018 by Pediatric
Infectious Diseases Society -Available online at
www.cocukenfeksiyon.org

©Telif Hakkı 2018

Çocuk Enfeksiyon Hastalıkları Derneği -Makale metnini
www.cocukenfeksiyon.org web sayfasından ulaşılabilir

Case Report

A 14 year-old female with a history of operation for VSD at 6 years old of age and insertion of pacemaker to the right atrium and ventricle at 7 years old of age due to antioventricular block, presented with fever persisting for 10 days which started after change of pacemaker. Due to persistent fever and increase in blood C-reactive protein (CRP), she had been hospitalized at an other center for 20 days. She was admitted to our center, because of fever persisting for 1 month, fatigue and chest pain. On admission, she was mildly ill appearing. Physical examination revealed a grade II-III/VI systolic ejection murmur. Laboratory evaluation noted bicytopenia (hemoglobin: 9.5 g/dL, thrombocyte count: $127 \times 10^3/\mu\text{L}$, leucocyte count: $5.9 \times 10^3/\mu\text{L}$), elevated erythrocyte sedimentation rate: 26 mm/h (0-20), increased CRP: 7.8 mg/dL (0-0.8), rheumatic factor positivity: 23.7 IU/mL (0-20), normal biochemistry and serum complements C3 and C4 levels. Transthoracic echocardiography (ECHO) revealed 5 x 5 mm and 17 x 4 mm vegetations next to tricuspid valve, at right ventricle. She was hospitalized due to infective endocarditis and sulbactam ampicillin and amikacin were started after blood cultures were taken. *P. oleovorans* was isolated from two of the blood cultures taken on admission. Cultured *P. oleovorans* was susceptible to amikacin, ciprofloxacin, colistin, gentamicin, imipenem, piperacillin tazobactam, but resistant to cefepime, ceftazidime and meropenem. Due to persistent fever and blood culture positivity, therapy was changed to piperacillin tazobactam plus amikacin. Blood culture was sterile at the seventh day of antibiotic treatment. She was operated at the thirteenth day of hospitalisation, infected pacemaker was extracted and residual VSD was closed. Tissue culture taken perioperatively from vegetation resulted as negative which was attributed to antibiotic therapy during blood culturing. Repeated transthoracic ECHO was obtained and there was no evidence of vegetation at the 21th day of therapy. After completion of 6 weeks of total antibiotic therapy, the patient was discharged and no complication was noted at control visits.

Discussion

Most children and adolescents with IE have an identifiable risk factor, and congenital heart defects (CHDs) as in our case, represent the major risk factor for IE today. CHDs are the most common birth defects worldwide, and affect approximately 1 per 100 live births (4). Survival of patients with CHDs has improved, and the population of children and adolescents with CHDs is expanding. Despite the improved survival, CHDs still represent an important cause of morbidity and mortality at young ages. A significant number of deaths occur among patients with infections. The previously reported incidence of IE in patients with CHDs is 15-140 times higher than that in the

general population, and CHDs also represent an important risk factor for increased mortality and complications in patients with IE (5). A particular form of IE is an infection where bacteria adhere to the leads of transvenously coronary implanted electronic devices, such as a pacemaker. In this type of infection, adherence of the bacteria to the foreign material and a subsequent formation of biofilm are believed to be important such as in our case.

Modified Duke criteria are commonly used for clinical diagnosis of IE (6). Though *Streptococcus viridans* and *Staphylococcus aureus* are the main pathogens of the disease, other pathogens have also been isolated (7). In a systematic review of 105 studies from 36 countries and a total of 33,214 cases *S. aureus* was found to be the most common microorganism, being the most frequent in 54.3% of studies (n= 57). Viridans group streptococci (VGS), coagulase-negative staphylococci (CNS), *Enterococcus* spp. and *Streptococcus bovis* were among the most common causative agents (8). Gram-negative pathogens account for less than 10% of all cases of infective endocarditis worldwide and endocarditis due to *Pseudomonas* spp. is rare (9).

In a study of 19 cases that fulfilled the criteria of definite endocarditis according to the modified Duke criteria, one (5.3%) blood culture was isolated as *P. aeruginosa* (10). The members of the genus *Pseudomonas* are known opportunistic pathogens. On the basis of 16S rRNA analysis, *Pseudomonas* spp. have been reorganized under various groups, each with several species. *P. oleovorans* was first described by Lee and Chandler in 1941 and taxonomically, it belongs to 16S rRNA group I (*P. aeruginosa* group) (11). *P. oleovorans* is a versatile bacterium capable of carrying out industrially important biocatalytic reactions such as oxidative assimilation of alkanes epoxidation of alkanes and fatty acids. *P. oleovorans* is an environmental isolate rarely detected in clinical specimens (12). Gilardi in 1972 reported the identification of this organism from many clinical specimens, but it was not considered pathogenic in any of these instances (13). Cowlshaw et al. reported the first case of *P. oleovorans* meningitis in an 8-month-old infant (1). Another case of peritoneal dialysis-associated peritonitis was reported by Hage et al (2). A case of septicemia due to *P. oleovorans* in a patient with disseminated staphylococcal infection who was on prolonged mechanical ventilation was also reported (3).

The diagnosis of IE is difficult and often made late, which highlights the need to consider the diagnosis of IE in any patient with a CHD and pacemaker, as the major risk factor for IE in our case, who presents with fever or other signs of infection. We conclude that *P. oleovorans* may be an important emerging non-*P. aeruginosa* or an under-diagnosed pathogen

in patients with underlying disease. It should be kept in mind that atypical and rare pathogens as in this case report can be causative agent. This report highlights the potential of unusual environmental species of *Pseudomonas* as opportunistic human pathogens. Microbiologists and clinicians should be aware of emerging pathogenic species of *Pseudomonas*, and these clinical conditions may predispose the patients to bacterial infection.

Informed Consent: Written consent was obtained from the patient's family.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept - KA; Design - KA; Supervision - HHA, AK, ABC; Data Collection and/or Processing - KA; Analysis and/or Interpretation - KA, HHA; Literature Review - KA; Writing - KA; Critical Review - HHA, AK, ABC.

Conflict of Interest: The authors have not reported a conflict of interest.

Financial Disclosure: No financial disclosure was declared by the authors.

References

1. Cowlishaw WA, Hughes ME, Simpson HC. Meningitis caused by an alkali-producing *Pseudomonas* spp. *J Clin Pathol* 1976;29:1088-90.
2. Hage JE, Schoch PE, Cunha BA. *Pseudomonas pseudoalcaligenes* peritoneal dialysis-associated peritonitis. *Perit Dial Int* 2013;33:223-4.
3. Gautam L, Kaur R, Kumar S, Bansal A, Gautam V, Singh M, et al. *Pseudomonas oleovorans* sepsis in a child: the first reported case in India. *Jpn J Infect Dis* 2015;68:254-5.
4. Leirgul E, Fomina T, Brodwall K, Greve G, Holmstrøm H, Vollset SE, et al. Birth prevalence of congenital heart defects in Norway 1994-2009 a nationwide study. *Am Heart J* 2014;168:956-64.
5. Verheugt CL, Uiterwaal CS, van der Velde ET, Meijboom FJ, Pieper PG, Veen G, et al. Turning 18 with congenital heart disease: prediction of infective endocarditis based on a large population. *Eur Heart J* 2011;32:1926-34.
6. Durack DT, Lukes AS, Bright DK. New criteria for diagnosis of infective endocarditis: utilization of specific echocardiographic findings. Duke Endocarditis Service. *Am J Med* 1994;96:200-9.
7. Quagliarello V. Infective endocarditis: global, regional, and future perspectives. *JAMA* 2005;293:3061-2.
8. Vogkou CT, Vlachogiannis NI, Palaodimos L, Kousoulis AA. The causative agents in infective endocarditis: a systematic review comprising 33,214 cases. *Eur J Clin Microbiol Infect Dis* 2016;35:1227-45.
9. Ambrosioni J, Hernandez-Meneses M, Tellez A, Pericàs J, Falces C, Tolosana JM, et al. The changing epidemiology of infective endocarditis in the twenty-first century. *Curr Infect Dis Rep* 2017;19:2.
10. Wei C, Wu KG, Syc L, Chen C, Tanga R. Infectious endocarditis in pediatric patients: analysis of 19 cases presenting at a medical center. *J Microbiol Immunol Infect* 2010;43:430-7.
11. Lee M, Chandler AC. A study of the nature, growth and control of bacteria in cutting compounds. *J Bacteriol* 1941;41:373-86.
12. Van Beilen JB, Eggink G, Enequist H, Bos R, Witholt B. DNA sequence determination and functional characterization of the OCT-plasmid-encoded alkJKL genes of *Pseudomonas oleovorans*. *Mol Microbiol* 1992; 6: 3121-36.
13. Gilardi GL. Infrequently encountered *Pseudomonas* species causing infection in humans. *Ann Intern Med* 1972;77:211-5.