Relating Brunn-Minkowski and Rogers-Shephard inequalities with the Minkowski asymmetry measure

Katherina von Dichter

Technische Universität München, Department of Mathematics
Boltzmannstraße 3, 85748 Garching by Munich, Germany
`dichter@ma.tum.de`

Abstract

In this paper we explain how the Minkowski asymmetry measure sharpens several classic results. Especially, we were able to tighten the Brunn-Minkowski and the Rogers-Shephard inequalities in terms of the Minkowski asymmetry measure using some stability results on those inequalities.

Keywords: Brunn-Minkowski inequality, Rogers-Shephard inequality, Minkowski asymmetry measure

MSC: 52A10, 52A38

1 Introduction

Let $\text{vol}(K)$ be the n -dimensional volume (or Lebesgue measure) of K . We will write $\text{vol}_n(K)$ whenever it is necessary to specify the dimension. Let $\mathcal{K}^n$ be the set of all convex compact sets in $\mathbb{R}^n$, $\mathbb{B}_2^n$ be the Euclidean ball of radius 1, $K + L$ be the Minkowski sum of convex bodies K and L , i.e. $K + L = \{a + b | a \in K, b \in L\}$. The Brunn-Minkowski inequality establishes that for any convex compact sets K, L holds

$$\text{vol}(K + L)^{\frac{1}{n}} \geq \text{vol}(K)^{\frac{1}{n}} + \text{vol}(L)^{\frac{1}{n}}.$$

Equality holds if and only if one of the three following cases are true (see [FiMaPr]):

- (i) in case for sets K and L with positive volume if and only if K is a homothet of L ,
- (ii) in case when one of them has volume 0, namely $\text{vol}(K) = 0$ and $\text{vol}(L) > 0$, then if and only if K is a singleton,
- (iii) in case when both of them have volume 0, then if and only if K and L are contained in parallel hyperplanes.

The Brunn-Minkowski inequality was proved in the 19th century by Brunn for compact convex sets in low dimensions ($n \leq 3$) (see [Bru]), and Minkowski for general compact convex sets in $\mathbb{R}^n$ (see [Mi]). With the time the Brunn-Minkowski inequality became the starting point of the Brunn-Minkowski theory and a powerful tool for problems involving metric quantities such as volume, surface area and mean width.

The Rogers-Shephard inequality (see [RoSh, Thm. 1]), which can be regarded as a reverse inequality to the Brunn-Minkowski inequality, yields

$$\text{vol}(K + L) \leq \binom{2n}{n} \frac{\text{vol}(K)\text{vol}(L)}{\text{vol}(K \cap (-L))}.$$

Moreover, equality holds if and only if $K = -L$ is an n -dimensional simplex.

The Minkowski asymmetry $s(K)$ of a convex compact sets K is the smallest rescaling of K which contains a translation of $-K$. It is known that $1 \leq s(K) \leq n$ with $s(K) = 1$ if and only if K is centrally symmetric, while $s(K) = n$ if and only if K is an n -dimensional simplex (see [BrKo, Cor. 2.7]).

Computing the volume is always a computationally hard task. However, computing the Minkowski asymmetry measure may be a computationally easy task (see [BrKo]). We wonder whether we could improve the Brunn-Minkowski and the Rogers-Shephard inequalities for prescribed $s(K) = s \in [1, n]$.

In order to start finding values of $c(s)$ and $C(s)$, we need to recall the proofs of the Brunn-Minkowski and the Rogers-Shephard inequalities. Secondly, we will show also the characterizations of their equality cases. Finally, we will show the ideas in order to obtain some stability results, in sense of near-equalities, of the Brunn-Minkowski and the Rogers-Shephard inequalities, which will help in providing bounds for $c(s)$ and $C(s)$ from some particular values of $s \in [0, 1]$.

The results presented in this paper are based on the master thesis (see [Di]).

1.1 Minkowski asymmetry measure

For an n -dimensional convex and compact set K , its Minkowski asymmetry (measure) $s(K)$ (cf. Minkowski 1911) is the smallest $\rho > 0$ such that $-K \subset x + \rho K$ for some $x \in \mathbb{R}^n$, i.e.

$$s(K) := \inf\{\lambda > 0 \mid K \subset x + \lambda(-K) \text{ for some } x \in \mathbb{R}^n\}.$$

Note that for Minkowski asymmetry $s(K)$ holds that $K \subset x + s(K)(-K)$ for some $x \in \mathbb{R}^n$. Figure 1 shows two examples of computing the Minkowski asymmetry measure.

Figure 1: $K \subset x + s(K)(-K)$ for some $x \in \mathbb{R}^n$ and $\Delta \subset y + 2(-\Delta)$ for some $y \in \mathbb{R}^n$

1.2 The Brunn-Minkowski inequality

Figure 2 shows the Minkowski sum of two 0-centered triangles.

Figure 2: Minkowski sum of the regular triangles Δ and ∇

We are now ready to state the Brunn-Minkowski inequality for compact sets, which gives a lower bound for $\text{vol}(K + L)$ in terms of $\text{vol}(K)$ and $\text{vol}(L)$.

Theorem 1 (Brunn 1887, Minkowski 1896). *For any compact sets K, L holds*

$$\text{vol}(K + L)^{\frac{1}{n}} \geq \text{vol}(K)^{\frac{1}{n}} + \text{vol}(L)^{\frac{1}{n}}. \quad (1)$$

The Brunn-Minkowski inequality (1) implies a well-known fact from Analysis, namely the Isoperimetric inequality.

Theorem 2 (Isoperimetric inequality). *Let K be a convex set in $\mathbb{R}^n$ and $P(K)$ its surface area. Then*

$$P(K) \geq n \text{vol}(\mathbb{B}_2^n)^{\frac{1}{n}} \text{vol}(K)^{\frac{n-1}{n}}.$$

The Isoperimetric inequality implies that among sets with a fixed surface area, Euclidean balls maximize the volume.

Next let us state the Rogers-Shephard inequality, which gives us an upper bound for $\text{vol}(K + L)$ in terms of $\text{vol}(K)$, $\text{vol}(L)$ and $\text{vol}(K \cap (-L))$. Note that it is in general impossible to find an upper bound for $\text{vol}(K + L)$ just in terms of $\text{vol}(K)$ and $\text{vol}(L)$.

Theorem 3 (Rogers and Shephard, 1958). *Let K, L be convex bodies. Then*

$$\text{vol}(K + L) \text{vol}(K \cap (-L)) \leq \binom{2n}{n} \text{vol}(K) \text{vol}(L). \quad (2)$$

There exist also another version of the Rogers-Shephard inequality, namely with projections and intersections.

Theorem 4 (Rogers and Shephard, 1958). *Let K be a compact body, H an i -dim subspace of $\mathbb{R}^n$. Then*

$$\text{vol}_i(P_H K) \max_{x \in \mathbb{R}^n} \text{vol}_{n-i}(K \cap (x + H^\perp)) \leq \binom{n}{i} \text{vol}_n(K). \quad (3)$$

An application of the Rogers-Shephard inequality (3) can be the following. One could approximate the volume $\text{vol}(K)$ of a compact set $K \in \mathbb{R}^n$ (for instance a brain tumor) by measuring the volume of the projection $\text{vol}_i(P_H K)$ and the volume of the biggest section $\max_{x \in \mathbb{R}^n} \text{vol}_{n-i}(K \cap (x + H^\perp))$.

In order to see that (3) implies (2), we introduce the set $M := \{(z_1, z_2)^T \in \mathbb{R}^{2n} | z_2 \in K, z_1 \in z_2 + L\}$. Using the definition of the set M it can be easily seen that $P_H M = K + L$, $M \cap H^\perp = K \cap (-L)$ and $\text{vol}_{2n}(M) = \text{vol}_n(K) \text{vol}_n(L)$. And therefore two versions of the Rogers-Shephard inequality (2) and (3) are equivalent.

Choosing $L = -K$ and combining the Rogers-Shephard and the Brunn-Minkowski inequalities leads to

$$2^n \leq \frac{\text{vol}(K - K)}{\text{vol}(K)} \leq \binom{2n}{n}.$$

We aim to find constants $c(s), C(s)$ with $2^n \leq c(s) \leq C(s) \leq \binom{2n}{n}$, such that for every convex body K such that $s(K) = s \in [1, n]$ holds

$$c(s) \leq \frac{\text{vol}(K - K)}{\text{vol}(K)} \leq C(s).$$

In order to do it we study the equality cases of the Brunn-Minkowski inequality (1) and the Rogers-Shephard inequalities (2) and (3), their stability results and then involve the Minkowski asymmetry measure in that inequalities.

1.3 Equality case of the Brunn-Minkowski inequality

The equality case of the Brunn-Minkowski inequality (1) is the following.

Theorem 5 (Klain, 2011, Kneser, Süss, 1932). *Let K and L be convex compact sets with positive volumes. Then*

$$\text{vol}(K + L)^{\frac{1}{n}} = \text{vol}(K)^{\frac{1}{n}} + \text{vol}(L)^{\frac{1}{n}}$$

holds if and only if K and L are homothets.

Moreover, if exactly one of the sets is lower dimensional, namely $\text{vol}(K) = 0$, and $\text{vol}(L) > 0$, then equality holds if and only if K is a singleton; and if $\text{vol}(K) = \text{vol}(L)$, then equality holds if and only if K and L are contained in parallel hyperplanes.

1.4 Stability of the Brunn-Minkowski inequality

We consider the stability as a sharpening of the inequality at the near-quality case. Let $K \triangle L$ be the symmetric difference of K and L , i.e. $K \triangle L = (K \setminus L) \cup (L \setminus K)$. For the Brunn-Minkowski inequality (1) it holds that

Theorem 6 (Figalli, 2009). *Let K and L be convex compact sets in $\mathbb{R}^n$. Then*

$$\text{vol}(K + L)^{\frac{1}{n}} \geq \left(\text{vol}(K)^{\frac{1}{n}} + \text{vol}(L)^{\frac{1}{n}} \right) \left(1 + \frac{A(K, L)^2}{c(n)\sigma(K, L)^{\frac{1}{n}}} \right)$$

with $A(K, L) := \inf_{x \in \mathbb{R}^n} \left\{ \frac{\text{vol}(K \triangle (x + \lambda L))}{\text{vol}(K)} \right\}$, $\lambda = \left(\frac{\text{vol}(K)}{\text{vol}(L)} \right)^{\frac{1}{n}}$, $\sigma(K, L) := \max \left\{ \frac{\text{vol}(K)}{\text{vol}(L)}, \frac{\text{vol}(L)}{\text{vol}(K)} \right\}$ and $c(n) = 14n^2 4^{n-1}$.

One can see that due to the stability result in case of near-equality of (1) we have that K and L are almost homothets. Therefore $\text{vol}(K \triangle (x + \lambda L)) \approx 0$ and therefore $A(K, L) \approx 0$.

1.5 Equality case of the Rogers-Shephard inequality

The equality case of the Rogers-Shephard inequality (3) can be stated as following.

Theorem 7 (Rogers and Shepard, 1958). *Let K be a convex compact sets in $\mathbb{R}^n$, H an i -dim subspace of $\mathbb{R}^n$ and $x \in \mathbb{R}^n$. Then*

$$\text{vol}_i(P_H K) \max_{x \in \mathbb{R}^n} \text{vol}_{n-i}(K \cap (x + H^\perp)) = \binom{n}{i} \text{vol}_n(K)$$

if and only if

$$K \cap (x + H^\perp) =_h K \cap H^\perp \quad \forall x \in \mathbb{R}^n$$

and

$$\text{vol}_{n-i}(K \cap (x + H^\perp)) = (1 - \|x\|_K)^{n-i} \text{vol}_{n-i}(K \cap H^\perp).$$

Next we present a more recent result that is the equality case of the Rogers-Shephard inequality (2).

Theorem 8 (Alonso-Gutierrez, Jimenez, Villa, 2013). *Let K and L be convex compact sets in $\mathbb{R}^n$. Then*

$$\text{vol}_n(K + L) \text{vol}_n(K \cap (-L)) = \binom{2n}{n} \text{vol}_n(K) \text{vol}_n(L)$$

if and only if K and L are simplices such that $K = -L$.

1.6 Stability of the Rogers-Shephard inequality

Let $d_{BM}(K, L) := \min\{\rho \geq 1 | K \subset A(L) \subset x + \rho K \text{ for some } x \in \mathbb{R}^n\}$ with $A(L)$ being any affine transformation of L be the Banach-Mazur distance of K with respect to L and T be an n -simplex.

Theorem 9 (Boroczsky, 2005). *Let K be a convex compact sets in $\mathbb{R}^n$. If*

$$\text{vol}(K - K) = (1 - \varepsilon) \binom{2n}{n} \text{vol}(K),$$

then

$$1 + \frac{1}{n} \varepsilon \leq d_{BM}(K, T) \leq 1 + n^{50n^2} \varepsilon.$$

1.7 Results

Let us recall that the Brunn-Minkowski and Rogers-Shephard inequalities for convex compact sets K and $-K$ state that

$$2^n \leq \frac{\text{vol}(K - K)}{\text{vol}(K)} \leq \binom{2n}{n}.$$

Moreover, $\frac{\text{vol}(K - K)}{\text{vol}(K)} = 2^n$ if and only if K is symmetric and $\frac{\text{vol}(K - K)}{\text{vol}(K)} = \binom{2n}{n}$ if and only if K is an n -dimensional simplex.

But at the same time we have that

$$1 \leq s(K) \leq n.$$

Moreover, $s(K) = 1$ if and only if K is symmetric and $s(K) = n$ if and only if K is an n -dimensional simplex.

Combining those facts enable us to involve the asymmetry measure into these inequalities. We state some improvements on the Brunn-Minkowski and the Rogers-Shephard inequalities by means of the Minkowski asymmetry measure (see [Di]).

Theorem 10. *Let K be a convex compact set in $\mathbb{R}^n$ and $s = s(K)$. Then*

$$c(s) \geq 2^n \left(1 + \frac{1}{n 4^{n-1}} \left(\frac{(s-1)^n \text{vol}_{n-1}(\mathbb{B}_2^{n-1})}{2^{n-1} n^{2n} \text{vol}_n(\mathbb{B}_2^n)} \right)^2 \right)^n \quad (4)$$

and

$$C(s) \leq (1 + s)^n. \quad (5)$$

Moreover, if $n - \frac{1}{4n} < s < n$, then

$$c(s) \geq \binom{2n}{n} (1 - 4n^2(n - s)) \quad \text{and} \quad C(s) \leq \binom{2n}{n} \left(1 - \frac{n - s}{n^{1+50n^2}} \right). \quad (6)$$

It is worth mentioning that (4) and (5) (resp. (6)) are specially good when $s(K) \approx 1$ (resp. $s(K) \approx n$).

Theorem 11. *The diagram $f(\mathcal{K}^n)$, where $f : \mathcal{K}^n \rightarrow [1, n] \times \left[2^n, \binom{2n}{n}\right]$ is given by $f(K) := \left(s(K), \frac{\text{vol}(K-K)}{\text{vol}(K)}\right)$, is simply connected, contains $(1, 2^n)$ and $(n, \binom{2n}{n})$.*

We now investigate $f(\mathcal{K}^2)$.

Remark 1. *Let $T = \text{conv}(\{(0, 1)^T, (\pm\sqrt{3}/2, -1/2)^T\})$, let $s \in [1, 2]$, and let*

$$K_s := T \cap (-sT) \quad \text{and} \quad C_s := \text{conv}(T \cup (-sT)).$$

Then $s(K_s) = s(C_s) = s$, $\text{vol}(K_s) = \frac{2s-(s-1)^2}{4}$, $\text{vol}(K_s - K_s) = \frac{(s+1)^2}{2}$, $\text{vol}(C_s) = \frac{3\sqrt{3}}{2}s$, and $\text{vol}(C_s - C_s) = 3\sqrt{3}s(1+s)$.

We finally provide upper and lower bounds for the constants $c(n), C(n)$ in the planar case derived from the Theorem 10 and Remark 1.

Corollary 1. *Let $K \in \mathcal{K}^2$ and let $s = s(K)$. Then*

$$4 \left(1 + \frac{(s-1)^4}{2^{11}\pi^2}\right)^2 \leq c(s) \leq 2 \frac{(s+1)^2}{2s - (s-1)^2}$$

and

$$2(s+1) \leq C(s) \leq (1+s)^2.$$

Moreover, if $s > \frac{8}{7}$, then

$$6(16s - 31) \leq c(s) \quad \text{and} \quad C(s) \leq 6 \left(1 - \frac{2-s}{2^{201}}\right).$$

Figure 3: The diagram in Theorem 11 and the bounds obtained in Theorem 11 (dashed blue and red lines, respectively) and Corollary 1 (dashed black lines) in the case of $n = 2$; the light grey area is obtained in Theorems 1 and 3.

References

- [AlArGoJiVi] D. Alonso-Gutiérrez, S. Artstein-Avidan, B. González Merino, C. H. Jiménez, R. Villa, Rogers-Shephard and local Loomis-Whitney type inequalities, arXiv:1706.01499, 2017.
- [AJV] D. Alonso-Gutiérrez, C.H. Jiménez, R. Villa, Brunn–Minkowski and Zhang inequalities for convolution bodies, *Adv. Math.*, 238 (2013), 50–69.
- [BiGaGr] G. Bianchi, R.J. Gardner, P. Gronchi, Symmetrization in Geometry, *Adv. Math.*, 36 (2017), 51–88.
- [Bl] W. Blaschke, Eine Frage über konvexe Körper, *Jahresber. Deutsch. Math.-Verein.*, 25 (1916), 121–125.
- [Bo] F. Bohnenblust, Convex regions and projections in Minkowski spaces, *Ann. of Math.* 39 (1938), no. 2, 301–308.
- [Bor] K. Boroczsky, The stability of the Rogers-Shephard inequality and some related inequalities, *Adv. Math.*, 190 (2005), no. 1, 47–76.
- [BrG] R. Brandenburg, B. González Merino, A complete 3-dimensional Blaschke-Santaló diagram, *Math. Ineq. Appl.* 20 (2017), no. 2, 301–348.
- [BrG2] R. Brandenburg, B. González Merino, The asymmetry of complete and constant width bodies in general normed spaces and the Jung constant, *Israel J. Math.* 218 (2017), no. 1, 489–510.
- [BrG3] R. Brandenburg, B. González Merino, Minkowski concentricity and complete simplices, *J. Math. Anal. Appl.* 454 (2017), no. 2, 981–994.
- [BrKo] R. Brandenburg and S. König, No dimension-independent core-sets for containment under homothetics, *Discrete Comput. Geom.*, 49 (2013), no 1, 3–21.
- [Bru] H. Brunn, Über Ovale und Eiflächen, Inaugural Dissertation, München (1887).
- [BrK2] R. Brandenburg, S. König, Sharpening geometric inequalities using computable symmetry measures, *Mathematika*, 61 (2015), no. 3, 559–580.
- [Da] P.J. Davis, 6. Gamma function and related functions, in Abramowitz, Milton; Stegun, Irene A., *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, New York: Dover Publications, 1972.

- [Di] K. von Dichter, Volume estimates via the Asymmetry Measure of Minkowski, Master Thesis (2018).
- [Fe] H. Federer, Geometric measure theory, Die Grundlehren der mathematischen Wissenschaften, Band 153, New York: Springer-Verlag New York Inc., pp. xiv+676, 1969.
- [FiMaPr] A. Figalli, F. Maggi, A. Pratelli, A refined Brunn–Minkowski inequality for convex sets, *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 26 (2009), no. 6, 2511–2519.
- [FiMaPr2] A. Figalli, F. Maggi, A. Pratelli, A mass transportation approach to quantitative isoperimetric inequalities, *Invent. Math.*, 182 (2010), no. 1, 167–211.
- [Gl] E. D. Gluskin, Diameter of the Minkowski compactum is approximately equal to n , *Funct. Anal. Appl.* 15 (1981), 57–58.
- [GoHe] B. González Merino, M. A. Hernández Cifre, El teorema del elipsoide de John, <http://hdl.handle.net/10201/30278>, Universidad de Murcia, 2009.
- [Gr] B. Grünbaum, Measures of symmetry for convex sets, *Proc. Sympos. Pure Math.* 7 (1963), 233–270.
- [GuJ] Q. Guo, H. Jin, Asymmetry of convex bodies of constant width, *Discrete Comput. Geom.* 47 (2012), no. 2, 415–423.
- [GuK] Q. Guo, S. Kaijser, On asymmetry of some convex bodies, *Discrete Comput. Geom.* 27 (2002), no. 2, 239–247.
- [HaOh] H. Hadwiger, D. Ohmann, Brunn-Minkowskischer Satz und Isoperimetrie, *Math. Z.*, 66 (1956), no. 1, 1–8.
- [Kl] D. A. Klain, On the equality conditions of the Brunn-Minkowski theorem, *Proc. Amer. Math. Soc.*, 139 (2011), no. 10, 3719–3726.
- [La] M. Lassak, Approximation of convex bodies by inscribed simplices of maximum volume, *Beitr. Algebra Geom.* 52 (2011), no. 2, 389–394.
- [McGu] R. J. McCann, N. Guillen, Five lectures on optimal transportation: geometry, regularity and applications. *Analysis and geometry of metric measure spaces*,

- Lecture notes of the séminaire de Mathématiques Supérieure (SMS) Montréal, 145–180, 2011.
- [Mc] P. McMullen, Inequalities between intrinsic volumes, *Monatsh. Math.*, 111 (1991), no. 1, 47–53.
- [Mi] H. Minkowski, *Geometrie der Zahlen*, Leipzig: Teubner, 1896.
- [RoSh] C. A. Rogers, G. C. Shephard, Convex bodies associated with a given convex body, *J. Lond. Math. Soc.*, 1 (1958), no. 3, 270–281.
- [Ru] M. Rudelson, Distance between non-symmetric convex bodies and the MM^* -estimate, *Positive* 4 (2000), no. 2, 161–178.
- [Sa] L. Santaló, Sobre los sistemas completos de desigualdades entre tres elementos de una figura convexa plana, *Math. Notae* 17 (1961), 82–104.
- [Sch] R. Schneider, *Convex bodies: the Brunn–Minkowski theory*, no. 151, Cambridge university press, 2014.
- [So] V. Soltan, A characterization of homothetic simplices, *Discr. Comput. Geom.*, 22 (1999), no. 2, 193–200.
- [St] P. Steinhagen, Über die größte Kugel in einer konvexen Punktmenge, *Abh. Hamb. Sem. Hamburg* 1 (1921), 15–26.
- [TJ] N. Tomczak-Jaegermann, *Banach-Mazur Distances and Finite-Dimensional Operator Ideals*, Pitman Monographs and Surveys in Pure and Applied Mathematics 38, Longman Scientific and Technical, New York (1989).