27. Ross WD, Marfell-Jones RJ (1995). Cinantropometria. In Duncan J, MacDougall H., Wenger A., Green HJ (eds.), *Evaluación fisiológica del deportista*, Paidotribo. Barcelona.
28. Seabra A, Maia J, Garganta R. (2001). Crescimento, maturação, aptidão física, força explosiva e habilidades motoras específicas. Estudo em jovens futebolistas e não futebolistas do sexo masculino dos 12 aos 16 anos de idade. *Ver Port Cie Desporto* 1: 22-35.
29. Semenick D (1990). The T-Test. *NSCA J* 1(1): 36-37.
30. Torres-Unda J, Zarrazquin I, Gil J, Ruiz F, Irazusta A, Kortajarena M, Seco J, Irazusta J (2013). Anthropometric, physiological and maturational characteristics in selected elite and non-elite male adolescent basketball players. *J Sports Sci* 31(2): 196-203.
31. Torres-Unda J, Zarrazquin I, Gravina L, Zubero J, Seco J, Gil S, Gil J, Irazusta J (2016). Basketball performance is related to maturity and relative age in elite adolescent players. *J Strength Cond Res* 30(5): 1325-1332.
32. Vaeyens R, Malina RM, Janssens M, Van Renterghem B, Bourgois J, Vrijens J, Philippaerts RM (2006). A multidisciplinary selection model for youth soccer: the Ghent Youth Soccer Project. *Br J Sports Med* 40: 928-934.

AUTHOR:José A Rodrigues dos Santos ¹

¹ CIFI²D – Faculty of Sport
University of Porto – Portugal

<https://doi.org/10.5628/rpcd.17.01.75>

Nutrition for Soccer**KEY WORDS:**

Carbohydrate. Fat. Protein.

Fibers. Vitamins. Minerals.

SUBMISSÃO: 15 de Novembro de 2016

ACEITAÇÃO: 03 de Abril de 2017

ABSTRACT

Soccer is characterized by repeated sprint ability which is conditioned by the efficiency of both the anaerobic and aerobic energy systems. Energy supply for soccer performance relies directly on the nutritional habits, usually inadequate in many soccer players. So, correct nutrition is a crucial factor toward soccer highest performance. The type and quantity of foods and timing of meals can influence both sport performance and recovery's capacity. Optimal nutrition contributes to improve physical conditioning, mental accuracy and faster recovery after workouts or matches. Soccer players need to achieve good nutritional habits in order to get the body weight related to the maximum performance and to maintain health. This short review intended to establish some nutritional guidelines which can help soccer players to achieve and maintain an optimal body composition and maximize energy availability for training and matches.

Corresponding: José Augusto Rodrigues dos Santos, Faculdade de Desporto da Universidade do Porto, Rua Dr. Plácido Costa, 91, 4200-450 Porto, Portugal; jaugusto@fade.up.pt

RESUMO

O futebol é caracterizado pela capacidade de sprint repetida a qual está condicionada por ambos os sistemas energéticos – aeróbio e anaeróbio. A provisão de energia para a performance no futebol apoia-se diretamente nos hábitos nutricionais que são usualmente inadequados em muitos jogadores de futebol. Por isso, uma nutrição correta é um fator crucial para a consecução da mais alta performance no futebol. O tipo e quantidade de alimentos e o *timing* das refeições podem influenciar quer a performance desportiva quer a capacidade de recuperação. Uma nutrição ótima contribui para a melhoria da condição física, da acurácia mental e uma rápida recuperação após treinos e jogos. Os jogadores de futebol necessitam de adquirir de forma a obter o peso corporal mais ajustado à obtenção da máxima performance bem como para a manutenção da saúde. Esta curta revisão pretende estabelecer algumas regras nutricionais que podem ajudar os jogadores de futebol a obter uma ótima composição corporal e maximizar a energia disponível para o treino e competição.

PALAVRAS CHAVE:

Hidratos de carbono. Gorduras.

Proteínas. Fibras. Vitaminas. Minerais.

INTRODUCTION

Soccer is an extraordinary social phenomenon mobilizing millions of people and acting as an important financial and economic lever in modern societies. The increasing social importance of soccer determined its professionalization and new exigencies for training and recovery.

Soccer's performance is a construct derived from the harmonious conjugation of the three main training factors: load-stimulus (exercise), rest and nutrition. While physical exertion is under coach's responsibility, rest and nutrition lies mainly on player's responsibility. So, adjusted rest and proper nutrition are important means to match the physical challenge imposed by soccer training and competition. How better you eat how better you fit. Nutrition is one of the basements for physical and mental performance.

Proper eating presupposes the selection of the better food materials which are fundamental for implementing a correct nutrition status, essential to maintain health and to improve physical conditioning in soccer players.

Overwhelming evidence supports the importance of the nutrition as *sine qua non* condition for soccer's performance.

Soccer's highest performance is correlated to high training intensity which promotes increased energy requirements. High levels of training and competition elicit increased nutritional demands.

With this review, we intend to focus on the nutritional needs for soccer's performance, trying to highlight the main aspects for an adequate nutrition for soccer players.

SOCCER CHARACTERIZATION

Soccer involves several muscular and organic capacities as well as technical and tactical qualities. Soccer can be described as stochastic (randomly determined), acyclical, and intermittent activity expressed through its variability and unpredictability ⁽¹⁰⁾. Soccer is a physically demanding sport that is characterized by intermittent bursts of high-intensity effort superimposed upon a background of continuous movement lasting approximately 90 min where players can cover distances between 9 and 12 km. The continuity of the effort during the game progressively increases fatigue, as is evidenced by greater distances being covered in the first versus second half and that sprint performance decreases over the match period. It was estimated for elite soccer players (outfield players) an average work rate during the match corresponding to approximately 70% of the maximal oxygen uptake ⁽⁴⁾. About 90% of the energy production corresponds to the aerobic metabolism. It has been estimated that 80-90% of the effort corresponds to low to moderate intensity activities whereas 10-20% corresponds to high intensity activities ⁽⁴⁾.

Although soccer games and practices are mainly supported by aerobic production of energy, the quality of soccer performance is given by the short bursts demanding anaerobic energy production. Elite soccer players perform 150-250 brief intense actions during a game⁽⁵⁾ which give the extent of the anaerobic contribution to the overall energy production.

Trying to elucidate which are the specific attributes and skills which characterize a successful soccer player, several soccer coaches gathered in England by UEFA (1977) established the following hierarchy: First, tactics; second, technical ability; and third, physical attributes. So, training must search for the tactical matrix of the team and the development of other factors will be the result of tactical realization. The crucial quality of the game is under tactical supervision. Tactical performance in soccer must be the projection of the coach's idea for the game. This is the main objective for training and determines team's competitive level. However, tactics are conditioned by technical skills and both supported by the physical conditioning (cardiorespiratory fitness, general and functional muscular strength, endurance, and power).

The individual skills and recurrent movements in soccer are diverse (passing and striking, receiving, dribbling, feinting, tackling, jumping, heading and throwing, accelerating and turning) and support different actions (defence, attack, patterns of play or strategy, set plays) which are the "soul" of the game.

Soccer training must respect several principles with special emphasis to the relationship between practice and recovery. Sometimes players have not an adequate period of rest between seasons and after hard workouts and matches. Nowadays, at top level, soccer players are continuously engaged in national and international games, travelling to places with different environmental conditions (temperature, humidity, altitude, etc.) which put increased stress to the players. So, in these conditions, nutrition and hydration gain special importance and can contribute for better recovery, avoiding situations of overreaching or overtraining.

ENERGY DEMANDS ON SOCCER

Human diet provides energy mainly from carbohydrates and fat. Carbohydrates, whether sugars or starches, are broken down in the body and stored as glycogen – a polymer of α -glucose. Some of the glycogen is stored in the liver which can also support skeletal muscle work but is mainly directed to brain and nervous system energy supply. The major amount of glycogen is stored in the muscles. Water is also stored with the glycogen; three grams of water for each gram of glycogen⁽²⁹⁾. Weight loss noticed after only one or two training sessions is a combination of the fuel burned up (minor contribution for weight loss) and the stored water lost as sweat (major contribution for weight loss). Even if the athlete drinks *ad libitum*, water loss is not completely replaced immediately after prolonged exercise.

Fat is stored in the adipose tissue and muscle cells. Adipose tissue cells are simply storage containers that expand when increase fat or carbohydrate intake and contract during fasting. Fat molecules are broken down to release fatty acids and glycerol. These can then be carried by the blood to the muscles (fatty acids) and liver (glycerol) for further use by other cells. The body fat content of an average lean adult man is about 15% and for lean adult women about 25%. Elite endurance runners have much less body fat, but even so it is enough to provide their muscles with fuel for several days of low intensity exercise. European elite soccer players have a percentage of body fat on average ranging from 10 to 12%^(58, 65).

During exercise, muscles use energy at a rate directly proportional to intensity. During exercise the working muscles convert chemical energy into kinetic energy and heat. Heat must be dissipated through thermoregulatory mechanisms.

Energy is aerobically produced when muscle cells oxidize carbohydrate and fatty acids to make ATP (aerobic metabolism). ATP is actually the direct source of muscle energy. Muscle ATP is never depleted. Even after a very intense exercise bout ATP concentration only declines from ≈ 25.0 to ≈ 16.0 mmol.kg⁻¹ (dry weight) while Phosphocreatine, the immediate source for ATP resynthesis, declines largely from ≈ 79.0 to ≈ 20.0 mmol.kg⁻¹ (dry weight). ATP concentration seems to be insensitive to training⁽⁷⁾. ATP can also be produced through anaerobic metabolism (phosphocreatine and glycolytic pathways).

The selection of the different energy production pathways depends on many factors:

Exercise intensity. Games like soccer involve brief periods of high intensity efforts mixed with periods of rest or low intensity exercise. As energy demands change so the body will rely on the different fuel sources.

Player's physical conditioning. Individuals vary considerably on their capacity to perform exercise and in their metabolic response to exercise.

Muscles' metabolic profile. Some muscles work more aerobically, which means they can use fat or carbohydrate as a fuel. Others work mainly anaerobically which are mainly supported by carbohydrate metabolism.

Training effect. Training alters muscles' metabolic profile. Training enables the muscles to take up more oxygen from the blood supply and hence produce more aerobic energy. The increased oxygen utilization by the body is a key factor in determining fuel use and performance efficiency.

Although, aerobic metabolism is important for soccer, the anaerobic sources of energy support the skills expressed by power and velocity.

ADEQUATE NUTRITIONAL INTAKE IS CRUCIAL FOR SOCCER PERFORMANCE

Food provides the essential nutrients to keep body functions, maintaining life and growth. Food is constituted by five essential nutrients: protein (essential for growth and repair of body tissues), fat (insulation and energy for body functions and physical activity), carbohydrate (main source for high intensity physical activities), minerals and vitamins (both control metabolic processes; they act as bio regulators). Although not a nutrient, water is essential to life as it helps the body to metabolize nutrients efficiently. With reduced nutritional value, fiber is an important component in a healthy diet contributing for the good functioning of the gut besides other important health functions. Nutritional cares for soccer players are directed for – refuelling, rehydration, and repair.

WATER

Water is not a nutrient but is fundamental for life, with special relevance in hot environments. An adequate water intake is crucial for a balanced body metabolism and thermoregulation. The major mechanism for thermoregulation during soccer is sweating. The metabolic efficiency during physical activity (i.e., the transformation of the energy stored in muscle into mechanical work) is about 20%, liberating 80% of energy as heat. About 50% of the energy from ATP hydrolysis is dissipated as friction between the motor and its substrate. Conformational changes of muscle filaments lead to additional energy dissipation ⁽¹¹⁾. One liter of water lost as sweat, at 30° C corresponds to 580 kcal lost as heat ⁽⁶⁸⁾.

Water is supplied by the ingestion of both foods and drinks and is also produced by the body during oxidation ⁽⁵¹⁾. The complete oxidation of 100g lipids yields about 107 ml water while a similar weight of carbohydrates yields 55 ml and of proteins about 41 ml water but is roughly offset by respiratory water losses ⁽⁶²⁾. Around two litres of water are needed daily to replenish water lost through breathing, sweating and excretion of liquids and solids.

Water is lost through breathing (500 ml), sweating (600 ml in temperate climate), urine (750 ml) and faeces (150 ml). The amount of water lost through sweating depends on thermoregulatory strain. Water lost through sweating increases salt concentration and triggers the sensation of thirst. When the quantity of water is adjusted to the body's needs, thirst sensation is cancelled. Large amounts of plain water intake can induce water intoxication (hyposmolality/hyponatremia) affecting health and performance ⁽⁴³⁾. Several factors influence sweating during exercise – environment's temperature, humidity, wind, intensity of sunshine, clothing, exercise duration and intensity, player's training level, body composition, etc.

The sweating rate is lower in people acclimatized to extreme hot conditions. However, dehydration can occur if the water loss is not compensated by fluids ingestion. Water balance during soccer's games and training needs to be solved individually.

WATER AND EXERCISE

Body's capability to store heat is very short; so, during exercise blood flow to the skin increases toward the elimination of the heat produced by exercising muscles. However, during high intensity exercise cardiac output is insufficient to meet the demands of both muscle and skin blood flow and, the latter will be reduced affecting heat dissipation and promoting a rapid rise in body temperature. Earlier and systematic fluid ingestion can counterbalance the negativity of these alterations.

It is generally accepted that warming up is a crucial step before training and competition. In hot and humid climates warming up must be shortened to avoid extensive water lost. Moreover, excessive timing for warming up reduces muscle glycogen stores what can impair physical performance in the latest periods of the game.

People exercising hardly (high volume of training and high load's intensity) also need to ensure adequate water intake, as dehydration can cause the body to overheat. Endurance events, like soccer matches, are deeply affected when ambient temperature and humidity are high. As the ambient temperature and humidity increases, sweating rate increases too and peripheral blood flow must also increase to limit the rise in body temperature. Both dehydration and hyperthermia can negatively impair soccer performance ⁽³⁸⁾.

Thirst is an unreliable indicator of water loss during exercise as the thirst mechanism lags behind actual need. It is therefore important to drink before thirst occurs. Assessing body weight before and after exercise permits the soccer player calculate how much water is needed to rehydrate.

Every 0.5 kg of weight lost in sweat is equivalent to 378 ml of water and this needs to be replaced before, during and after exercise to maintain performance; however, water ingestion for rehydration must be higher than water lost by sweating. The athlete must drink between 450 to 675 mL of fluid for every 0,5 kg of body weight lost during exercise ⁽⁴⁰⁾. For better rehydration fluid replacement must be got through beverages that are palatable to the athlete.

Soccer players must start the match fully hydrated and, during the match, mainly when the environmental conditions are extremely hot, must ingest carbohydrate-electrolyte containing drinks which are the best for fluid replacement adding some energy for muscle function. Usually, fluids in sport add some nutrients to water providing both water and energy. The rates at which substrate and water can be supplied during exercise are limited by gastric emptying and intestinal absorption.

Increasing the carbohydrate content of fluid slows the rate of gastric emptying; however, when the gut receives both glucose and sodium, water absorption is increased. Table 1 shows the characteristics of the different drinks. Table 2 show the differences on the absorption rate of different commercial sport drinks compared to water when exercising ⁽³⁸⁾.

TABLE 1. Composition of the three sports drinks (adapted from Hill et al. ⁽³⁸⁾)

	DRINK 1 (PER 100 ML)	DRINK 2 (PER 100 ML)	DRINK 3 (PER 100 ML)
Energy	29 kcal	32 kcal	21 kcal
Carbohydrate (g)	6.7	8.0	5.0
Type (g)	Maltodextrin (1.9)	Sucrose (6.0)	Glucose (5.0)
	Sucrose (3.0)	Maltodextrin (2.0)	
	Fructose (1.3)		
	Glucose (0.5)		
Sodium (mg)	41.0	25.0	23.0
Potassium (mg)	11.7	14.0	19.5
Magnesium (mg)	0	0	2.4

TABLE 2. Exercise group mean ($\pm$ SEM) drink absorption parameters (adapted from Hill et al. ⁽³⁸⁾)

	DRINK 1	DRINK 2	DRINK 3	WATER
T1 (min)	14.3 $\pm$ 1.2 ^a	13.4 $\pm$ 1.0 ^a	14.2 $\pm$ 1.2 ^a	19.8 $\pm$ 1.3
T2 (min)	40.7 $\pm$ 6.3	49.0 $\pm$ 9.6	41.1 $\pm$ 5.8	41.2 $\pm$ 5.2
T1/2 (min)	17.7 $\pm$ 1.6	19.7 $\pm$ 2.6	17.7 $\pm$ 1.8	20.3 $\pm$ 1.6

T1 – time at which the absorption rate was maximum
T2 – time when the rate of absorption returned to zero
T1/2 – time when 50% of the drink had been absorbed
^a significantly different from water

Although the study from Hill et al. ⁽³⁸⁾ had verified that water and some commercial sports drinks are similar in the rate of hydration, it clearly concludes that the enrichment in carbohydrates and some electrolytes don't delay the absorption rate. So, soccer players can hydrate adequately with commercial drinks, the chose depending on the palatability and on the flavour the athlete enjoys more.

What thinking about beer for rehydration? Castro-Sepulveda et al. ⁽¹⁸⁾ stated that non-alcoholic beer rehydrates while maintaining electrolyte homeostasis; however, alcoholic beer reduced plasma Na⁺ and increased plasma K⁺ during exercise, what can be deleterious

for health and physical performance. Alcoholic drinks must be completely avoided before, during and immediately after exercise. Rehydration must be achieved through non-alcoholic drinks. Lately in this paper we shall analyse some deleterious effects of alcoholic drinks.

PROTEIN

Body proteins are in a dynamic state, with tissue proteins and other nitrogenous compounds being resynthesized and degraded continuously.

Protein is used by the body for the growth and repair of tissues such as skin, muscle, internal organs, hair and nails. Proteins are built from some twenty amino acids (building blocks) arranged in many different combinations. Amino acids are called “essential”, because they cannot be synthesized in the body and must be provided by the diet; and “non-essential” which can be built within the body from the contribution of essential amino acids. Twelve amino acids are non-essential: alanine, arginine, aspartic acid, cysteine, cystine, glutamic acid, glutamine, glycine, ornithine, proline, serine, taurine and tyrosine. The essential amino acids are leucine, isoleucine, valine, threonine, lysine, methionine, phenylalanine and tryptophan.

Protein can be obtained from both vegetable and animal sources. Although the most common protein sources are meat, milk, eggs, and fish, significant amounts of protein can be obtained from foods such as beans, nuts, cereals and some vegetables. Animal proteins have more biological and nutritional value because their amino acid profile is closer to human requirements.

The recommended dietary allowance (RDA) for protein for adults is 0.8 g/kg of body weight per day. For a 70-kg man, this is 56 g of protein per day, and for a 55-kg woman, 44 g. For children, adolescents, and pregnant or lactating women, the RDA is slightly higher to ensure a satisfactory growth rate (JADA, 2000). This is the amount present in a 160-185 g lean grilled steak, 180-210 g hard cheese, 7-8 eggs or 17-20 slices of wholemeal bread. Western diet currently provides daily more than these values.

PROTEINS AND EXERCISE

Many athletes, especially those participating in power sports and strength training, believe that a high protein diet supplies extra energy, enhances athletic performance, and increases muscle mass. The misbelieve conception that large amounts of protein intake are crucial for muscle and strength improvement, is responsible for the huge proliferation of commercial supplementations which claim to transform a normal subject into a clone of Schwarzenegger. There is no evidence that excessive protein intake improves athletic

capability, and research has shown that muscle mass is determined mainly by the genetic matrix and by progressive weight training.

Compared to sedentary subjects, endurance athletes and weight training athletes may have an increased dietary requirement for protein ⁽⁵⁰⁾, but no scientific data justify the huge amounts of protein and isolated amino acids that are normally ingested by some athletes. The amount of protein intake must be related to the level of muscle destruction. Protein synthesis is stimulated by resistance exercise. Body builders and weight lifters undergoing large muscle disruptive training workouts must counterbalance protein breakdown with increased protein intake. For these athletes, the RDA are out of the normal range. Our studies (not published) with body builders estimated more than 4 g protein/kg/day during the hypertrophic phase, clearly exceeding the highest values of normal references for athletes. Protein intake must be directed related to the level of protein destruction.

PROTEINS FOR SOCCER

The amount of protein needed to satisfy requirements depends on its quality, which is determined by the relative amounts of essential amino acids present in the diet. When protein destruction is not relevant, excessive protein intake can be deleterious for performance because obliges the body, mainly the liver, to extra work for protein metabolism. Nitrogen, which is not used for protein synthesis, is mainly excreted as urea in the urine, and the rest of the amino acid molecule is used to synthesize glucose, fatty acids, or used as source of energy. Excessive protein intake increases muscle acidosis and hepatic stress.

The amount of protein required for strength and speed athletes ranges from 1.2 to 1.7 g.kg of body mass (BM)⁻¹.day⁻¹, whereas for endurance athletes the reference values range from 1.2 to 1.4 g.kg BM⁻¹.day⁻¹ (JADA, 2000). For soccer players, whose specific activity is a mixture of sprint and endurance efforts, adequate protein intake must vary between 1.2 to 1.5 g.kg BM⁻¹.day⁻¹, values easily achieved by a normal balanced diet. Branched-chain amino acid (BCAA) supplementation has been claimed to offset central fatigue provoked by long lasting exercise. Endurance exercise performance in two consecutive days is enhanced with BCAA plus arginine and citrulline supplementation. The underlying reasons are possible connected with the inhibition of cerebral serotonin synthesis and the prevention of hyperammonemia increasing urea synthesis through arginine/ citrulline ingestion ⁽²⁰⁾. It seems that power-producing capability is faster recovered after BCAA supplementation ⁽³⁴⁾. Subjects engaged on a caloric-restricted diet during strength training seem to preserve lean body mass and improve skeletal muscle performance with BCAA supplementation ⁽²⁶⁾. Other benefits were seen after BCAA supplementation. Tang and Chan ⁽⁶⁶⁾ suggest that BCAA supplementation improves substrate stores for athletes engaged in endurance exercise, reducing Purine Nucleotide Cycle metabolism, and decreasing uric acid production.

However, the results of the different studies about the benefits of BCAA supplementation are conflictual. The impaired macrophage function induced by strenuous exercise is not improved by BCAA supplementation ⁽⁷⁰⁾. It seems, also, that BCAA supplementation neither decrease some indicators of fatigue nor enhance immune function after maximal intensity exercise ⁽⁴⁴⁾. When carbohydrate is co-ingested with BCAA before and after exercise the signalling markers of muscle protein synthesis did not augment when compared to carbohydrate ingestion alone ⁽³⁰⁾.

A balanced diet whose protein intake varies within 12 to 15% of the total caloric intake provides the amount of protein necessary for muscle reconstruction in soccer.

It is important to note that research clearly stated that when glycogen depots are empty the rate of protein degradation is much higher. So, adequate muscle glycogen stores provide a significant protein-sparing effect during soccer matches and hard workouts. Some studies highlight the inadequacy of carbohydrate intake in professional soccer players ^(2, 8) what can elicit a larger incidence on protein metabolism to support energy demands.

A mixed diet is the best source of good dietary protein. Strict vegetarian athletes must plan their diet carefully to ensure an intake of enough good quality protein. The use of amino acid supplements is most likely useless and may even be harmful in the long run. Commercial protein supplements for athletes offer no advantage over natural good proteins, and they are substantially more expensive.

FAT

Fats are important sources of energy and act as insulators and protectors. They also give many foods their characteristic flavours, aromas and textures.

Fatty acids may be either saturated or unsaturated. Saturated fatty acids contain the maximum possible amount of hydrogen. Fats with a high proportion of saturated fatty acids, such as whole milk, cheese, butter, hardened margarines and meat fat, tend to be solid. Butyric acid and Palmitic acid present in butter and palm oil, respectively, are good sources of saturated fatty acids.

Mono-unsaturated fatty acids have two hydrogen atoms missing and are the most commonly occurring fatty acids. Olive oil (oleic acid) is particularly rich in mono-unsaturated fatty acids, although it is widely distributed in all fat-containing foods. Polyunsaturated fats have more hydrogen atoms missing. Corn, soya, sunflower and safflower oils, which are liquids at room temperature, are particularly good sources. Linoleic acid and α-Linolenic acid are examples of polyunsaturated fats. Fat is the richest nutritional source of energy providing 9 kcal per gram. It also acts as a carrier for the fat-soluble vitamins (A, D, E and K).

There are three fatty acids which are essential for the body's survival. Two cannot be synthesised by the body from other food stuffs so must be provided by the diet. These are the

polyunsaturated fatty acids, linoleic acid and α -linolenic acid. Arachidonic acid (precursor for eicosanoid synthesis essential for some physiological and pathological functions as inflammation and childbirth) is also very important but can be formed in the body from linoleic acid.

In developed countries, over 80 per cent of fat on an average diet comes from milk, cheese, cream, meat, butter and margarine. Average fat intake, expressed as a percentage of total energy intake, is between 35 and 40 per cent in many industrialised countries. These high values are not compatible with sport's activity and healthy living profile. To achieve optimal health, it is recommended that fat intake should be reduced to 30-35 per cent, of which no more than one-third should be saturated. Typically, total fat intake should be around 95 grams of fat per day for a man, 75 grams for a woman. However, fat intake is strictly conditioned by living and nutritional patterns. Ecological conditions are of surmount importance for nutritional behaviours. Some ethnic groups have a diet mainly vegetarian and develop excellent physical capabilities for sport. Kalenjin runners in Kenya, very successful athletes in middle and long distance running, are characterized by a diet with a reduced fat intake, about 7.5% of the total caloric intake ⁽²¹⁾, what matches fat intake of another ethnic group as the Raramuri (Tarahumara) of Mexico whose average fat intake is 9.0% of total caloric intake ⁽¹⁹⁾. More recent studies confirm this trend. Fudge et al. ⁽³¹⁾ stated that elite Kenyan runners have a diet high in carbohydrates ($67.3 \pm 7.8\%$), sufficient in protein (15.3 ± 4.0) and low in fat ($17.4 \pm 3.9\%$) while Onywera et al. ⁽⁵³⁾ assessing macronutrient intake of another group of elite Kenyan runners stated the following mean distribution: CHO – 76.5%, protein – 10.1%, fat – 13.4%. So, a low-fat diet is compatible with high sport performance.

To reduce fat intake some changes must be done as using low-fat instead of whole milk, using low-fat spread instead butter or margarine, trimming fat off meat, removing chicken skin and grilling instead of frying, can reduce fat intake substantially. In food preparation, switching to vegetable oils and eating more fatty fish favourably alters the unsaturated to saturated ratio in diet.

More important than the quantity of ingested fat is its quality. In a study of over 80.000 nurses, Harvard researchers actually found that saturated fats and trans fats are the bad fats which contribute to increase heart disease risk. Trans fats are processed by heating liquid vegetable oils in the presence of hydrogen. This process called hydrogenation permits the oils become harder at room temperature. There are different levels of hydrogenation. So, spreadable tub margarine has a hydrogenation level lower than stick margarine. Trans fatty acids are present in commercially prepared baked goods, margarines, snack foods, and processed food.

Saturated fats are mainly present in animal foods but are also present on vegetable foods as coconut, coconut oil, palm oil, palm kernel oil. Saturated fats when ingested act

as precursors of hepatic cholesterol raising total blood cholesterol more than the cholesterol rich foods present in diet. It seems that with a very functioning gut, cholesterol absorption is controlled by feedback mechanism. Saturated fats boost both good (HDL) and bad (LDL) cholesterol while trans-fat is much more dangerous because not only rises bad cholesterol but also reduces the good one. There are sufficient epidemiologic evidences that increased ingestion of saturated and trans fats contribute to the emergence of the metabolic syndrome ⁽⁴⁹⁾. However, the studies are conflictual. Lawrence ⁽⁴⁸⁾, in his review, stated that the adverse health effects usually associated with saturated fatty acids are not thoroughly supported by scientific evidence. Respecting the limits proposed by the OMS, saturated fats are essential for healthy life.

Although some nutritional behaviours can develop dyslipidaemias. Our studies (not published) with a 1st division soccer team showed, starting the season, more than half of the team displayed total cholesterol levels higher than the normal values; at the end of the season the results were similar. It seems that diet more than exercise is the main regulator for cholesterol metabolism. Although soccer players are not a group with special risk for dyslipidemia and heart disease, they need to take care about fat profile of the diet to avoid future health problems. Soccer players must recurrently control their blood lipid profile. Normal values: total cholesterol – below 200 mg/dL; LDL cholesterol – 100-129 mg/dL; HDL cholesterol – equal or above 60 mg/dL; triglycerides – below 150 mg/dL; ratio total Cholesterol: HDL – below 5.0.

The best way to decrease “bad” cholesterol (LDL-c) and to increase “good” cholesterol (HDL-c) is to alter some life habits as:

- exercise regularly even during out season periods
- cessation of smoking
- weight reduction
- decrease saturated fat intake
- maintain good body composition (reduced percent of fat)
- increase dietary fiber
- increase the amount of aerobic exercise.

Mainly in the beginning of the season, some soccer players arrive to the team with high percent of body fat and an unhealthy blood lipid profile. This situation can be easily corrected with adequate diet and increased aerobic training.

FATS AND EXERCISE

Usually, there is no problem with the quantity of fat in the athlete's diet. Too much fat in the diet, rather than too little, is the problem most athletes face, what carries several problems as insufficient intake of other macronutrients or excessive caloric intake. There is enough scientific evidence about the contribution of fatty acids to energy metabolism during exercise. It is well known that low to moderate intense exercise is mainly supported by fat metabolism. The relative contribution of fatty acids is maximal over a working intensity eliciting 65% VO_2max , decreasing the energetic importance of fatty acids when the intensity of exercise increases above that level ⁽⁶⁰⁾. It's important to highlight that the maximal fat oxidation rate during exercise varies largely among athletes. In soccer players, mean maximal fat oxidation rate is at 51.8% VO_2max , but varies between 22.9% and 88.5% VO_2max ⁽⁵⁶⁾.

It seems that there is a continuous fatty acid turnover between adipocyte and intramuscular triglycerides depots during submaximal exercise. If the goal is to reduce, by exercise, the peripheral fat accumulated as subcutaneous skinfolds, the best selected exercise must respect low to moderate intensity and must last as long as possible. However, in overweight persons, weight reduction only by exercise can be frustrated because fat oxidation is not enough to result into significant body weight decrease. Diet plus exercise is the ideal conjugation to reduce excessive body fat.

Although fatty acid metabolism plays a supporting role in energy production during endurance exercise, there is no performance benefit to increase dietary fat intake out the references (25-30% total caloric intake) preconized by the World Health Organization.

FATS FOR SOCCER

Western diets, as well as some Eastern diets, in our days, are characterized by increased intake of fats. For soccer players, increased amounts of fat intake impair the possibility to get adequate amounts of carbohydrates. Although, some authors alert against to eventual health problems produced by the reduced fat intake, the scientific data about nutritional habits of some African middle and long-distance runners point out that dietary fat is not so important for performance in sport (see above). No negative effects have been reported connected to low fat intake in elite athletes. However, essential fatty acids must be part of an athlete's diet with significant reduction of saturated fatty acid intake. Some health departments recommend the following values for fat intake: saturated fatty acids, less than 10% of the total energy intake; monounsaturated fatty acids about 15%; polyunsaturated fatty acids – 6%; linoleic acid – 1%; linolenic acid – 0,2%; trans-fatty acids less than 2%. It well known that carbohydrate metabolism *per se* is not enough to support long last-

ing exercise, like soccer. So, fat metabolism becomes energetically important; however, reference values ranging from 30 to 35% of fat intake seem be excessive. Respecting the specific intake of the essential fatty acids, values within the range 20-25% (about 1 g/kg/day) are adequate for sport.

Some fatty acids have a crucial role to health and body function and must be part of a soccer player's normal diet. For example, oleic acid is a monounsaturated fatty acid very common in Mediterranean diet which protects against heart disease and lowering the incidence of gallstone formation. High amounts of oleic acid are found in olive oil, canola oil, peanut oil, safflower oil, and chicken fat. The two essential fatty acids – linoleic acid (omega-6 or n-6) and α -linolenic acid (omega-3 or n-3), when missing or deficient in the diet, provoke cell deterioration and impairment of some body functions. Although n-3 fatty acid supplementation showed to improve anaerobic endurance capacity ⁽³⁶⁾, the balanced intake of essential fatty acids must be highlighted mainly for health reasons.

Balance between n-3 and n-6 fatty acid intake must be preoccupation for every athlete because in the last 50 years there was a dramatic shift toward omega-6 fatty acid consumption reducing the relative intake of omega-3 fatty acid. This change, coupled with the marked increase of hydrogenated and oxidized fats consumption is supposed to be one of the leading factors which contribute to chronic illnesses, especially cardiovascular diseases. So, it urges to increase n-3 oils intake reducing the ratio n-6: n-3. The n-3 fatty acid recommendation to achieve nutritional adequacy is 0.6-1.2% of the total energy intake for α -linolenic acid and about 500 mg of EPA and DHA ⁽³³⁾. EPA (eicosapentaenoic acid) and DHA (docosahexaenoic acid) nutritional needs for a soccer player are easily achieved consuming 2 fatty fish meals per week (herring, sardine, salmon, etc.). Some studies have shown the benefits of the n-3 fatty acids, mainly related with the prevention of heart disease. They can be summarized ⁽³³⁾:

- prevention of arrhythmias (ventricular tachycardia and fibrillation)
- formation of and competition against various prostaglandins and leukotrienes
- anti-inflammatory properties (partly by prostaglandin effect)
- antithrombotic effect
- hypolipidemic effect on triglycerides and VLDLs
- inhibition of atherosclerosis.

Although soccer players, by their age, are less prone to heart diseases, some problems can be avoided with a n-3 rich diet. When soccer player's diet lack fatty fish and green vegetables is very difficult to achieve a balanced n-6: n-3 ratio. In those cases, n-3 fatty acids supplementation is justified. Flaxseed oil or concentrated fish oil supplements can be optimal sources to regain the nutritional equilibrium between omega-6 and omega-3 fatty acids.

CARBOHYDRATES

Carbohydrates are the most readily available energy source for human work as well as for sport. Optimum diet for most sports must get about 60-70% of the total energy as carbohydrates, higher than the RDA (recommended dietary allowances) for sedentary people – 50 to 55%.

There are two main types of carbohydrates – sugars and starches. The most common food sugars are sucrose, glucose and fructose, which occur in fruit, vegetables and honey, and lactose which occurs in milk. Ordinary sugar, often referred to as table sugar, is sucrose, produced from sugar cane and sugar beet. Starches occur mainly in cereals and potatoes.

All carbohydrates, whether sugars or starches (both provide 4 kcal/g), are compounds made from one or more basic building blocks. Almost all of these blocks are ultimately transformed into glucose. Starches or “complex carbohydrates” must be broken down into glucose before they can be used by the body.

Which foods are more fattening? Fats or carbohydrate? The answer is not important in sport context because athletes need larger amounts of energy than sedentary people to cope with energetic demands of exercise. Soccer players must get a balanced diet to avoid that extra energy be stored as body fat. Overweight can occur, even in athletes, when diet is not balanced and extra energy intake is accumulated as fat.

CARBOHYDRATES AND EXERCISE

Carbohydrates are essential for physical activity. High intensity exercise is mainly supported by carbohydrates. These nutrients are stored in the body (muscles and liver), in the form of glycogen. They are easily converted into energy at a rate which is directly proportional to exercise intensity.

Fats and proteins are exclusively metabolized through aerobic pathway while carbohydrate can be metabolized by both aerobic and anaerobic pathways.

During soccer match, although aerobic metabolism is basic for energy providing, the high intensity skills and displacements are mainly supported by anaerobic metabolism. Plain anaerobic metabolism can only be used as short-term measure as it produces lactic acid which is rapidly dissociated into lactate and hydrogen ion; the latter increases acidosis which impairs muscle efficiency, producing an early onset of fatigue ⁽¹⁷⁾.

Soccer performance can be improved by:

Increasing muscle strength and body's efficiency. Endurance athletes (soccer is a special kind of endurance activity) can therefore maximise the use of fat at the start of an event. This spares muscle glycogen depots for the anaerobic bursts of high intensity exertions.

Eating large amounts of carbohydrates before training or competition.

Research has shown that carbohydrate supplementation can improve performance when compared to subjects on a mixed diet. Carbohydrate loading increases soccer players' capability for high intensity exercising. Carbohydrate ingestion before and during the match increases soccer performance towards the end of the game ⁽³⁾.

At low intensity exercise the body works aerobically. At less than 50% VO_2max , fat is the predominant fuel, accounting for more than half of total energy production ⁽¹⁵⁾. The energy stored as fat cannot be released rapidly enough for exercising more intensely than about 50% VO_2max . At about 60-65% VO_2max , the contributions of fat and carbohydrate are roughly equal. Above this level of exercise intensity carbohydrate is the major fuel and maximum availability is essential ⁽⁶¹⁾.

In very short intense exercise, for example the 100m sprint or a burst of maximal running in soccer, nearly all the energy is supplied anaerobically. Anaerobic metabolism uses up carbohydrate very quickly, increases muscle acidosis and reduces muscle contractility.

Carbohydrate (CHO) and fat are both metabolized during soccer's training and matches. The relative contribution of these two substrates depends on factors such as: exercise intensity, exercise duration, and player's physical training condition ⁽¹⁴⁾. Brooks and Mercier ⁽¹⁵⁾ stated that endurance training reduces CHO utilization and increases fatty acid oxidation during mild or moderate exercise intensity (> 45 and 50-55% VO_2max) while this not true for exercise intensities higher than 65% VO_2max . At such exercise intensity, even trained persons change the energy supply to CHO metabolism. This is the basement for the “crossover concept”. However, Coggan ⁽²³⁾, conflicting with the crossover concept, stated that training can change the relative contributions of CHO and fat even during exercise performed at 65-85% VO_2max . At this exercise intensity fat can still be a significant source of energy, which opens a new/ old field for endurance training in soccer.

CARBOHYDRATES FOR SOCCER

The profile of carbohydrate intake must be determined by the profile of training and competition as well as the motor pattern of the players during the match. Highly physical active people must increase their CHO consumption. Heavy exercise like soccer imposes greater demands on the body's CHO stores and fatigue is often associated with muscle glycogen depletion. Consuming foods which contain mainly CHO not only provides substrate to refuel muscle and hepatic stores of glycogen, but also help decrease fat intake. High intakes of fat can be deleterious for soccer performance. Kenyan success on the international athletic scene in middle and long distance running is well known and among the factors that contribute to success there is the typical diet of Kenyan athletes. Kenyan runners ingest a nearly complete vegetarian diet with adequate amounts of protein, reduced fat intake and high CHO intake. Although the biomechanical and physiological features are different between run-

ners and soccer players, high carbohydrate intake is of extreme importance for both athletes. Endurance training can improve fat acid oxidation at high exercise intensities (65–85% $\text{VO}_{2\text{max}}$) which can be an effective support for energy supply during soccer's training and competition. Soccer diet must be matched with energy demands for training and competition in order to avoid energy failure which is always deleterious for performance.

STRATEGIES TO INCREASE CARBOHYDRATE CONSUMPTION

Carbohydrates are crucial for hard training and competition in soccer. Some strategies can be established in order to increase CHO intake. Hard training and competition for long lasting sports requires about 70% of the total caloric intake as carbohydrate. So, diet must respond to this nutritional demand:

Increase carbohydrate-rich foods intake (wholegrain breads and breakfast cereals, rice, oatmeal, maize meal and other grain foods, pasta and noodles, fruits, starchy vegetables (e.g. potatoes, corn, sweet potato, yams), legumes (lentils, beans, soy-based products), sweetened dairy products (e.g., fruit-flavoured yoghurt, milk-shakes).

Pay attention about fat content of some foods. Many foods presumably high CHO are, in reality, high-fat foods.

Sugar and sugary foods provide a compact CHO source. They may be particularly useful in a high-energy diet, or, because their easily digestibility, when carbohydrate are needed before, during and after exercise.

CHO drinks (e.g., fruit juices, soft drinks, fruit/ milk smoothies) are good components for a high-carbohydrate diets.

If the athlete needs more energy, he must increase the number of meals and snacks rather the size of the meals to avoid larger release of pancreatic insulin.

Avoid foods with high fiber content when energy needs are high, or during pre-exercise meals (increased gastric work can impair performance if exercise starts during digestion).

To speed the recovery process, mainly the replenishment of glucose stores, a high-carbohydrate (foods with high glycaemic index) meal must be eaten within 30 minutes after a prolonged session of exercise.

During prolonged exercise (training or competition) the athlete must consume CHO (e.g., sports drinks) in order to supply energy and water (water supply can be crucial because in some situations water losses by sweating can be in order of 1–2 L per hour).

It is important to discriminate between meals for massive energy supply and meals for a healthy diet. The nutritional procedures cited above are adequate for an athlete exercising daily on a high intensity long lasting training regime; however, when the same athlete is out

of the intensive training phases, his diet must be more balanced in relation to macronutrient intake. It's important to highlight that some of the deleterious outcome attributed to fat ingestion, namely saturated fatty acids, is derived from massive intake of simple sugars as fructose whose oxidation rate is several times higher than glucose ⁽⁴⁸⁾.

DIETARY FIBER

Dietary fibers are a mixture of mainly indigestible substances which are found in the walls of plant cells and in plant stems. They are non-digestible polysaccharides known as microbiota-accessible carbohydrates which are the main source of energy for gut bacteria. So, dietary fibers modulate gut microbial composition and function with direct action over health and immunity ⁽²⁴⁾.

In the digestive system, dietary fibers affect the way the body absorbs and uses nutrients. It helps the excretion of bile acids, and soaks up water, thereby bulking up the faecal mass.

Dietary fibers among other substances are mainly made up by soluble (pectins, gums, and mucilages) and insoluble (cellulose, hemicellulose, and non-carbohydrate lignin molecules) materials. The suggested health benefits of dietary fiber include a decrease in constipation and diverticular disease, prevention of bowel cancer, lowering of the blood cholesterol level and better control of blood glucose levels after consumption of carbohydrates and, through the specific action of the glucomannans, increasing immune surveillance ⁽⁶⁷⁾.

A diet which provides around 25–30 grams of dietary fiber is well adequate to a good digestive function.

In our studies about the nutritional status of several male and female athletes, soccer players included, a large percentage didn't meet the daily recommendations for dietary fiber, incurring into a potential situation of health and immune fragility.

Consumption of fiber in developed countries is about half the recommended level and should be increased. Good sources of dietary fiber are fruits, nuts, cereals, vegetables (especially peas and beans) and wholemeal bread. For getting the adequate amount of dietary fiber is enough to ingest the following foods, or equivalents: 1 slice of whole bread, 1 banana, 1 apple, 2 servings of beans, 1 serving of rice, 1 orange and 1 raw carrot.

It is important to note that some individuals tend to overpass fiber deficits of the diet with large intake of bran supplements. This behaviour is not correct because huge intakes of dietary fiber could hamper mineral absorption conducting to mineral deficiency. Fatigue, mood disturbances, and gastrointestinal stress are common among athletes during training and competition. To avoid these constraints a balance diet is crucial. The gut microbiome, important for several body functions, changes its composition under physical and emotional stress. Diet is known to modulate the composition of gut microbiota which acts like an endocrine organ secreting serotonin, dopamine and other neurotransmitters

⁽²²⁾. Then, increased intake of plant polysaccharides must be recommended for elite athletes, targeting the increasing of microbiota and avoiding the gastrointestinal constraints induced by physical and psychological stress.

VITAMINS

Vitamins are micronutrients which help to control vital body processes. They are found in small amounts in food and because the body cannot make most of them, they must be got through the diet. An insufficient supply can cause poor health either by increasing susceptibility to illness or by developing a specific vitamin deficiency disease. Children's growth can be impaired by vitamin deficiency.

Although a growing number of individuals take vitamin supplements, this is rarely necessary. Biesalski and Tins ⁽⁹⁾ reviewed the effects of supplementation combining the intake of seven vitamins and three minerals and stated that none study found differences between treatment and placebo. Almost all studies point out to the definition of rationale criteria for micronutrient supplementation avoiding deficiency or excessive intake. Logically, the criteria for determining nutrient intake requirements should be based on the actual nutritional status not the performance potentiation.

Vitamins display several biological roles. We summarize the more important ⁽²⁷⁾:

Release of energy – Thiamine (B1), riboflavin (B2), Niacin (B3), Pantothenic acid (B5), Biotin (B7)

Blood function – Pyridoxine (B6), Cyanocobalamin (B12), Folic Acid (B9)

Healthy skin and eyes – Vitamin A (retinol)

Antioxidant action – Vitamin C (ascorbic acid), Vitamin E (tocopherol)

Bone metabolism – Vitamin D (calciferol), Vitamin K (phyllloquinone and menaquinone).

VITAMINS FOR SOCCER

Although the ergogenic benefits of vitamins and mineral supplements are very doubtful when diet is energetically balanced, diversified, and omnivore, soccer players as other athletes tend to speculate about the causes of their natural impairment when training is hard. For some athletes, when fatigue occurs, it is not accepted as the natural physiological response to training, but as a body deficiency which can be overcome by supplementation. Only assessing the athlete's nutritional status eventual deficit in some nutrient can be found.

Meta-analysis of the literature, as well our own findings with Portuguese soccer players, show that soccer players are usually in a balanced diet without vitamin or mineral deficits. In these conditions, any supplementation can be deleterious for health and performance. However, Brannstrom et al. ⁽¹²⁾ showed that the level of vitamin D is markedly low in adolescent female soccer players. In that cases supplementation seems advisable. When exposure to sunlight is limited it is important to measure the levels of 25(OH)D3, calcium, and parathormone. Kopec et al. ⁽⁴⁵⁾ stated that 50% of a group of Polish professional soccer players showed normal values of 25(OH)D3 after summer but only 16.7% after winter. As vitamin D metabolism is related to bone and muscle functions, supplementation can prevent physical performance impairment and health constraints.

Exercise increases oxidative stress (i.e., free radicals' formation is increased during exercise). Free radicals are highly reactive chemical species with one or more unpaired electrons in their outer orbit. Free radicals will readily react with cellular molecules, resulting in several damages. Cell damages are counterbalanced by antioxidant defences, some of them constituted by antioxidant enzymes other built by several vitamins (beta-carotene, vitamin C, vitamin E). Oxidative stress is aggravated by air pollution, smoking, alcohol, physical and emotional stress and poor diet. Free radicals are also a by-product of weight loss. Then, it is acceptable that during periods when athletes experiment energy restriction for body weight reduction purposes, they are not able to achieve an adequate vitamin status. In that case, to avoid vitamin deficiency, athletes must increase vitamin intake not through supplementation but ingesting more antioxidants rich-foods. The body naturally synthesizes its own antioxidants (e.g., superoxide dismutase, catalase), but, under extreme stress, the natural defences of the body may be insufficient to protect against the deleterious effects of free radicals. Antioxidants rich-foods can be the solution: fruits and berries, vegetables, legumes, nuts and grains, cocoa and coffee, tea, and spices are good sources of antioxidants.

The effects of vitamin supplementation in physical performance when nutrition is adequate are hard to verify. In moderately trained individuals without vitamin deficiency, the administration of vitamins C and E has no effect on endurance performance ⁽⁷¹⁾. Moreover, some studies point out for the deleterious effects of vitamin supplementation. Paulsen et al. ⁽⁵⁵⁾ verified that vitamin C and E supplementation hampers the cellular adaptation to endurance training decreasing the markers of mitochondrial biogenesis. Reinforcing this study, it was stated that vitamin C supplementation hampered endurance capability reducing the exercise-induced expression of key transcription factors involved in mitochondrial biogenesis ⁽³⁵⁾. Lamprecht et al. ⁽⁴⁷⁾ showed also that the administration of an encapsulated antioxidant concentrate (107 IU vitamin E, 450 mg vitamin C, 36 mg beta-carotene, 100 g selenium) in trained male cyclists increased lipid peroxidation and decreased plasma glutathione peroxidase after strenuous exercise. In his meta-analysis Stepanyan et al. ⁽⁶⁴⁾ stated that tocoph-

erol supplementation did not result in significant protection against either exercise-induced lipid peroxidation or muscle damage. Usually supplementation is ineffective. When vitamin supplementation is introduced in the athletes' nutritional habits one must be aware about the eventual reverse effects induced by large doses of vitamin intake.

MINERALS

The fundamental nature of life is chemical; the way for life starts with the inorganic elements ⁽⁶⁹⁾. All forms of living matter require many inorganic elements for their normal life processes. Virtually all the elements of the periodic table have been found in living cells, though not all are necessarily essential to life. The nutrients which are commonly referred to as mineral elements or inorganic nutrients, and which have definite demonstrable functions in the human body and metabolism are calcium, phosphorus, magnesium, sodium, potassium, sulphur, chlorine, iron, copper, cobalt, iodine, manganese, zinc, molybdenum, selenium, and perhaps fluorine. Aluminium, arsenic, nickel, and silicon might be mentioned as elements which occur very consistently in the human body, though for them doesn't exist any proof of human need. The minerals are interrelated and balanced against each other in human physiology. For example, calcium and phosphorus are in a close relationship in the formation of bones and teeth. Iron, copper, and cobalt (in vitamin B12) are interrelated in haemoglobin synthesis and red blood cell formation. Sodium, potassium, calcium, phosphorus, and chlorine serve several purposes in the body fluids. Calcium and magnesium are necessary for normal soft tissue and nerve cell function. Iodine is an essential constituent of the thyroid hormone, while zinc, molybdenum, and manganese serve as essential activators of a number of enzyme-catalysed metabolic reactions ⁽⁶⁹⁾.

About 13 minerals are known to be needed by the body, and all must be derived from the diet. All minerals are essential for athletes but the three elements whose supply is most likely to be critical are calcium, iron, and iodine.

MINERALS FOR SOCCER

In well fed athletes, mineral deficiencies are very rare. Most of the essential minerals for good body function (e.g., calcium, phosphorous, magnesium, potassium, iron, sodium, etc.) are needed in minute amounts and are so widely distributed in different foods that it is unlikely that anyone who eats a varied and balanced diet would be deficient in any of them. However, when some mineral deficiency is verified it must be rapidly corrected because mineral deficits impair both aerobic and anaerobic metabolism. Even, in situations of adequate daily intake of iron, increased serum levels of hepcidin were observed in female

long-distance runners what can increase the risk of iron deficiency ⁽³⁹⁾. In several sportsmen, including football players, it was shown that overtrained athletes are characterized by lower levels of zinc, ferritin and IGF-binding protein ⁽¹⁾. These features can alter blood rheology affecting lactate metabolism and overall physical performance ⁽⁴²⁾.

It is important to note that some athletes usually tend to cope with fatigue by introducing several supplementations in their diet. This can be harmful for health. In normal fed soccer players, the ergogenic benefits of micronutrient supplements are very doubtful and can result into body dysfunctions. So, it is necessary to analyze the athlete's nutritional status. The best way consists in assessing several indicators of micronutrient status by the determination of daily dietary intakes or chemical profiles of the blood. After that, any dietary inadequacy should be corrected by supplementation or increased intake of natural foods rich in the micronutrient in deficit.

The literature and our own studies ^(37, 59) showed that soccer players and other ball-game players have an overall mineral intake within the recommendations. Usually, calcium, magnesium, iron, zinc, sodium, phosphorus, potassium and selenium intakes are sufficient and higher than the RDA, in male players.

Although, male soccer players have usually an adequate mineral intake they need continuous nutritional education to avoid negative nutritional behaviours which can impair their capability to perform in the best conditions.

SOCCER'S PLAYER NUTRITIONAL PROFILE

Soccer embraces a diversity of skills which only can be realized at top level by intensive and systematic training. The primary requirement for the athlete in training is to consume a well-balanced, nutritionally complete diet, which will meet the additional nutrient demand imposed by training load. The nutritional strategies must regard the individual differences in the physical demands imposed to the players during matches and training. These differences are related to the training status, but also to the specific playing position and tactical exigencies.

Energy intake must be sufficient to maintain body mass, except when players are overweight and weight reduction becomes necessary. The athlete's diet should be low in fat and high in carbohydrate (mainly complex carbohydrates), with an adequate amount of protein. Soccer players must reduce fat intake, mainly saturated and hydrogenated fats, increasing the dietary amount of omega-3 fatty acids and avoiding alcohol intake.

FREQUENCY OF EATING IN SOCCER

Soccer players undergoing strenuous training often have energy expenditures 2-3 times greater than untrained individuals. High-intensity training elicits greatly increased carbohydrate needs. Carbohydrate is required to optimise fuel availability during training sessions and to restore muscle glycogen depots after exercise. High energy intake corresponds to a high frequency of food consumption. Some athletes report 8-10 eating occasions per day. Between-meals snacks have been found to account for approximately one third of the energy intake in a variety of athletic populations.

It appears that eating a series of small meals and snacks over the course of a day is a practical strategy which allows increased energy intake while reducing the gastric discomfort of infrequent large meals.

PRIOR TO EXERCISE

Nutritional errors before competition can impair performance while correct nutritional practices are difficult to enhance competitive performance. The purpose of the pre-competition meal is to maximize glycogen stores, mainly in skeletal muscles, and provide adequate hydration while avoiding gastric disorders. The literature, regarding to the pre-competition meals, pay attention to the following factors: (i) timing; (ii) amount; (iii) components of the meal: and (iv) fluids.

The main goals for pre-competition meal can be summarized as follows:

- to optimise muscle and liver glycogen stores

- to avoid excessive fullness and gastrointestinal discomfort (e.g. vomiting or diarrhoea) during exercise

- to attenuate the exaggerated increase in insulin levels normally experienced after a high carbohydrate meal.

It is important to realize that, before competition, any drastic change from the normal dietary habits can result in gastric distress, eventual diarrhoea and presumable physical impairment. So, the nutritional claims that put emphasis on a high carbohydrate diet before competition must be challenged to the athlete's normal diet. If the athlete is normally habituated to eat a high fat/protein diet, drastic change for a high carbohydrate diet can be deleterious for performance. Diet must be also a matter for training.

A carbohydrate-rich meal consumed 4 hours before exercise significantly increases muscle and liver glycogen content, mainly if the glycogen depots of those tissues are reduced. One study has shown that the intake of 300g of carbohydrates 4 hours before exer-

cise increased performance when compared to subjects who did not ingest carbohydrates. However, soccer players in a fed state didn't increase running and sprint performance during the match after consumption of 250 ml of 12% carbohydrate solution ⁽³²⁾. This highlights the importance of an adequate nutrition before competition. Eventual energy deficits in the days preceding the matches can be overreached through a larger breakfast in the match day, apparently without abdominal discomfort ⁽¹³⁾.

One single study reported earlier fatigue during exercise after consuming carbohydrates 30 minutes before exercise. This result was so widely reported that warnings to avoid carbohydrates in the hour prior to exercise have become part of sports nutrition dogma. However other studies didn't replicate that finding. Other investigations have found either no detrimental effect or even improvements in performance ranging from 7 to 20%. The problem concerns to the individual responsiveness to insulin increase after ingesting a carbohydrate-rich meal. Reduced blood glucose can impair physical performance in hypersensitive athletes while others can perform well in those conditions.

For individuals that experience exaggerated insulin levels when they eat CHO prior to exercise, the symptoms of eventual physical failure may be overcome by manipulating the timing of carbohydrate feedings or by choosing fructose and foods with low glycaemic index. However, for the majority of these individuals, any adverse metabolic effects are transient, of little physiological significance and are offset by the increased carbohydrate availability.

In general, 3-4 hours are sufficient for the digestion of a large carbohydrate-rich low fat meal. On the other hand, a smaller carbohydrate-rich snack or drink may be consumed without gastric problems 30-90 minutes prior to exercise. This may be a more suitable choice for athletes who are competing in early morning events, or when there is a long gap between the last meal and the exercise session.

DURING EXERCISE

There is strong evidence that increasing the availability of carbohydrates during prolonged exercise can improve physical work capability. While in some sports (e.g., cycling) nutritional choices can include some solid foods, during soccer match and rest time, the supply of carbohydrates can be done by sports drinks much easier to digest and absorb. To avoid gastrointestinal discomfort sports drinks utilization must be "trained" during workouts. During very hot and humid days, fluid replacement must begin as soon as possible during the match utilizing any stops to drink small quantities of fluid.

The optimum formulation for rehydration purposes during exercise will vary, depending on the relative requirements for fluid replacement and substrate provision, which in turn will depend on the environmental conditions and exercise intensity. When water replace-

ment is the first priority water (as is demonstrated by the study from Hill et al. ^[38]) or an isotonic solution containing glucose and sodium will be most effective.

It must be noted that only when some degree of dehydration occurs – perhaps as much as 1.5-2% of body weight – the stimulus for drinking begins. So, drinking behaviour prior to dehydration must be stimulated and trained during workouts. As beverage palatability stimulates drinking the athlete must choose the commercial or auto-constructed beverage he likes more changing the drink flavour every moment he begins to get nauseated what often occurs with many athletes. In those moments water is the best solution for hydration.

One good solution for energy and water replacement during the match can consist in ingesting a carbohydrate gel, along with water, what have been confirmed to improve performance during prolonged intermittent efforts.

POST-EXERCISE

As long as the timing and total amount of carbohydrate consumed is sufficient, muscle glycogen restoration is unaffected by the frequency of food intake. One study showed that muscle glycogen replenishment after 24 hours was similar when 525 g of carbohydrate were provided as either 2 or 7 meals. In another study, muscle glycogen depots were similar when 10 g carbohydrate/kg body mass were consumed as either 4 large meals or 16 snacks. However, it is important not to delay the beginning of carbohydrate ingestion. The first two hours after exercise are crucial for optimal muscle glycogen replenishment. The rate of glycogen restoration is increased in the first two hours, so, carbohydrate ingestion must start quickly after exertion.

While during normal meals low glycaemic index carbohydrates are the most important for any subjects' diet, post-exercise meal must be characterized by the ingestion of large amounts of moderate to high glycaemic index carbohydrates. Recovery meals must put emphasis in the ingestion of carbohydrates with moderate and high glycaemic index. Low glycaemic index carbohydrates should not make up more than one third of recovery meals.

After strenuous exercise, many athletes report diminished appetite. This symptom can last till 2 or 3 hours. Such persons may require nutritional counselling. They should try a variety of foods and drinks to see which suits them best. As solid or liquid carbohydrates have the same replenishing action, fluids can be a suitable choice to start early glycogen restoration.

After exercise, not only glycogen restoration is important. Replacement of water and electrolytes are crucial for maintenance of exercise capacity when repeated bouts of exercise have to be performed. When soccer players practice twice a day, recovery (mainly hydro-electrolyte balance) between workouts is crucial for maintaining work capacity. A dehydrated body is not capable to exercise intensively.

NUTRITIONAL CARES

FOR YOUNG SOCCER PLAYERS

Young soccer players necessitate appropriate caloric and selected nutrient intake, with special emphasis on carbohydrates, protein, and certain minerals (iron, calcium, and sodium), to meet the energy and nutrient demands for normal growth and development, as well as the additional nutritional requirements imposed by training and competition.

In addition to proper nutrition, sufficient rest and adequate recovery among training sessions are all equally fundamental for athlete's health, development, and adaptation to training. Proper nutrition and appropriate recovery time between bouts are essential to avoid health risks for young soccer players. Excessive strain from practice in hot environments must be avoided in young players putting special emphasis in adequate hydration the main factor for efficient thermoregulation. Proper hydration is the crucial completion for a good nutritional status for every athlete, particularly for younger. The general dietary needs of the young soccer players appear to be similar to those of the adult soccer players ⁽⁶⁾. However, there are several age – and maturation-related differences. Compared with adults, children's daily recommended protein intake (per kg of body mass) is higher; during exercise children's muscles rely more on fat than on carbohydrates ⁽⁶⁾, however, we think that young soccer players don't need higher daily fat intake. The main care the coaches need to observe in relation to children's training is to avoid excessive hot strain, increasing fluid intake which must have good palatability and easy stomach emptying.

As for everybody a healthy diet for children is one that provides the energy they need from the correct proportions of nutrients. They should have a wide variety of different foods and drinks to ensure they obtain all the vitamins and minerals they need.

ALCOHOL IN SOCCER

Alcohol is "genetic" incompatible with high physical performance. Even when some nutritionists claim about the health benefits from red wine intake it must be assumed that soccer players, like other young sportsmen, are prone to excessive drinking, so the better is to completely avoid alcoholic beverages. Studies show that ethanol consumption is associated with spermatogenesis damage ⁽²⁵⁾ and testosterone level alterations ⁽⁶³⁾.

Chronic intake of large amounts of alcohol is associated with the development of a fatty liver and, ultimately, irreversible cirrhosis of the liver. Liver is a crucial organ for sport. A marathoner's aphorism claims that the runner "run more with his liver than with their legs". In marathoners, hypertension is related with high frequency of alcohol consumption ⁽⁴²⁾ increasing the risk for cardiovascular events.

Alcohol as a food contributes with 7 kcal/g of ethanol oxidized, compared with carbohydrates and proteins (4 kcal/g) or fats (9 kcal/g). In Portugal, some studies we have done estimated that alcohol contributes 5 to 10 percent of the total caloric intake of the population. Athletes have the same nutritional pattern. Of alcoholic drinks most frequently consumed by athletes are beer and wine.

Although some fermented (not distilled) alcoholic beverages can be source of some important nutrients (pantothenic acid, folic acid, other B complex vitamins and phosphorus), this is not reason enough to admit them in the athlete's diet because alcohol affects both glycogen recovery ⁽¹⁶⁾ and protein synthesis ⁽⁵⁴⁾ after exercise.

Alcohol is absorbed, primarily in the stomach, by simple diffusive transport without need for prior digestion. Approximately 10-20% of ethanol is absorbed in the stomach and 75-80% is absorbed in the small intestine. Absorption of alcohol is more rapid from solutions containing 10 to 20 percent on an empty stomach; alcohol in the presence of large quantities of food is absorbed less rapidly.

The absorbed alcohol diffuses rapidly and uniformly throughout the fluid compartments of the body and it is found in different tissues in proportion to their water content. In the central nervous system ethanol acts as a depressant, first in the phylogenetically more recent regions of the brain (affecting at first such parameters as acquired, learned inhibitions, judgment, personal behaviour, and memory). Animal studies point out for the progressive deterioration of cognitive functions induced by alcoholic behaviour ⁽⁴⁶⁾. The progressive effects of alcohol roughly parallel its concentration in the blood.

Physical performance is impaired by alcohol which acts on the absorption of nutrients affecting intestinal mucosa and cellular transport processes. It inhibits active cellular transport of sodium and potassium, amino acids, and glucose. At a 2 percent concentration in the lumen of the small intestine (a concentration that can be present in moderate drinkers) the absorption of a number of essential amino acids is inhibited. Alcohol ingestion may lead also to decreased absorption of fat and vitamin B12. At the same time ethanol stimulates triglyceride synthesis by the intestine with subsequent increased lymph triglyceride content.

By all these reasons there is no rationality to accept alcohol drinking in athletes; not only because alcohol directly affect health and performance but also because alcohol ingestion is often related with malnutrition affecting the recovery process after training and competitions. Alcohol drinking is not compatible with the philosophy of life inherent to sportsmen and sportswomen who care to get in sport the realization of their more elevated goals.

NUTRITIONAL ERGOGENICS FOR SOCCER

Success in soccer, contrary to other sports, is not dependent upon genetic endowment. High commitment to training and strong will and personality can overpass eventual physical deficits. In soccer, tactical intelligence is the main factor to get success. Physical conditioning must be the result of tactical goals which put the emphasis in the continuous greater level of game's organization. Then, all motor capabilities are conditioned by the level of tactical demands. Nutrition must respond to these demands.

However, soccer players often believe that training maximization turn necessary the recurrence to ergogenic aids. Soccer is extremely permeable to faddism in matter of ergogenic aids. However, scientific research rarely supports supplementation benefits. While repeated sprint ability, crucial for soccer higher performance, is not enhanced by caffeine, arginine and branched-chain amino acids ⁽²⁸⁾, it seems that soccer performance is increased by ingestion of nitrate-rich beetroot juice ⁽⁵²⁾. However, any nitrate-rich food (e.g., arugula lettuce, spinach, celery and iceberg lettuce), are able to increase nitrate intake and boost body pool of nitric oxide which is known to have direct effect on circulatory function. We must refuse the miraculous effect of a single food; nutrition is a construct implying all the foods of the vegetal and animal nutritional spectrum. Another study ⁽³⁶⁾ showed the positive effects of n-3 fatty acid supplementation on anaerobic endurance capacity in soccer players. Instead of supplementation, soccer players must increase n-3 rich-foods intake whose benefits overpass the specific action of the n-3 fatty acids in the body.

The misconception that protein was the principal component of the human muscle was spread in sport. Actually, it is water. It was claimed that athletes have special protein needs what justified protein supplementation. Muscle building is not caused by eating extra protein but by intensive muscular work. Once basic protein needs have been met, the small additional amount needed during intense strength training is easily obtained from a balanced diet. So, the false and misleading claims which promote the massive ingestion of protein by athletes result from advertising campaigns taken by the industry of supplements looking for economic profits. Ingesting massive amounts of amino acids does not cause humans to release growth hormone with the subsequent increasing on muscle mass and muscle performance ^(44, 70).

Athletes who eat a balanced diet rarely need extra protein or vitamins. For example, one kilogram of muscle contains only about 200 grams of protein, since it is composed of more than 72 percent water. So, if an athlete is gaining one half kg of muscle every week through an excellent strength training program, he is adding only about 100 grams of protein each week, or about 15 grams of protein each day. Two cups of corn and beans will meet this need. Requirements for only four vitamins increase with exercise: thiamine, niacin, riboflavin, and pantothenic acid. These vitamins are used up minimally in the breakdown of

carbohydrates and, to a small degree, using protein for energy. They are abundant in food. Furthermore, deficiencies of these vitamins have never been reported in athletes.

Athletes believe that supplements can help them because their beliefs result from the coincidence promoted by the natural improvement induced by training and the effect placebo. Generally, supplementation is a waste of money and when uncontrolled can produce adverse effects for health and performance.

Some products are widely advertised as ergogenic without scientific support: wheat germ oil and bee pollen (falsely claimed to boost energy and endurance), ginseng, spirulina, royal jelly, ma huang, guarana, wheat grass, gotu kola, cayenne pepper and so on. The misconception that natural products are safer than artificial ones can be very dangerous. In South Korea, extracts of ginseng roots are claimed as good for almost every disease. Logically this is a fraud and every person must put some doubt about the miraculous properties of this natural product.

It's necessary to distinguish between the inexistence of scientific proofs about the ergogenic efficacy of a product and the placebo effect which promotes the faith about the benefits of that product. Sometimes coaches and nutritionists have to accept the supplementation because the psychological behaviour of the athlete is conditioned by his beliefs.

Although most purported nutritional ergogenic aids have not been shown to enhance physical performance in well-trained, well-nourished athletes, some reliable scientific data support some ergogenic efficacy of several substances, including caffeine, creatine and sodium bicarbonate, but for soccer the potential benefits need to be evaluated for each person because we have reported some gastric discomfort after supplementation with these products in several athletes. Some energy drinks, claimed to boost energy, actually induce lipid peroxidation and oxidative damage⁽⁵⁷⁾. Training for performance in sport is a matter of adaptation-loss of adaptation cycles. This alternation must be respected and not thwarted by ergogenic supplementation.

CONCLUSION

Nutrition in soccer must be part of nutrition for life. Meanwhile some specific nutritional guidance can be underlined for soccer players. Consumption of carbohydrate-rich foods for energy needs and glycogen resynthesis are key behaviours soccer players need to focus on a daily basis. To attain this goal there is no need of qualified nutritional counselling because everybody has sufficient knowledge about this subject. The problem is to adapt daily strategies towards improving food selections. This can be done by players, their parents, coaches and physical trainers. Nutrition education is a task to develop every day during the individual's lifespan.

REFERENCES

1. Aïssa Benhaddad A, Bouix D, Khaled S, Micallef JP, Mercier J, Bringer J, Brun JF (1999). Early hemorheologic aspects of overtraining in elite athletes. *Clin Hemorheol Microcirc* 20(2): 117-125.
2. Anderson L, Orme P, Naughton RJ, Close GL, Milsom J, Rydings D, O'Boyle A, Di Michele R, Louis J, Hambley C, Speakman JR, Morgans R, Drust B, Morton JP (2017). Energy intake and expenditure of professional soccer players of the English Premier League: evidence of carbohydrate periodization. *Int J Sport Nutr Exerc Metab* 4: 1-25. doi:10.1123/ijsnem.2016-0259
3. Baker LB, Rollo I, Stein KW, Jeukendrup AE (2015). Acute effects of carbohydrate supplementation on intermittent sports performance. *Nutrients* 7(7): 5733-5763. doi:10.3390/nu7075249
4. Bangsbo J (1994) Energy demands in competitive soccer. *J Sports Sci* 12(Spec No): S5-S12.
5. Bangsbo J, Mohr M, Krstrup P (2006) Physical and metabolic demands of training and match-play in the elite football player. *Journal of Sports Science*, 24(7):665-674.
6. Bar-Or O, Unnithan VB (1994). Nutritional requirements of young soccer players. *J Sports Sci* 12: S39-S42.
7. Barnett C, Carey M, Proietto J, Cerin E, Febbraio MA, Jenkins D (2004). Muscle metabolism during sprint exercise in man: influence of sprint training. *J Sci Med Sport* 7(3): 314-322.
8. Bettonviel AEQ, Brinkmans NYJ, Russcher K, Wardenaar FC, Witard OC (2016). Nutritional status and daytime pattern of protein intake on match, post-match, rest and training days in senior professional and youth soccer players. *Int J Sport Nutr Exerc Metab* 26(3): 285-293. doi:10.1123/ijsnem.2015-0218
9. Biesalski HK, Tinz J (2017). Multivitamin/mineral supplements: rationale and safety. *Nutrition* 36: 60-66. doi:10.1016/j.nut.2016.06.003
10. Bloomfield J, Polma R, O'Donoghue P (2007) Physical demands of different positions in FA Premier League soccer. *J Sports Sci Med* 6: 63-70.
11. Bormuth V, Varga V, Howard J, Schaffer E (2009). Protein friction limits diffusive and directed movements of kinesin motors on microtubules. *Science* 325(5942): 870-873.
12. Brannstrom A, YU JG, Jonsson P, Akerfeldt T, Stridsberg M, Svensson M (2017) Vitamin D in relation to bone health and muscle function in young female soccer players. *Eur J Sport Sci* 17(2): 249-256. doi:10.1080/17461391.2016.1225823
13. Briggs MA, Harper LD, McNamee G, Cockburn E, Rumbold PLS, Stevenson EJ, Russell M (2017). The effects of an increased calorie breakfast consumed prior to simulated match-play in academy soccer players. *Eur J Sport Sci* 17(7): 858-866.
14. Brooks GA (1997). Importance of the "crossover" concept in exercise metabolism. *Clin Exp Pharmacol Physiol* 24(11): 889-895.
15. Brooks GA, Mercier J (1994). Balance of carbohydrate and lipid utilization during exercise: the "crossover" concept. *J Appl Physiol* 76(6): 2253-2261.
16. Burke LM, Collier GR, Broad EM, Davis PG, Martin DT, Sanigorski AJ, Hargreaves M (2003). Effect of alcohol intake on muscle glycogen storage after prolonged exercise. *J Appl Physiol* (1985) 95(3): 983-990.
17. Cairns SP (2006) Lactic acid and exercise performance: culprit or friend? *Sports Med* 36(4): 279-291.
18. Castro-Sepúlveda M, Johannsen N, Astudillo S, Jorquera C, Álvarez C, Zbinden-Foncea H, Ramírez-Campillo R (2016). Effects of beer, non-alcoholic beer and water consumption before exercise on fluid and electrolyte homeostasis in athletes. *Nutrients* 8(6): 345. doi:10.3390/nu8060345
19. Cerqueira MT, Fry MM, Connor WE (1979). The food and nutrient intakes of the Tarahumara Indians of Mexico. *Am J Clin Nutr* 32(4): 905-915.
20. Cheng IS, Wang YW, Chen IF, Hsu GS, Hsueh CF, Chang CK (2016). The supplementation of branched-chain amino acids, arginine, and citrulline improves endurance exercise performance in two consecutive days. *J Sports Sci Med* 15(3): 509-515.
21. Christensen DL, van Hall G, Hambraeus L (1998). Food intake of Kalenjin runners in Kenya: A field study. Communication to the Second Annual Conference of the European College of Sport Science (Abstract). *J Sports Sci* 16(5): 500.
22. Clark A, Mach N (2016). Exercise-induced stress behaviour, gut-microbiota-brain axis and diet: a systematic review for athletes. *J Int Soc Sports Nutr* 13: 43. doi:10.1186/s12970-016-0155-6
23. Coggan AR (1997). The glucose crossover concept is not an important new concept in exercise metabolism. *Clin Exp Pharmacol Physiol* 24(11): 896-900.
24. Daien CI, Pinget GV, Tan JK, Macia L (2017). Detrimental impact of microbiota-accessible carbohydrate-deprived diet on gut immune homeostasis: an overview. *Front Immunol*, 8: 548. doi:10.3389/fimmu.2017.00548

25. Donnelly GP, McClute N, Kennedy MS, Lewis SE (1999). Direct effect of alcohol on the motility and morphology of human spermatozoa. *Andrologia* 31(1): 43-47.
26. Dudgeon WD, Kelley EP, Scheett TP (2016). In a single-blind, matched group design: branched-chain amino acid supplementation and resistance training maintains lean body mass during a caloric restricted diet. *J Int Soc Sports Nutr* 13: 1. doi:10.1186/s12970-015-0112-9
27. European Food Safety Authority (2006). Tolerable upper intake levels for vitamins and minerals. *European Food Safety Authority*. Disponível em <http://www.efsa.eu.int>
28. Ermolao A, Zanutto T, Carraro N, Fornasier T, Zaccaria M, Neunhaeuser D, Bergamin M (2017). Repeated sprint ability is not enhanced by caffeine, arginine, and branched-chain amino acids in moderately trained soccer players. *J Exerc Rehab* 13(1): 55-61.
29. Fernández-Elías VE, Ortega JF, Nelson RK, Mora-Rodríguez R (2015). Relationship between muscle water and glycogen recovery after prolonged exercise in the heat in humans. *Eur J Appl Physiol* 115(9): 1919-1926.
30. Ferreira MP, Li R, Cooke M, Kreider RB, Willoughby DS (2014). Periexercise coingestion of branched-chain amino acids and carbohydrate in men does not preferentially augment resistance exercise-induced increases in phosphatidylinositol 3 kinase/protein kinase B-mammalian target of rapamycin pathway markers indicative of muscle protein synthesis. *Nutr Res* 34(3): 191-198. doi:10.1016/j.nutres.2013.12.007
31. Fudge BW, Westerterp KR, Kiplamai FK, Onywera VO, Boit MK, Kayser B, Pitsiladis YP (2006). Evidence of negative energy balance using doubly labelled water in elite Kenyan endurance runners prior to competition. *Bri J Nutr* 95(1): 59-66.
32. Funnell MP, Dykes NR, Owen EJ, Mears SA, Rollo I, James LJ (2017). Ecologically valid carbohydrate intake during soccer-specific exercise does not affect running performance in a fed state. *Nutrients* 9(1) pii: E39. doi:10.3390/nu9010039
33. Gebauer SK, Psota TL, Harris WS, Kris-Etherton PM (2006). n-3 fatty acid dietary recommendations and food sources to achieve essentiality and cardiovascular benefits. *Am J Clin Nutr* 83(6Suppl): S1526-S1535.
34. Gee TI, Deniel S (2016). Branched-chain amino acid supplementation attenuates a decrease in power-producing ability following acute strength training. *J Sports Med Phys Fitness*, 56(12): 1511-1517.
35. Gomez-Cabrera MC, Domenech E, Romagnoli M, Arduini A, Borrás C, Pallardo FV, Sastre J, Viña J (2008). Oral administration of vitamin C decreases muscle mitochondrial biogenesis and hampers training-induced adaptations in endurance performance. *Am J Clin Nutr* 87(1): 142-149.
36. Gravina L, Brown FF, Alexander L, Dick J, Bell G, Witard OC, Galloway SD (2017). n-3 fatty acid supplementation during 4 weeks of training leads to improved anaerobic endurance capacity, but not maximal strength, speed, or power in soccer players. *Int J Sport Nutr Exerc Metab* 7: 1-25. doi:10.1123/ijsnem.2016-0325
37. Guimarães B, Rodrigues dos Santos JA, Natal A (2004) Nutrição no futebol: Estudo comparativo do perfil nutricional dos jogadores de futebol de diferentes níveis competitivos. *Rev Port Med Desportiva* 22: 89-100.
38. Hill RJ, Bluck LJC, Davies PSW (2008). The hydration ability of three commercially available sports drinks and water. *J Sci Med Sport* 11: 116-123.
39. Ishibashi A, Maeda N, Sumi D, Goto K (2017). Elevated serum hepcidin levels during an intensified training period in well-trained female long-distance runners. *Nutrients* 9(3) Pii: E227. doi:10.3390/nu9030277
40. Journal of the American Dietetic Association (2000). Position of the American Dietetic Association, Dietitians of Canada, and the American College of Sports Medicine: Nutrition and Athletic Performance. *J Am Diet Assoc* 100(12): 1543-1556.
41. Khaled S, Brun JF, Micallet JP, Bardet L, Cassanas G, Monnier JF, Orsetti A (1997). Serum zinc and blood rheology in sportsmen (football players). *Clin Hemorheol Microcirc* 17(1): 47-58.
42. Kim YJ, Park Y, Kang DH, Kim CH (2017). Excessive exercise habits in marathoners as novel indicators of masked hypertension. *Biomed Res Int* doi:10.1155/2017/1342842.
43. Kolka MA, Latzka WA, Montain SJ, Corr WP, O'Brien KK, Sawka MN (2003). Effectiveness of revised fluid replacement guidelines for military training in hot weather. *Aviat Space Environ Med* 74(3): 242-246.
44. Koo GH, Woo J, Kang S, Shin KO (2014). Effects of supplementation with BCAA and L-glutamine on blood fatigue factors and cytokines in juvenile athletes submitted to maximal intensity rowing performance. *J Phys Ther Sci* 26(8): 1241-1246. doi:10.1589/jpts.26.1241
45. Kopec A, Solarz K, Maida F, Slowinska-Lisowska M, Medras M (2013). An evaluation of the levels of vitamin D and bone turnover markers after the summer and winter periods in Polish professional soccer players. *J Hum Kinet* 38: 135-140. doi:10.2478/hukin-2013-0053
46. Kus K, Rataiczak P, Czaia N, Zaprutko T, Nowakowska E (2017) Effect of combined administration of aripiprazole and fluoxetine on cognitive functions in female rats exposed to ethyl alcohol. *Acta Neurobiol Exp (Wars)* 77(1): 86-93.
47. Lamprecht M, Hofmann P, Greilberger JF, Schwabberger G (2009). Increased lipid peroxidation in trained men after 2 weeks of antioxidant supplementation. *Int J Sports Nutr Exerc Metab* 19(4): 385-399.
48. Lawrence GD (2013) Dietary fats and health: dietary recommendations in the context of scientific evidence. *Adv Nutr* 4(3): 294-302. doi:10.3945/an.113.003657
49. Molendi-Coste O, Legry V, Leclercq IA (2011). Why and how meet n-3 PUFA dietary recommendations? *Gastroenterol Res Pract* 2011(2011): Article ID 364040. doi:10.1155/2011/364040
50. Moore DR, Camera DM, Areta JL, Hawley JA (2014). Beyond muscle hypertrophy: why dietary protein is important for endurance athletes. *Appl Physiol Nutr Metab* 39(9): 987-997.
51. Mora-Rodríguez R, Ortega JF, Fernandez-Elias VE, Kapsokefalou M, Malisova O, Athanasatou A, Husemann M, Domnik K, Braun H (2016). Influence of physical activity and ambient temperature on hydration: The European Hydration Research Study (EHRS). *Nutrients* 8(5) pii: E252. doi:10.3390/nu8050252
52. Nyakayiru J, Jonvik KL, Trommelen J, Pinckaers PJ, Senden JM, van Loon LJ, Verdijk LB (2017). Beetroot juice supplementation improves high-intensity intermittent type exercise performance in trained soccer players. *Nutrients* 9(3) Pii: E314. doi:10.3390/nu9030314
53. Onywera VO, Kiplamai FK, Boit MK, Pitsiladis YP (2004). Food and macronutrient intake of elite Kenyan distance runners. *Int J Sport Nutr Exerc Metab* 14(6): 709-719.
54. Parr EB, Camera DM, Areta JL, Burke LM, Phillips SM, Hawley JA, Coffey VG (2014). Alcohol ingestion impairs maximal post-exercise rates of myofibrillar protein synthesis following a single bout of concurrent training. *PLoS One*, 9(2): e88384. doi:10.1371/journal.pone.0088384
55. Paulsen G, Cumming KT, Holden G, Hallén J, Ronnestad BR et al. (2014). Vitamin C and E supplementation hampers cellular adaptation to endurance training in humans: a double-blind, randomised controlled trial. *J Physiol* 592(8): 1887-1901.
56. Randell RK, Rollo I, Roberts TJ, Dalrymple KJ, Jeukendrup AE, Carter JM (2017). Maximal fat oxidation rates in an athletic population. *Med Sci Sports Exerc* 49(1): 133-140.
57. Reis R, Charehsaz M, Sipahi H, Ekici AI, Macit C, Akkaya H, Aydin A (2017). Energy drink induced lipid peroxidation and oxidative damage in rat live and brain when used alone or combined with alcohol. *J Food Sci* 82(4): 1037-1043.
58. Rodrigues dos Santos JA (1999). Estudo comparativo, fisiológico, antropométrico e motor entre futebolistas de diferente nível competitivo. *Rev Paul Educ Fis* 13(2): 146-159.
59. Rodrigues dos Santos JA, Vasconcelos CEGC (2009). Nutrição e composição corporal de jovens futebolistas. *Rev Bras Fisiol Exerc* 8(3): 113-120.
60. Romijn JA, Coyle EF, Sidossis LS, Gastaldelli A, Horowitz JF, Ender E, Wolfe RR (1993). Regulation of endogenous fat and carbohydrate metabolism in relation to exercise intensity and duration. *Am J Physiol* 265(3 Pt 1): E380-E391.
61. Sahlin K, Harris RC (2008). Control of lipid oxidation during exercise: role of energy state and mitochondrial factors. *Acta Physiol (Oxford)* 194(4): 283-291. doi:10.1111/j.1748-1716.2008.01879
62. Sawka MN, Cheuvront SN, Kenefick RW (2015). Hypohydration and human performance: impact of environment and physiological mechanisms. *Sports Med* 45(Suppl 1): S51-S60. doi:1007/s40279-015-0395-7
63. Shayakhmetova GM, Bondarenko LB, Matvienko AV, Kovalenko VM (2014). Chronic alcoholism-mediated metabolic disorders in albino rat testes. *Interdiscip Toxicol* 7(3): 165-172.
64. Stepanyan V, Crowe M, Haleagrahara N, Bowden B (2014). Effects of vitamin E supplementation on exercise-induced oxidative stress: a meta-analysis. *Appl Physiol Nutr Metab* 39(9): 1029-1037.
65. Sutton L, Scott M, Wallace J, Reilly T (2009). Body composition of English Premier League soccer players: influence of playing position, international status, and ethnicity. *J Sports Sci* 27(10): 1019-1026.
66. Tang FC, Chan CC (2016). Contribution of branched-chain amino acids to purine nucleotide cycle: a pilot study. *Eur J Clin Nutr* 71(5): 587-593. doi:10.1038/ejcn.2016.161
67. Tester R, Al-Ghazzewi FH (2017). Role of glucomannans in immunology. *J Pharm Pharm Sci* 20(0): 97-114. doi:10.18433/J31P52.
68. Wenger CB (1972). Heat of evaporation of sweat: thermodynamic considerations. *J Appl Physiol* 32(4): 456-459.
69. Williams RJP (2002). The fundamental nature of life as a chemical system: the part played by inorganic elements. *J Inorg Biochem* 88: 241-250.
70. Xiao W, Chen P, Liu X, Zhao L (2015) The impaired function of macrophages induced by strenuous exercise could not be ameliorated by BCAA supplementation. *Nutrients* 7(10): 8645-8656. doi:10.3390/nu71054525
71. Yfanti C, Akerström T, Nielsen S, Nielsen AR, Mounier R, Mortensen OH, Lykkesfeldt J, Rose AJ, Fischer CP, Pedersen BK (2010). Antioxidant supplementation does not alter endurance training adaptation. *Med Sci Sports Exerc* 42(7): 1388-1395.