Statistica Sinica Preprint No: SS-2019-0467	
Title	Estimation and Inference for Dynamic Single-Index Varying-Coefficient Models
Manuscript ID	SS-2019-0467
URL	http://www.stat.sinica.edu.tw/statistica/
DOI	10.5705/ss.202019.0467
Complete List of Authors	Xin Guan, Hua Liu, Jinhong You and Yong Zhou
Corresponding Author	Jinhong You
E-mail	johnyou07@163.com
Notice: Accepted version subject to English editing.	

ESTIMATION AND INFERENCE FOR DYNAMIC SINGLE-INDEX VARYING-COEFFICIENT MODELS

Xin Guan¹, Hua Liu², Jinhong You^{2,3}, and Yong Zhou⁴

¹*Zhongnan University of Economics and Law,*

²*Shanghai University of Finance and Economics,*

³*Shanghai Lixin University of Accounting and Finance,*

⁴*East China Normal University*

Abstract: Motivated by applications, we propose a class of dynamic single-index varying-coefficient models to explore the varying interaction effects on the response variable among a set of covariates. That is to say, the interaction effects are allowed to change with some factors of interest, such as time, spatial location, or other covariates. A spline-based approach is developed to estimate the index and varying-coefficient functions. The convergence rates and asymptotic normalities of the resulting estimators are established. It is also shown that the resulting estimators exhibit an oracle property. A penalized method is presented to select related covariates, and the consistency of the penalized estimator is proved. A test statistic is provided to check whether the interaction effect also varies with the factors of interest, and the asymptotic normality of the test statistic is

1. INTRODUCTION₂

established. Simulation studies and two real data analyses illustrate the good performance of the proposed model and the corresponding statistical inference methods for finite samples.

Key words and phrases: Interaction effect, Single-index varying-coefficient regression model, Spline approximation, Variable selection.

1. Introduction

Single-index varying-coefficient regression models (SICMs), proposed by Xia and Li (1999), have attracted much attention over the past few decades and have proven to be effective in practical applications; see Fan, Yao and Cai (2003), Ma and Song (2015), Liu, Cui and Li (2016), and references therein. SICMs can capture the nonlinear relationship between the response and related covariates, and they can also achieve dimensional reduction by taking a linear combination of covariates as an index. A classical SICM is defined by

$$Y = \sum_{l=1}^d g_l(\mathbf{Z}^T \boldsymbol{\beta}) X_l + \varepsilon, \quad (1.1)$$

where Y is the response variable, $(\mathbf{Z}^T, \mathbf{X}^T)^T$ is a vector of covariates consisting of $\mathbf{X} = (X_1, X_2, \dots, X_d)^T$ with $X_1 = 1$ and $\mathbf{Z} \in \mathbb{R}^p$, $\boldsymbol{\beta} \in \mathbb{R}^p$ is the coefficient parameter vector, $g_l(\cdot), l = 1, \dots, d$, are unknown smooth functions that describe the nonlinear relationship between Y and the covariates,

1. INTRODUCTION₃

ε is the error term with mean zero, and superscript T denotes the transpose of a vector or matrix. From (1.1), it is easy to see that the effects of interaction between $\mathbf{X}$ and $\mathbf{Z}$ in the SICM are static. However, this may be not true in many applications, where the interaction effects of some covariates may depend on factors of interest, such as time and spatial location, or on other covariates. For example, in an analysis of air pollution data in Guangzhou, China, we found that the interaction effects on the air quality index of the next day between four air pollutants (nitrogen dioxide, ozone, fine particles, and inhalable particles) and two environmental factors varied with wind direction; see Figures 3(a), 4(a), and 4(b). As another example, for the WeChat public accounts data, the interaction relationship between reading volume and the attributes of the posted articles varies with the proportion of articles posted during 18:00–02:00, as can be seen in Figures 3(b) and 5(a)–5(e). Thus, SICMs may be inappropriate in these situations with varying interaction effects. To explore the possible varying interaction effect patterns, we develop a class of dynamic single-index varying-coefficient models (DSICMs) defined by

$$Y = \sum_{l=1}^d g_l(\mathbf{Z}^T \boldsymbol{\beta}(U)) X_l + \varepsilon, \quad (1.2)$$

where U is a scalar, which may be one of the entries of $\mathbf{Z}$ or $\mathbf{X}$ or some other variable, $\boldsymbol{\beta}(u) = (\beta_k(u), 1 \leq k \leq p)^T$ is a p -dimensional varying-coefficient

---

1. INTRODUCTION4

---

function with nonparametric covariate U , and ε is an independent and identically distributed (i.i.d.) random error satisfying $E(\varepsilon|\mathbf{Z}, \mathbf{X}, U) = 0$ and $\text{Var}(\varepsilon|\mathbf{Z}, \mathbf{X}, U) = \sigma^2(\mathbf{Z}, \mathbf{X}, U)$.

Obviously, the model (1.2) includes many important models as special cases. For example, when $\beta(U) \equiv \beta$, it reduces to the classical SICM (1.1), and when $d = 1$ and $X_l = 1$, it reduces to the class of single-index varying-coefficient models (SIVCs) studied by Luo, Zhu and Zhu (2016) and Ma (2016) or the generalized varying-coefficient models with unspecified link functions studied by Chen, Hall and Müller (2011), Kuruwita, Kulasekera and Gallagher (2011), Zhang, Li and Xia (2015), and others. In fact, Chen, Hall and Müller (2011) were the first to study the generalized functional linear model with an unspecified link function. They approximated the link function by kernel smoothing and established a polynomial convergence rate for it. Kuruwita, Kulasekera and Gallagher (2011) developed a locally constant approximation to the varying-coefficient functions and the link function. They established pointwise-asymptotic properties of the estimated varying-coefficient functions. Zhang, Li and Xia (2015) provided an iterative procedure to approximate the varying-coefficient functions and the unspecified link function by using a two-dimensional kernel. Luo, Zhu and Zhu (2016) proposed a SIVC model and derived asymptotic properties

1. INTRODUCTION⁵

---

of varying-coefficient functions and index functions by optimal integration of information across all grid points. Ma (2016) considered a functional single-index model and established uniform convergence rates of coefficient functions and index functions, based on B-spline basis expansions. The convergence rates in their work were slower than the classical uniform non-parametric rates.

We propose a B-spline-based method to estimate the unknown functions in a DSICM with the model identification criterion from Ma (2016). The basic idea is to approximate the smooth index functions $g_l(\cdot)$ via a linear assumption and give initial estimates of the varying-coefficient functions $\beta_k(\cdot)$. We then fit two varying-coefficient models (VCMs) (Hastie and Tibshirani, 1993) to obtain the final spline estimators of the index functions and varying-coefficient functions, respectively. We show that the resulting estimators are consistent and that each estimator attains the optimal convergence rate of a univariate nonparametric function. We derive the asymptotic normality of the estimators and present asymptotic simultaneous confidence bands (SCBs) for each univariate function. Variable selection is an important aspect of nonparametric models. As in the case of the traditional single-index model (1.1), our primary concern is to select relevant predictors $\mathbf{Z}$ that modify the relationship between Y and $\mathbf{X}$, that is, to pick

---

1. INTRODUCTION₆

out nonzero functions $\beta_k(\cdot)$ in the model (1.2) and improve the estimation accuracy. Many methods and inferences for nonparametric model selection have been developed; see, for example, Wang, Li and Huang (2008) and Hu, Huang and You (2019). We propose an identification method based on a SCAD penalty function for varying-coefficient functions and estimate the varying-coefficient functions and the index functions simultaneously. We show that our estimation procedure is consistent, that is, the true model is correctly selected with probability tending to one. Another important contribution of this paper is that we develop a test statistic to check whether the varying-coefficient function vector $\beta(\cdot)$ varies with the factors of interest, and we derive the asymptotic distribution of the proposed test statistic.

The rest of this article is organized as follows. The three-step estimation procedure is described in Section 2. The asymptotic theory of the proposed estimators is presented in Section 3. Section 4 describes the model identification method and the hypothesis testing procedure for the varying-coefficient components. Sections 5 and 6 present simulation studies and two real data analyses to demonstrate the finite-sample performance of the DSICM. Some conclusions are discussed in Section 7. A description of notation and technical proofs are relegated to the Supplementary Materials.

2. Spline Estimation Method

2.1 Model identifiability

A single-index model is not identifiable in the absence of constraints on its structure. For the semiparametric single-index model (1.1), Xia et al. (2002) and Fan, Yao and Cai (2003) developed a standard model identification method by assuming that the parameter vector has a positive first component and norm equal to one. For the nonparametric single-index model (1.2), Kuruwita, Kulasekera and Gallagher (2011) and Luo, Zhu and Zhu (2016) employed a pointwise identification condition by simply borrowing the idea from the model (1.1). Zhang, Li and Xia (2015) pointed out that the restriction to pointwise identifiability imposed some limitations, and they proposed a weaker model identification condition, based on a binary function that separated the direction and norm of each varying-coefficient function. In the present paper, similarly to Ma (2016), we adopt a model identification condition based on a spline approximation as follows.

Proposition 1. *Assume that $\|\beta(u)\|_{L_2} = 1$ and that $\beta_1(u)$ is nonconstant and monotone nondecreasing over $u \in S_U$. Then, the model (1.2) is identifiable.*

2. SPLINE ESTIMATION METHOD8

2.2 Estimation

Suppose $\{(Y_i, \mathbf{X}_i^T, \mathbf{Z}_i^T, U_i), 1 \leq i \leq n\}$ are the i.i.d. realizations of $(Y, \mathbf{X}^T, \mathbf{Z}^T, U)$. Let $\mathbf{B}_1(w) = (B_{m,1}(w), 1 \leq m \leq J_{n,1})^T$ and $\mathbf{B}_2(u) = (B_{s,2}(u), 1 \leq s \leq J_{n,2})^T$ be B-spline bases of orders q_1 and q_2 , respectively. The number of basis functions is defined as $J_{n,t} = K_t + q_t$, where K_t is the number of interior knots for a knot sequence $\zeta_1 = \dots = 0 = \zeta_{q_t} < \zeta_{q_t+1} < \dots < \zeta_{q_t+K_t} < 1 = \zeta_{q_t+K_t+1} = \dots = \zeta_{2q_t+K_t}$ and K_t increases with the sample size n , $t = 1, 2$. Here we adopt a spline-based method to estimate the varying-coefficient functions and index functions in three steps:

Step 0: Obtain an initial spline approximation $\hat{\beta}_k^{(0)}(\cdot) \approx \mathbf{B}_2(u)^T \hat{\boldsymbol{\delta}}_k^{(0)}$ by fitting a linear structure to the index functions, where $\hat{\boldsymbol{\delta}}_k^{(0)} = (\hat{\delta}_{sk}^{(0)}, 1 \leq s \leq J_{n,2})^T$ are the estimated spline coefficients.

Step 1: Denote $\Phi_{i,sk} = B_{s,2}(U_i)Z_{ik}$ and $\boldsymbol{\Phi}_i = (\boldsymbol{\Phi}_{i,1}^T, \dots, \boldsymbol{\Phi}_{i,p}^T)^T$, with $\boldsymbol{\Phi}_{i,k} = (\Phi_{i,1k}, \dots, \Phi_{i,J_{n,2}k})^T$. At iteration s , given $\hat{\boldsymbol{\delta}}^{(s-1)} = (\hat{\boldsymbol{\delta}}_1^{(s-1),T}, \dots, \hat{\boldsymbol{\delta}}_p^{(s-1),T})^T$, we obtain for the model (1.2) the following approximated VCM: $Y_i \approx \sum_{l=1}^d g_l(\boldsymbol{\Phi}_i^T \hat{\boldsymbol{\delta}}^{(s-1)})X_{il} + \varepsilon_i$. Suppose that $\hat{\boldsymbol{\lambda}}^{(s)}(\hat{\boldsymbol{\delta}}^{(s-1)}) = (\hat{\lambda}_{m,l}^{(s)}, 1 \leq m \leq J_{n,1}, 1 \leq l \leq d)^T$ is obtained by minimizing $\sum_{i=1}^n \left[Y_i - \sum_{l=1}^d \sum_{m=1}^{J_{n,1}} B_{m,1}(\boldsymbol{\Phi}_i^T \hat{\boldsymbol{\delta}}^{(s-1)}) \lambda_{m,l} X_{il} \right]^2$. Denote $W_i = \mathbf{Z}_i^T \boldsymbol{\beta}(U_i)$ and $\widehat{W}_i^{(s)} = \mathbf{Z}_i^T \widehat{\boldsymbol{\beta}}^{(s)}(U_i) = \boldsymbol{\Phi}_i^T \hat{\boldsymbol{\delta}}^{(s)}$. Let $\mathbf{D}_i(W_i) = \{D_{i,ml}(W_i), 1 \leq l \leq d, 1 \leq m \leq J_{n,1}\}^T$,

2. SPLINE ESTIMATION METHOD9

with $D_{i,ml}(W_i) = B_{m,1}(W_i)X_{i,l}$. Similarly, we define $\mathbf{D}_i(\widehat{W}_i^{(s)})$ by replacing W_i with $\widehat{W}_i^{(s)}$. Let $\mathbf{W} = (W_1, \dots, W_n)^T$ and $\mathbf{D}(\widehat{\mathbf{W}}^{(s)}) = [(\mathbf{D}_1(\widehat{W}_1^{(s)}), \dots, \mathbf{D}_n(\widehat{W}_n^{(s)}))^T]_{n \times J_{n,1}d}$. Thus,

$$\hat{\boldsymbol{\lambda}}^{(s)}(\hat{\boldsymbol{\delta}}^{(s-1)}) = [\mathbf{D}(\widehat{\mathbf{W}}^{(s-1)})^T \mathbf{D}(\widehat{\mathbf{W}}^{(s-1)})]^{-1} \mathbf{D}(\widehat{\mathbf{W}}^{(s-1)})^T \mathbf{Y}. \quad (2.1)$$

The spline estimator of the index function is then given by $\hat{g}_l^{(s)}(w, \hat{\boldsymbol{\delta}}^{(s-1)}) = \sum_{m=1}^{J_{n,1}} B_{m,1}(w) \hat{\lambda}_{m,l}^{(s)}(\hat{\boldsymbol{\delta}}^{(s-1)})$, $l = 1, \dots, d$.

Step 2: At iteration s , when the index components are known, the model (1.2)

is specified as a VCM. Suppose that $\hat{\boldsymbol{\delta}}^{(s)}$ minimizes

$$L(\boldsymbol{\delta}) = \frac{1}{2} \sum_{i=1}^n \left[Y_i - \sum_{l=1}^d \mathbf{B}_1(\boldsymbol{\Phi}_i^T \boldsymbol{\delta})^T \hat{\boldsymbol{\lambda}}_l^{(s)}(\hat{\boldsymbol{\delta}}^{(s-1)}) X_{il} \right]^2, \quad (2.2)$$

subject to $\delta_{11} \leq \dots \leq \delta_{J_{n,2}1}$. Then, $\hat{\beta}_k^{(s)}(u) = \sum_{s=1}^{J_{n,2}} B_{s,2}(u) \hat{\delta}_{sk}^{(s)}$ for $k = 1, \dots, p$, and we further identify it according to Proposition 1.

The constraint condition $\delta_{11} \leq \dots \leq \delta_{J_{n,2}1}$ is only used to ensure that $\beta_1(u)$ is nondecreasing. We use the “constrOptim” function in R to deal with the constrained optimization problem (2.2), and the gradient function is given in the Supplementary Materials.

Step 3: Repeat Steps 1 and 2 until convergence, and we then obtain $\hat{\beta}_k(\cdot)$ and $\hat{g}_l(\cdot)$.

3. ASYMPTOTIC PROPERTIES OF THE ESTIMATION METHOD₁₀

3. Asymptotic Properties of the Estimation Method

In this section, we obtain asymptotic results for the three-step spline estimators of the varying-coefficient components and the index components.

Like Li and Lv (2020) and Ma and Song (2015), we focus on the space $\mathcal{M}$ as the following collection of dynamic single-index varying-coefficient functions

with finite L_2 norm: $\mathcal{M} = \left\{ f : f(\mathbf{x}, \mathbf{z}, u) = \sum_{l=1}^d f_l(\mathbf{z}^T \boldsymbol{\beta}(u)) x_l, E f^2(\mathbf{x}, \mathbf{z}, u) < \infty \right\}$, with $\mathbf{z} = (z_1, \dots, z_p)^T$, $\mathbf{x} = (x_1, \dots, x_d)^T$, and $u \in [0, 1]$. For

$1 \leq k \leq p$, $1 \leq s \leq J_{n,2}$, we assume that f_{sk}^0 satisfies

$$\mathbb{P}(\Phi_{sk}) = \sum_{l=1}^d f_{l,sk}^0(\mathbf{z}^T \boldsymbol{\beta}^0(u)) x_l = \arg \min_{f \in \mathcal{M}} E\{\Phi_{sk} - f(\mathbf{z}^T \boldsymbol{\beta}^0(u), \mathbf{x})\}^2, \quad (3.1)$$

which implies a projection of the variable $\boldsymbol{\Phi}$ onto the space $\mathcal{M}$. Let $\mathbb{P}(\boldsymbol{\Phi}) =$

$\{\mathbb{P}(\boldsymbol{\Phi}_1)^T, \dots, \mathbb{P}(\boldsymbol{\Phi}_p)^T\}^T$ with $\mathbb{P}(\boldsymbol{\Phi}_k) = \{\mathbb{P}(\boldsymbol{\Phi}_{1k}), \dots, \mathbb{P}(\boldsymbol{\Phi}_{J_{n,2}k})\}^T$, $k = 1, \dots, p$.

Denote $\tilde{\boldsymbol{\Phi}} = \boldsymbol{\Phi} - \mathbb{P}(\boldsymbol{\Phi})$.

3.1 Assumptions

We make the following assumptions:

- (C1) The density function $f_W(\cdot)$ of the random variable $W = \mathbf{Z}^T \boldsymbol{\beta}(U)$ is bounded away from 0 on S_W , and $f_W(\cdot) \in C^{0,1}(S_W, c)$ for $\boldsymbol{\beta}(u)$ in the neighborhood of $\boldsymbol{\beta}^0(u)$, with $S_W = \{\mathbf{Z}^T \boldsymbol{\beta}(U), \mathbf{Z} \in S_Z, U \in S_U\}$, where S_Z and S_U are compact support sets of $\mathbf{Z}$ and U , respectively.

3. ASYMPTOTIC PROPERTIES OF THE ESTIMATION METHOD₁₁

Without loss of generality, we assume that $S_U = [0, 1]$, $S_W = [a, b]$, and $\mathbf{X}$ is bounded.

(C2) $\beta_k(\cdot) \in C^{(r)}[0, 1]$, $1 \leq k \leq p$, and $g_l(\cdot) \in C^{(r)}[a, b]$, $1 \leq l \leq d$, for some integer $r \geq 2$, and the spline order satisfies $q \geq r$.

(C3) The conditional variance function $\sigma^2(\mathbf{x}, \mathbf{z}, u)$ is measurable and bounded away from some constant $0 < C_\sigma < \infty$.

(C4) There exist constants $0 < c_Q \leq C_Q < \infty$ such that $c_Q \leq Q(\mathbf{z}, u) = E(\mathbf{X}\mathbf{X}^T | \mathbf{Z} = \mathbf{z}, U = u) \leq C_Q$ for all $\mathbf{z} \in S_Z$ and $u \in [0, 1]$.

(C5) For $1 \leq k \leq p$, $1 \leq s \leq J_{n,2}$, $f_{sk}^0(\cdot) \in C^{(2)}[a, b]$.

(C6) $\Xi_1 = E\{\mathbf{D}_i(W_i)\mathbf{D}_i(W_i)^T\}$, and the eigenvalues of Ξ_1 are bounded away from zero and infinity.

(C7) $\Xi_2 = E\{[\sum_{l=1}^d \dot{g}_l(W_i)X_{il}\tilde{\Phi}_i]^{\otimes 2}\}$, and the eigenvalues of Ξ_2 are bounded away from zero and infinity.

Remark 1. Condition (C1) is the same as in Tang and Cheng (2008), Ma and Song (2015), and Zhao, Lian and Liang (2017). The boundedness of X is assumed mainly for convenience in the proof and can be replaced by a weaker assumption; see Zhao, Lian and Liang (2017). Conditions (C2)–(C4) are common conditions for nonparametric models; see Ma and Song (2015),

3. ASYMPTOTIC PROPERTIES OF THE ESTIMATION METHOD₁₂

Ma (2016), and Liu, Cui and Li (2016). Condition (C5) is a smoothness condition on the function f_{sk}^0 defined in (3.1). Conditions (C6) and (C7) are necessary to derive the asymptotic theorem.

3.2 Asymptotic results

According to de Boor (2001), for $\beta_k(\cdot)$ and $g_l(\cdot)$ satisfying Condition (C2), there exist spline functions $g_l^0(w) = \sum_{m=1}^{J_{n,1}} B_{m,1}(w) \lambda_{m,l}^0 = \mathbf{B}_1(w)^T \boldsymbol{\lambda}_l^0$ with $\boldsymbol{\lambda}_l^0 \in \mathbb{R}^{J_{n,1}}$ and $\beta_k^0(u) = \sum_{s=1}^{J_{n,2}} B_{s,2}(u) \delta_{s,k}^0 = \mathbf{B}_2(u)^T \boldsymbol{\delta}_k^0$ with $\boldsymbol{\delta}_k^0 \in \mathbb{R}^{J_{n,2}}$ such that

$$\sup_{w \in S_W} |g_l^0(w) - g_l(w)| = O(J_{n,1}^{-r}), \quad \sup_{u \in S_U} |\beta_k^0(u) - \beta_k(u)| = O(J_{n,2}^{-r}). \quad (3.2)$$

Denote $\boldsymbol{\delta}^0 = (\boldsymbol{\delta}_1^{0T}, \dots, \boldsymbol{\delta}_p^{0T})^T$. The following proposition gives the uniform convergence rates of the spline estimators $\hat{g}_l(w; \boldsymbol{\delta})$ and $\hat{g}_l(w; \boldsymbol{\delta})$ near $\boldsymbol{\delta}^0$.

Proposition 2. *Under Conditions (C1) and (C2), when $J_{n,1}^3 n^{-1} \log n = o(1)$, $J_{n,1} J_{n,2}^{-r} = o(1)$, and $a_n = o(J_{n,1}^{-1})$, for $l = 1, \dots, d$, we have (i) $\sup_{\|\boldsymbol{\delta} - \boldsymbol{\delta}^0\|_\infty \leq a_n} \sup_{w \in S_W} |\hat{g}_l(w, \boldsymbol{\delta}) - g_l(w)| = O_p(a_n + J_{n,1}^{-r} + J_{n,2}^{-r} + (\log n)^{1/2} J_{n,1}^{1/2} n^{-1/2})$; (ii) $\sup_{\|\boldsymbol{\delta} - \boldsymbol{\delta}^0\|_\infty \leq a_n} \sup_{w \in S_W} |\hat{g}_l(w, \boldsymbol{\delta}) - \dot{g}_l(w)| = O_p(J_{n,1} a_n + J_{n,1} J_{n,2}^{-r} + J_{n,1}^{1-r} + (\log n)^{1/2} J_{n,1}^{3/2} n^{-1/2})$.*

Remark 2. Let $\hat{\boldsymbol{\delta}}$ be the minimizer of $L(\boldsymbol{\delta})$ given in (2.2) satisfying $\|\hat{\boldsymbol{\delta}} - \boldsymbol{\delta}^0\| \leq a_n$ with probability tending to one. This implies that we are assuming

3. ASYMPTOTIC PROPERTIES OF THE ESTIMATION METHOD₁₃

that there exists a local minimizer in the neighborhood of δ^0 in probability.

A similar restriction has been made in Xia and Li (1999) and Ma (2016).

Let $\mathbb{B}_1(w) = \mathbf{I}_d \otimes \mathbf{B}_1(w)^T$ and $\mathbb{B}_2(u) = \mathbf{I}_p \otimes \mathbf{B}_2(u)^T$, where $\mathbf{I}_d$ and $\mathbf{I}_p$ are the $d \times d$ and $p \times p$ identity matrices. Let $\mathbf{e}_l$ be the $d \times 1$ vector with the l th element being 1 and all other elements being 0 and let $\mathbf{b}_k$ be the $p \times 1$ vector with the k th element being 1 and all other elements being 0. Define $d_{1n} = 1 - [2 \log(K_1 + 1)]^{-1} \{ \log[-0.5 \log(1 - \alpha)] + 0.5 \{ \log[\log(K_1 + 1)] + \log(4\pi) \} \}$, and let d_{2n} be the same as d_{1n} except that K_1 is replaced by K_2 . Let $Q_{n1}(\alpha) = [2 \log(K_1 + 1)]^{1/2} d_{1n}$ and $Q_{n2}(\alpha) = [2 \log(K_2 + 1)]^{1/2} d_{2n}$.

The following theorems give the asymptotic results and construct asymptotic simultaneous confidence bands (SCBs) for the varying-coefficient component $\hat{\beta}_k(\cdot)$ and the index component $\hat{g}_l(\cdot)$.

Theorem 1. *Under Conditions (C1)–(C7), if $n^{1/(2r+2)} \ll J_{n,1} \ll n^{1/4}$ and $n^{1/(2r+2)} \ll J_{n,2} \ll n^{1/4}$, then we have (i) $\|\hat{\beta}(u) - \beta(u)\|_{L_2} = O_p(J_{n,2}^{-r} + J_{n,2}^{1/2} n^{-1/2})$; (ii) $\sup_{u \in S_U} |\hat{\beta}(u) - \beta(u)| = O_p(J_{n,2}^{-r} + (\log n)^{1/2} J_{n,2}^{1/2} n^{-1/2})$; (iii) if $n J_{n,2}^{-(2r+1)} = o(1)$, then $\Omega_2^{-1}(u) [\hat{\beta}(u) - \beta(u)] \xrightarrow{d} N(0, 1)$, where $\Omega_2(u) = n^{-1} \mathbb{B}_2(u) \Xi_2^{-1} E \left\{ \sigma^2(U_i, \mathbf{Z}_i, \mathbf{X}_i) \left[\sum_{l=1}^d \dot{g}_l(W_i) X_{il} \tilde{\Phi}_i \right]^{\otimes 2} \right\} \Xi_2^{-1} \mathbb{B}_2(u)^T$. Further, for $k = 1, \dots, p$, we have $\sigma_{2,k}^{-1}(u) [\hat{\beta}_k(u) - \beta_k(u)] \xrightarrow{d} N(0, 1)$, where $\sigma_{2,k}^2(u) = \mathbf{b}_k^T \Omega_2(u) \mathbf{b}_k$.*

Remark 3. Let $J_{n,2} \asymp n^{1/(2r+1)}$, then the spline estimates of the varying-

3. ASYMPTOTIC PROPERTIES OF THE ESTIMATION METHOD₁₄

coefficient components attain the optimal L_2 -norm nonparametric rate

$O_p(n^{-r/(2r+1)})$ as well as the optimal uniform nonparametric rate $O_p((\log n)^{1/2} n^{-r/(2r+1)})$.

Theorem 2. *Under Conditions (C1)–(C7), if $n^{1/(2r+2)} \ll J_{n,1} \ll n^{1/4}$ and $n^{1/(2r+2)} \ll J_{n,2} \ll J_{n,1}$, then, for $l = 1, \dots, d$, we have (i) $\sup_{w \in S_W} |\hat{g}_l(w; \hat{\boldsymbol{\delta}}) - g_l(w)| = O_p(J_{n,1}^{-r} + J_{n,2}^{-r} + (\log n)^{1/2} J_{n,1}^{1/2} n^{-1/2} + (\log n)^{1/2} J_{n,2}^{1/2} n^{-1/2})$; (ii) if $J_{n,2} \ll J_{n,1}$ and $nJ_{n,2}^{-(2r+1)} = o(1)$, then $\sigma_{1,l}^{-1}(w)[\hat{g}_l(w; \hat{\boldsymbol{\delta}}) - g_l(w)] \xrightarrow{d} N(0, 1)$, where $\sigma_{1,l}^2(w) = n^{-1} \mathbf{e}_l^T \mathbb{B}_1(w) \Xi_1^{-1} [E(\sigma^2(U_i, \mathbf{Z}_i, \mathbf{X}_i) \mathbf{D}_i(W_i)^{\otimes 2})] \Xi_1^{-1} \mathbb{B}_1^T(w) \mathbf{e}_l$.*

Remark 4. Let $J_{n,1} \asymp J_{n,2} \asymp n^{1/(2r+1)}$, then the spline estimates of the index components attain the optimal uniform nonparametric rate $O_p((\log n)^{1/2} n^{-r/(2r+1)})$. The conditions $J_{n,2} \ll J_{n,1}$ and $nJ_{n,2}^{-(2r+1)} = o(1)$ are only imposed to make the bias asymptotically negligible. It can be seen that the asymptotic variance of $\hat{\beta}_k(\cdot)$ obtained from Theorem 1 is the same as would be obtained if all the index functions $g_l(\cdot), l = 1, \dots, d$, were known. Analogously, the asymptotic variance of $\hat{g}_l(\cdot)$ obtained from Theorem 2 is the same as would be obtained if all the $\beta_k(\cdot), k = 1, \dots, p$, were known. Hence, the proposed spline estimator has the oracle property. If $\sigma^2(\mathbf{Z}, \mathbf{X}, U) = \sigma^2$, then the asymptotic variance matrices are reduced to $\sigma_{1,l}^2(w) = n^{-1} \sigma^2 \mathbf{e}_l^T \mathbb{B}_1(w) \Xi_1^{-1} \mathbb{B}_1^T(w) \mathbf{e}_l$ and $\sigma_{2,k}^2(u) = n^{-1} \sigma^2 \mathbf{b}_k^T \mathbb{B}_2(u) \Xi_2^{-1} \mathbb{B}_2^T(u) \mathbf{b}_k$.

3. ASYMPTOTIC PROPERTIES OF THE ESTIMATION METHOD¹⁵

Theorem 3. Under Conditions (C1)–(C7), if $n^{1/(2r+2)} \ll J_{n,1} \ll n^{1/4}$, $n^{1/(2r+2)} \ll J_{n,2} \ll n^{1/4}$, and $nJ_{n,2}^{-(2r+1)} = o(1)$, then, for any $\alpha \in (0, 1)$, we have $\lim_{n \rightarrow \infty} P\left[\sup_{u \in S_{n,2}} |\sigma_{2,k}^{-1}(u)\{\hat{\beta}_k(u) - \beta_k(u)\}| \leq Q_{n2}(\alpha)\right] = 1 - \alpha$, where $S_{n,2}$ is a subset of S_U that becomes denser as $n \rightarrow \infty$. Therefore, an asymptotic $100(1 - \alpha)\%$ SCB for $\beta_k(u)$ over $u \in S_{n,2}$ is $\hat{\beta}_k(u) \pm \sigma_{2,k}(u)Q_{n2}(\alpha)$.

Theorem 4. Under Conditions (C1)–(C7), if $n^{1/(2r+2)} \ll J_{n,1} \ll n^{1/4}$, $n^{1/(2r+2)} \ll J_{n,2} \ll n^{1/4}$, $J_{n,2} \ll J_{n,1}$, and $nJ_{n,2}^{-(2r+1)} = o(1)$, then, for any $\alpha \in (0, 1)$, we have $\lim_{n \rightarrow \infty} P\left[\sup_{w \in S_{n,1}} |\sigma_{1,l}^{-1}(w)\{\hat{g}_l(w; \hat{\boldsymbol{\delta}}) - g_l(w)\}| \leq Q_{n1}(\alpha)\right] = 1 - \alpha$, where $S_{n,1}$ is a subset of S_W that becomes denser as $n \rightarrow \infty$. Therefore, an asymptotic $100(1 - \alpha)\%$ SCB for $\hat{g}_l(w; \hat{\boldsymbol{\delta}})$ over $w \in S_{n,1}$ is $\hat{g}_l(w; \hat{\boldsymbol{\delta}}) \pm \sigma_{1,l}(w)Q_{n1}(\alpha)$.

Remark 5. Compared with the asymptotic $100(1 - \alpha)\%$ point confidence bands (PCBs): $\hat{g}_l(u) \pm \sigma_{1,l}(u)z_\alpha$ and $\hat{\beta}_k(u) \pm \sigma_{2,k}(u)z_\alpha$, the widths of the SCBs are inflated by rates $[2 \log(K_1 + 1)]^{1/2}d_{1n}/z_\alpha$ and $[2 \log(K_2 + 1)]^{1/2}d_{2n}/z_\alpha$, respectively, where z_α is the α -quantile of the standard normal distribution.

To establish the asymptotic confidence bands of $\hat{g}_l(\cdot)$ and $\hat{\beta}_k(\cdot)$, the key issue is to obtain accurate variance estimates $\hat{\sigma}_{1,l}$ and $\hat{\sigma}_{2,k}$. On the suggestion of Yang (2008) and Ma et al. (2015), we adopt a wild bootstrap procedure to approximate the asymptotic variances, since the plug-in method may lead to inaccurate results for finite samples.

4. Model Selection

4.1 Model identification

In this subsection, we present a model identification procedure based on a penalized method to select significant variables Z_k with nonzero varying-coefficient functions $\beta_k(\cdot)$. Without loss of generality, we assume that $\beta_k(u), k = 1, \dots, p_1$, are nonzero varying-coefficient functions and $\beta_k(u) = 0$ for $k = p_1 + 1, \dots, p$ in the true model.

Denote $\mathbf{H} = (h_{ij})_{J_{n,2} \times J_{n,2}}$ with entries $h_{ij} = \int_U B_{i,2}(u)B_{j,2}(u) du$. Let $\boldsymbol{\pi} = (\boldsymbol{\pi}_1^T, \dots, \boldsymbol{\pi}_p^T)^T$, with $\boldsymbol{\pi}_k = (\pi_{1k}, \dots, \pi_{J_{n,2}k})^T$, $1 \leq k \leq p$. For $\beta_k(u) = \mathbf{B}_2(u)^T \boldsymbol{\pi}_k$, let $\|\beta_k(u)\|_{L_2}^2 = \boldsymbol{\pi}_k^T \mathbf{H} \boldsymbol{\pi}_k = \|\boldsymbol{\pi}_k\|_{\mathbf{H}}^2$. We adopt the penalized approach for varying-coefficient functions presented in Wang, Li and Huang (2008). Define the objective function with penalization as

$$\begin{aligned} Q(\boldsymbol{\pi}) &= \frac{1}{2} \sum_{i=1}^n \left[Y_i - \sum_{l=1}^d \hat{g}_l(\boldsymbol{\Phi}_i^T \boldsymbol{\pi}) X_{il} \right]^2 + n \sum_{k=2}^p p_{\alpha_n}(\|\boldsymbol{\pi}_k\|_{\mathbf{H}}) \\ &= L(\boldsymbol{\pi}) + n \sum_{k=2}^p p_{\alpha_n}(\|\boldsymbol{\pi}_k\|_{\mathbf{H}}), \end{aligned} \quad (4.1)$$

where the functions $\hat{g}_l(\cdot)$ are known, p_{α_n} is the SCAD penalty function proposed by Fan and Li (2001), and α_n is the tuning parameter. We find the minimizer of (4.1) by the following steps:

Step 1: Initialize $\boldsymbol{\pi} = \boldsymbol{\pi}^{(1)}$.

4. MODEL SELECTION₁₇

Step 2: Given $\boldsymbol{\pi}^{(m)}$, update $\boldsymbol{\pi}^{(m+1)}$ by minimizing (4.1).

Step 3: Iterate Step 2 until $\|\boldsymbol{\pi}_k^{(m+1)} - \boldsymbol{\pi}_k^{(m)}\|_H < \epsilon_1$.

In the implementation of the algorithm, the initial value $\boldsymbol{\pi}^{(1)}$ is given by the unpenalized estimators. To update $\boldsymbol{\pi}^{(m+1)}$ in Step 2, we apply an iterative process similar to Step 2 in Section 2, but with an additional penalty term. If the m th iterative estimator $\boldsymbol{\pi}^{(m)}$ is close to zero, that is, $\|\boldsymbol{\pi}^{(m)}\|_H$ is smaller than a positive threshold value ϵ_2 , then we set $\hat{\beta}_k(u) = 0$.

We set $\epsilon_1 = \epsilon_2 = 0.01$ in the simulation.

The penalized estimator of $\boldsymbol{\pi}$ is denoted by $\hat{\boldsymbol{\pi}} = \{\hat{\pi}_{sk}, 1 \leq s \leq J_{n,2}, 1 \leq k \leq p\}^T = \arg \min_{\boldsymbol{\pi}} Q(\boldsymbol{\pi})$, and the penalized spline estimators are given by $\hat{\beta}_k^P(u) = \mathbf{B}_2(u)^T \hat{\boldsymbol{\pi}}_k$, $k = 1, \dots, p$. We then have the following results.

Theorem 5. *Under Conditions (C1)–(C7), if $\alpha_n \rightarrow 0$, $J_{n,2}^{-r}/\alpha_n \rightarrow 0$, $J_{n,2}^{1/2}n^{-1/2}/\alpha_n \rightarrow 0$ as $n \rightarrow \infty$, then we have (i) $\hat{\beta}_k^P(u) = 0, k = p_1 + 1, \dots, p$, with probability tending to 1; (ii) $\|\hat{\beta}_k^P(u) - \beta_k(u)\|_{L_2} = O_p(J_{n,2}^{1/2}n^{-1/2} + J_{n,2}^{-r}), k = 1, \dots, p_1$.*

4.2 Hypothesis testing

The model (1.2) is quite general, since it does not depend on any specific parametric structure. However, it is worth considering whether it can be

4. MODEL SELECTION₁₈

reduced to a simpler form. In this subsection, we propose an L_2 -distance statistic to complete the testing problem.

We are interested in testing whether the model (1.2) can degenerate to the single-index model (1.1), that is,

$$H_0 : \beta(u) = \beta, \quad \text{a.s. on } [0, 1].$$

The null hypothesis can be rewritten as $H_0 : \int_0^1 \|\beta(u) - \beta\|^2 du = 0$. Let $\hat{\beta}(u)$ be the spline estimators of the model (1.2) and let $\hat{\beta}$ be the parametric estimator of the model (1.1). We define the L_2 -distance test statistic as

$$\mathcal{T} = \frac{n}{J_{n,2}} \int_0^1 \|\hat{\beta}(u) - \hat{\beta}\|^2 du. \quad (4.2)$$

The following theorem presents the asymptotic behavior of the test statistic under the null hypothesis.

Theorem 6. *Under Conditions (C1)–(C7), if $n^{1/(2r+1)} \ll J_{n,1} \ll n^{1/4}$ and $n^{1/(2r+1)} \ll J_{n,2} \ll n^{1/4}$, then, under H_0 , $\mathcal{T} - B \xrightarrow{d} N(0, V)$, where $B = (n/J_{n,2}) \int_0^1 \Omega_2(u) du$ and $V = (8n^2/J_{n,2}^2) \int_0^1 \int_0^1 \Omega_2(u)\Omega_2(u') du du'$.*

To demonstrate the power of the test, we define a sequence of local alternatives that converge to the null hypothesis with increasing sample size. We consider $H_1 : \beta_n(u) = \beta + c_n \Delta(u)$, where $c_n \rightarrow 0$ as $n \rightarrow \infty$, and $\Delta(u)$ is a continuous function. We show the asymptotic distribution of the test statistic under H_1 in the following theorem.

5. NUMERICAL STUDIES¹⁹

Theorem 7. *Under Conditions (C1)–(C7), if $n^{1/(2r+1)} \ll J_{n,1} \ll n^{1/4}$ and $n^{1/(2r+1)} \ll J_{n,2} \ll n^{1/4}$, and we let $c_n^2 = J_{n,2}/n$, then, under H_1 , $\mathcal{T} - B \xrightarrow{d} N(\tilde{\Delta}, V)$, where $\tilde{\Delta} = \int_0^1 \Delta^2(u) du$.*

Although a plug-in procedure can be used to evaluate the asymptotic distribution of the test statistic, we adopt a wild bootstrap method to improve the performance of the test. The steps can be found in the Supplementary Materials.

5. Numerical Studies

In this section, we conduct simulation studies to illustrate the good performance of the proposed model and the corresponding statistical inference methods for finite samples.

5.1 Selecting the smoothing parameter

We use spline order $q_1 = q_2 = 3$ throughout the numerical simulation studies. For each univariate function, the positions of interior knots are chosen as equally spaced sample quantiles. The smoothing parameters K_1 and K_2 are selected by minimizing the following Bayesian information criterion (BIC): $\text{BIC}(K_1, K_2) = \log \left\{ n^{-1} \sum_{i=1}^n \left[Y_i - \sum_{l=1}^d \hat{g}_l(\mathbf{Z}_i^T \hat{\beta}(U_i)) X_{il} \right]^2 \right\} + df \log(n)/n$, where $df = p(q + K_1) + d(q + K_2)$. That is, the optimal num-

5. NUMERICAL STUDIES₂₀

ber of interior knots is given by $(\hat{K}_1, \hat{K}_2) = \arg \min_{K_1, K_2 \in K_T} \text{BIC}(K_1, K_2)$, where K_T is the potential number of interior knots.

For the model identification procedure, we set $a = 3.7$ in the SCAD function as in Fan and Li (2001). The tuning parameter α_n is chosen by minimizing $\text{BIC}(\alpha_n) = \log \left\{ n^{-1} \sum_{i=1}^n \left[Y_i - \sum_{l=1}^d \hat{g}_l(\mathbf{Z}_i^T \hat{\boldsymbol{\beta}}^P(U_i)) X_{il} \right]^2 \right\} + (p - p_1)(q + K_1) \log(n)/n$. The optimal choice is obtained as $\hat{\alpha}_n = \arg \min \text{BIC}(\alpha_n)$.

5.2 Numerical examples

Example 1. We consider the model:

$$Y_i = g_1(\mathbf{Z}_i^T \boldsymbol{\beta}(U_i)) X_{i1} + g_2(\mathbf{Z}_i^T \boldsymbol{\beta}(U_i)) X_{i2} + \varepsilon_i, \quad i = 1, \dots, n. \quad (5.1)$$

We independently generate covariates X_1 and X_2 from a bivariate normal distribution with mean 0, variance 1, and covariance 0.3. Let $(Z_{i1}^*, Z_{i2}^*, Z_{i3}^*)^T$ follow a multivariate normal distribution with mean 0, variance 1, and covariance 0.2, and then scale it as $Z_{ik} = \Phi(Z_{ik}^*) - 0.5$, where $\Phi(\cdot)$ is the cumulative distribution function of the standard normal distribution. U_i is generated from the uniform distribution $U(0, 1)$. The varying coefficients $\boldsymbol{\beta}(u) = (\beta_1(u), \beta_2(u), \beta_3(u))^T$ are given by $\beta_1(u) = 1 + u^2$, $\beta_2(u) = (1 - u)^2$, and $\beta_3(u) = -1 + 4(u - 0.5)^2$, and they are then identified as $\boldsymbol{\beta}(u)/\|\boldsymbol{\beta}(u)\|_{L_2}$. Set the index functions as $g_1(w) = 10 \exp(5w)/\{1 + \exp(5w)\} - E[10 \exp(5W)/\{1 + \exp(5W)\}]$, $g_2(w) = 5 \sin(\pi w) - E[5 \sin(\pi W)]$.

5. NUMERICAL STUDIES₂₁

The error term ε_i follows an independent normal distribution $N(0, \sigma^2(\mathbf{X}_i, \mathbf{Z}_i, U_i))$ with $\sigma^2(\mathbf{X}_i, \mathbf{Z}_i, U_i) = 0.2 \text{Var}\{E(Y_i | \mathbf{X}_i, \mathbf{Z}_i, U_i)\}$.

To evaluate the performance of the proposed spline estimators, we use the root mean squared error (RMSE) criterion. For any function $f(u)$, the RMSE is given as $\text{RMSE}(f) = \left\{ n^{-1} \sum_{i=1}^n [\hat{f}(u_i) - f(u_i)]^2 \right\}^{1/2}$. Table 1 lists the means and standard deviations (in parentheses) of the RMSE of the estimators for each component function with sample size $n = 300, 500$, or 700 , based on 500 Monte Carlo replications. It is obvious that the mean and standard deviation of RMSE of all estimators decrease as the sample size increases.

Figures 1(a)–1(e) show the estimated functions (dashed), the 95% simultaneous confidence bands (SCBs, dash-dotted), and the pointwise confidence bands (PCBs, dotted) for each estimator with sample size $n = 300$. The estimated functions are close to the true curves (solid), and they all fall into the PCBs and SCBs. We also compare the 95% PCBs based on the asymptotic results and the wild bootstrap method described in Section 4.2. It can be seen from Figure 1 in the Supplementary Materials that for all component functions, these two types of confidence bands are close to each other, which means that the wild bootstrap method performs well.

Example 2. To evaluate the power of the test presented in Section 4.2, we

5. NUMERICAL STUDIES₂₂

Table 1: Means and standard deviations (in parentheses) of the RMSE of the estimators for each component function.

n	β_1	β_2	β_3	g_1	g_2
300	0.0682(0.0313)	0.0711(0.0270)	0.0717(0.0258)	0.4450(0.1076)	0.4923(0.1132)
500	0.0523(0.0222)	0.0509(0.0186)	0.0508(0.0181)	0.3210(0.0698)	0.3546(0.0761)
700	0.0430(0.0167)	0.0397(0.0134)	0.0402(0.0143)	0.2619(0.0546)	0.2872(0.0557)

test whether the varying-coefficient components $\beta(u)$ in the model (5.1) are constant. A sequence of alternative models is specified as $H_1 : \beta(u, \theta) = \mathbf{c} + \theta\beta(u)$, where $\mathbf{c} = (c_k, 1 \leq k \leq 3)^T$, with $c_k = \int_0^1 \beta_k(u) du$ and $\theta = \{0, 0.1, 0.2, 0.3, 0.5, 0.8\}$. Let the index components $\mathbf{g}(\cdot)$ and covariates $\mathbf{Z}$ and $\mathbf{X}$ be the same as in Example 1. Let U be equidistant sequences from 0 to 1. The error term follows the same distribution as in Example 1. When $\theta = 0$, it reduces to the null-hypothesis model.

For different sample sizes and θ , we carry out the test procedure 300 times with sample sizes $n = 300, 500$, and 700 . For each replication, we draw $B = 500$ bootstrap samples. The significance level α is 0.05. We give the simulated powers in Figure 1(f). We can conclude that under the null hypothesis, the estimated powers are close to 0.05, which is consistent with the theoretical result. Furthermore, the estimated powers increase to one as θ increases, indicating that the proposed test statistic works well.

5. NUMERICAL STUDIES²³

Figure 1: (a)–(c) Curves of varying-coefficient functions $\beta_k(\cdot)$, $k = 1, 2, 3$. (d) and (e) Curves of index functions $g_l(\cdot)$, $l = 1, 2$. The true functions are shown by solid curves, the spline estimators by dashed curves, the 95% PCBs by dotted curves, and the 95% SCBs by dash-dotted curves. (f) Simulated powers for different sample sizes.

5.3 Comparison with the Random Forest Approach

Prediction is an important concern in statistical modeling. To investigate the prediction performance of the proposed model and estimation, we compare the coverage probability and width of the prediction confidence intervals with three random forest approaches discussed in Zhang et al. (2020), which are out-of-the-bag prediction intervals (OOB), quantile regression

6. REAL DATA ANALYSIS₂₄

forests (QRF), and split conformal prediction intervals (SC). We generate the data according to the settings in Example 1 and rewrite the model (5.1) as $Y = \mathbf{g}(\tilde{\mathbf{X}}) + \epsilon$, where the predictor $\tilde{\mathbf{X}} = (\tilde{X}_1, \dots, \tilde{X}_{\tilde{p}})^T = (\mathbf{X}^T, \mathbf{Z}^T, U)^T$ with $\tilde{p} = 6$.

We set the nominal level at $(1 - \tilde{\alpha})$, with $\tilde{\alpha} = 0.1$. Let the training sample sizes $n = 200, 300, 500$, and 700 and the test sample sizes $n_0 = 20$. Based on 200 replications, Figures 2(a) and 2(b) present the estimated coverage probability and width of the prediction confidence interval for OOB, QRF, SC, and the proposed model (DSICM), respectively. The triangle in each boxplot is the mean of the coverage probability or width of the prediction confidence interval. As shown in Figure 2(a), the coverage probabilities of OOB, SC, and DSICM are close to the nominal level of 0.9, while the QRF estimators are over-covered. Figure 2(b) shows that the width of the prediction confidence interval of DSICM is shorter than those based on other methods.

6. Real Data Analysis

In this section, we illustrate the proposed model and statistical inference methods via the analysis of two real data applications.

6. REAL DATA ANALYSIS

Figure 2: (a) and (b) show the estimated coverage probability and width of the prediction confidence interval for OOB, QRF, SC, and DSICM in boxplots. Each triangle is the mean of the estimated coverage probability or width of the prediction confidence interval.

6.1 Air pollution data for Guangzhou, China

This dataset consists of 2174 observations between January 1, 2014 and December 31, 2019, after the exclusion of several missing samples. The data are available on the website: <https://www.aqistudy.cn>. The aim of this example is to study the potential relationship between the air quality index (AQI) and predictors of the previous day.

Let Y be the logarithm of the AQI for the next day. The predictors include the daily average levels of pollutants commonly used in the literature, namely, ozone Z_1 ($\mu\text{g}/\text{m}^3$), nitrogen dioxide Z_2 ($\mu\text{g}/\text{m}^3$), fine particles Z_3 ($\mu\text{g}/\text{m}^3$), and inhalable particles Z_4 ($\mu\text{g}/\text{m}^3$), and two environmental fac-

6. REAL DATA ANALYSIS 26

tors, namely, temperature X_2 and humidity X_3 . Owing to the particular location and landform of Guangzhou, its climate is mainly influenced by the East Asian monsoon, and so the interaction effect of air pollutants and environmental factors may vary with wind direction.

To begin, we split the data into three groups according to the angles of wind direction. The angle of the first group is 1° – 120° , that of the second group 120° – 240° , and that of the third group 240° – 360° . We then fit each group by the single-index varying-coefficient model

$$Y = g_1(\mathbf{Z}^T \boldsymbol{\beta}) + g_2(\mathbf{Z}^T \boldsymbol{\beta})X_2 + g_3(\mathbf{Z}^T \boldsymbol{\beta})X_3 + \varepsilon, \quad (6.1)$$

where $g_l(\cdot)$, $l = 1, 2, 3$, are unknown smooth functions, $\mathbf{Z} = (Z_1, Z_2, Z_3, Z_4)^T$, and $\boldsymbol{\beta} = (\beta_1, \beta_2, \beta_3, \beta_4)^T$ is the coefficient vector. The covariates $\mathbf{Z}$ are normalized. Figure 3(a) shows the estimated coefficient vectors $(\hat{\beta}_1, \hat{\beta}_2, \hat{\beta}_3, \hat{\beta}_4)^T$ for three groups: Group 1 (solid lines), Group 2 (long dashed lines), and Group 3 (dashed lines). It is worth noting that the parameters $\boldsymbol{\beta}$ are different in the three groups, and so the interaction pattern may be affected by the wind direction. To further explore the dynamic interaction between the response and covariates, we let $U = \text{Direction}/360$ and consider the following model:

$$Y = g_1(\mathbf{Z}^T \boldsymbol{\beta}(U)) + g_2(\mathbf{Z}^T \boldsymbol{\beta}(U))X_2 + g_3(\mathbf{Z}^T \boldsymbol{\beta}(U))X_3 + \varepsilon, \quad (6.2)$$

6. REAL DATA ANALYSIS 27

where $\beta(\cdot) = (\beta_k(\cdot), 1 \leq k \leq 4)^T$.

Figure 3: The lines connect the estimated coefficient vectors of different groups for two sets of real data: (a) Group 1 (solid lines), Group 2 (long dashed lines), and Group 3 (dashed lines); (b) work-hour (solid lines) and off-hour (dashed lines).

By conducting the variable selection described in Section 4.1, we find that the covariates Z_1 and Z_2 are significant. The optimal parameters are chosen as $(q_1, q_2, K_1, K_2) = (3, 3, 2, 2)$ according to the BIC. We perform the hypothesis test developed in Section 4.2 with 500 wild bootstrap resamples, and find that the test p -value is 0.036. Thus, we reject H_0 [the model (6.1)] at a 0.05 significance level.

Figures 4(a) and 4(b) show the resulting spline estimators (solid curves) and the corresponding 95% simultaneous confidence bands (dashed curves) based on 500 wild bootstrap samples for $\beta_k(u)$, $k = 1, 2$. From Figure 4(a), it can be seen that the interaction effect of ozone increases with increas-

6. REAL DATA ANALYSIS

Figure 4: (a) and (b) Estimators (solid curves) of varying-coefficient functions and 95% simultaneous confidence bands (dashed curves) for $\beta_k(\cdot)$, $k = 1, 2$, respectively. (c)–(e) Spline estimates (solid curves) and 95% simultaneous confidence bands (dashed curves) for index functions $g_l(\cdot)$, $l = 1, 2, 3$. (f) *QQ* plot of residuals.

ing wind direction and that their combination has a positive effect on the response. Figure 4(b) reveals that the interaction relationship between nitrogen dioxide and wind direction has a slowly decreasing trend as the wind direction increases and that their interaction has a positive nonlinear impact on the response. The spline estimators of the index functions $g_l(\cdot)$, $l = 1, 2, 3$, and the corresponding 95% simultaneous confidence bands are depicted in Figures 4(c)–4(e). Figure 4(c) reveals that there is a truly non-

6. REAL DATA ANALYSIS 29

---

linear relationship between the median value $\log(\text{AQI})$ and the predictors. Figures 4(d) and 4(e) show that the modifications due to the temperature and humidity of the previous day are altered by the admixture of other variables. In addition, the QQ plot of the residuals in Figure 4(f) shows that the model (6.2) is a reasonable option for this data set.

6.2 WeChat public accounts data

WeChat has become one of the most popular mobile applications in recent years. The public account is one of the most important and most popular functions of WeChat, providing users with information and services. On the other hand, WeChat public accounts have a mature profit model, which is related to the amount of reading. The reading volume of a public account reflects the number of its active fans and directly affects the income of the official accounts. The purpose of this study is to analyze the association between reading volume and related covariates.

The dataset is obtained from a public account analysis website (<http://top.askci.com>) and contains the information of 624 public accounts in March 2017. The response Y is the logarithm of reading volume. The covariates in the index $\mathbf{Z} = (Z_1, Z_2, Z_3, Z_4, Z_5, Z_6, Z_7)^T$ include the type of public account ($Z_1 = 1$ for enterprises and $Z_1 = 0$ for individuals), the

6. REAL DATA ANALYSIS 30

average number of daily posts (Z_2), the ratio of original posts (Z_3), the proportion of posts with video (Z_4), the average title length of the last ten articles (Z_5), the title punctuation index of the last ten articles (Z_6), and the length of the account name (Z_7). Covariates of interest also include the positive emotion score of titles in the last ten articles (X_2). Here we take $X_1 = 1$ as the intercept term and let $\mathbf{X} = (X_1, X_2)^T$.

This dataset records the proportion of articles posted during six time periods (from 18:00, every four hours is a period). We found that the time of posting articles is concentrated in two periods, 18:00–22:00 and 22:00–02:00. Note that the interaction effects between $\mathbf{X}$ and the index $\mathbf{Z}$ may vary with the time of posting articles. Let U be the proportion of articles posted during 18:00–02:00. We first divide the data into two groups according to U , namely work-hour ($U < 0.5$) and off-hour ($U \geq 0.5$). For each group, we run a regression analysis using the single-index varying coefficient model

$$Y = g_1(\mathbf{Z}^T \boldsymbol{\beta}) + g_2(\mathbf{Z}^T \boldsymbol{\beta})X_2 + \varepsilon, \quad (6.3)$$

where $g(\cdot)$ is an unknown smooth function and $\boldsymbol{\beta} = (\beta_1, \beta_2, \beta_3, \beta_4, \beta_5, \beta_6, \beta_7)^T$ is a loading parameter vector. All predictors are centered and standardized. The lines in Figure 3(b) connect the variant estimated coefficient vectors $(\hat{\beta}_1, \hat{\beta}_2, \hat{\beta}_3, \hat{\beta}_4, \hat{\beta}_5, \hat{\beta}_6, \hat{\beta}_7)$, with solid lines corresponding to the work-hour group and dashed lines to the off-hour group. It is clear that the

6. REAL DATA ANALYSIS 31

proportion of articles posted during 18:00–02:00 does indeed change the interaction patterns, which inspires us to further consider the model $Y = g_1(\mathbf{Z}^T \boldsymbol{\beta}(U)) + g_2(\mathbf{Z}^T \boldsymbol{\beta}(U))X_2 + \varepsilon$.

First, we select significant variables Z_1 , Z_2 , Z_3 , Z_4 , and Z_5 in the index by a variable-selection procedure and choose the optimal parameters $(q_1, q_2, K_1, K_2) = (3, 3, 2, 2)$ according to the BIC. Next, we conduct the hypothesis test from Section 4.2 with 500 wild bootstrap resamples, and the test p -value is 0.008. Thus, we reject H_0 [the model (6.3)] at a 0.05 significance level.

Figure 5 shows the resulting spline estimators (solid curves) and the corresponding 95% simultaneous confidence bands (dashed curves) based on 500 wild bootstrap samples for $\beta_k(u)$, $k = 1, \dots, 5$. Figures 5(a)–5(e) show that the indices of Z_1 , Z_2 , Z_3 , Z_4 , and Z_5 change with U , especially the index of Z_2 . The spline estimators of the index functions $g_l(\cdot)$, $l = 1, 2$, and the corresponding 95% simultaneous confidence bands are depicted in Figures 5(f) and 5(g), respectively. Figure 5(f) shows that there is a truly nonlinear relationship between the median value of the reading volume and the predictors. Figure 5(g) shows that the modification due to the positive emotion score of titles in the last ten articles is altered by the admixture of other variables and that this influence changes from positive to

6. REAL DATA ANALYSIS 32

Figure 5: (a)–(e) Estimators (solid curves) of varying-coefficient functions and 95% simultaneous confidence bands (dashed curves) for $\beta_k(\cdot)$, $k = 1, \dots, 5$. (f) and (g) Spline estimates (solid curves) and 95% simultaneous confidence bands (dashed curves) for index functions $g_l(\cdot)$, $l = 1, 2$. (h) QQ plot of residuals.

negative with a decreasing trend. In addition, the QQ plot of the residuals in Figure 5(h) shows that the model (6.2) is a reasonable option for this data set.

7. Conclusion

The dynamic single-index coefficient model proposed in this paper could be regarded as an extension of many existing nonparametric and semiparametric models. We have come up with a spline-based method to approximate the unknown functions and establish the asymptotic properties of the resulting estimators. We have also proposed a model identification procedure to select significant covariates, and we have developed a hypothesis test to check whether the proposed model could be reduced to a simpler form.

There are at least two directions in which this work could be extended. First, motivated by Ma and Song (2015), we could extend the proposed model to the form $Y = \sum_{l=1}^d g_l(\mathbf{Z}^T \beta_l(U)) X_l + \varepsilon$, where the coefficient functions $\beta_l(\cdot)$ vary across X_l . Second, we could consider a semiparametric model $Y = \sum_{l=1}^d g_l(\mathbf{Z}^T \beta(U) + \mathbf{G}^T \gamma) X_l + \varepsilon$, where the covariate $\mathbf{G} \in \mathbb{R}^q$ and $\gamma = (\gamma_1, \dots, \gamma_q)^T$ is unknown.

Supplementary Materials

The Supplementary Materials include the notation used in this paper, the implementation of the gradient function, the wild bootstrap procedure for hypothesis testing, technical proofs of theorems, an additional numerical example, and two additional analyses of real data.

Acknowledgements

The authors would like to thank the Editor, Associate Editor, and two referees for their helpful comments and suggestions that have substantially improved the paper. You's research was supported in part by the National Natural Science Foundation of China (11971291). Zhou's work was supported by the State Key Program of National Natural Science Foundation of China (71931004 and 92046005) .

References

- Chen, D., Hall, P. and Müller, H. G. (2011). Single and multiple index functional regression models with nonparametric link. *Ann. Stat.* **39**, 1720-1747.
- de Boor, C. (2001). *A Practical Guide to Splines*. Springer, New York.
- Fan, J. and Li, R. (2001). Variable selection via nonconcave penalized likelihood and its oracle properties. *J. Amer. Statist. Assoc.* **96**, 1348-1360.
- Fan, J., Yao, Q. and Cai, Z. (2003). Adaptive varying-coefficient linear models. *J. Roy. Statist. Soc. Ser. B* **65**, 57-80.
- Hastie, T. and Tibshirani, R. (1993). Varying-coefficient models. *J. Roy. Statist. Soc. Ser. B* **55**, 757-796.
- Hu, L., Huang, T. and You, J. (2019). Estimation and identification of a varying-coefficient

REFERENCES³⁵

- additive model for locally stationary processes. *J. Amer. Statist. Assoc.* **114**, 1191-1204.
- Kuruwita, C., Kulasekera, K. and Gallagher, C. (2011). Generalized varying coefficient models with unknown link function. *Biometrika* **98**, 701-710.
- Luo, X., Zhu, L. and Zhu, H. (2016). Single-index varying coefficient model for functional responses. *Biometrics* **72**, 1275-1284.
- Li, J. and Lv, J. (2020). High-dimensional varying index coefficient quantile regression model. *Statist. Sinica*, accepted.
- Liu, X., Cui, Y. and Li, R. (2016). Partial linear varying multi-index coefficient model for integrative gene-environment interactions. *Statist. Sinica* **26**, 1037-1060.
- Ma, S., Carroll, R. J., Liang, H. and Xu, S. (2016). Estimation and inference in generalized additive coefficient models for nonlinear interactions with high-dimensional covariates. *Ann. Stat.* **43**, 2102-2131.
- Ma, S. and Song, P. (2015). Varying index coefficient models. *J. Amer. Statist. Assoc.* **110**, 341-356.
- Ma, S. (2016). Estimation and inference in functional single-index models. *Ann. Inst. Statist. Math.* **68**, 181-208.
- Tang, Q. and Cheng, L. (2008). M-estimation and B-spline approximation for varying coefficient models with longitudinal data. *J. Nonparam. Statist.* **20**, 611-625.
- Wang, L., Li, H. and Huang, J. (2008). Variable selection in nonparametric varying coefficient

REFERENCES³⁶

models for analysis of repeated measurements. *J. Amer. Statist. Assoc.* **103**, 1556-1569.

Xia, Y. and Li, W. (1999). On single-index coefficient regression models. *J. Amer. Statist. Assoc.* **94**, 1275-1285.

Xia, Y., Tong, H, Li, W. and Zhu, L. (2002). An adaptive estimation of dimension reduction space. *J. Roy. Statist. Soc. Ser. B* **64**, 363-410.

Yang, L. (2008). Confidence band for additive regression model. *J. Data Sci.* **6**, 207-217.

Zhang, W., Li, D. and Xia, Y. (2015). Estimation in generalised varying-coefficient models with unspecified link functions. *J. Econometrics* **187**, 238-255.

Zhang, H., Zimmerman, J., Nettleton, D. and Nordman, D. J. (2020). Random forest prediction intervals. *Amer. Statist.* **74**, 392-406.

Zhao, W., Lian, H. and Liang, H. (2017). Quantile regression for the single-index coefficient model. *Bernoulli* **23**, 1997-2027.

School of Statistics and Mathematics, Zhongnan University of Economics and Law, Wuhan, China

E-mails: guanxin1112@163.com

School of Statistics and Management, Shanghai University of Finance and Economics, Shanghai, China

liuhua_sufe@163.com

School of Statistics and Management, Shanghai University of Finance and Economics, Shanghai,

---

REFERENCES³⁷

China, and Shanghai Lixin University of Accounting and Finance, Shanghai, China

johnyou07@163.com

Key Laboratory of Advanced Theory and Application in Statistics and Data Science, MOE, and

Academy of Statistics and Interdisciplinary Sciences, East China Normal University, Shanghai,

China

E-mail: yzhou@amss.ac.cn