The epidemiological situation in Africa

JA CAP

Institut de Léprologie Appliquée, B. P. 11023, Dakar, Senegal

Out of a total of 12,000,000 estimated leprosy patients, nearly 4,000,000 patients, approximately one third are living in Africa, in a population of 460,000,000, or an overall estimated prevalence rate of nearly 10 o/oo (1).

The various estimates made by WHO have not changed considerably over the years: in 1965, the number of cases was estimated at 3,868,000 and in 1970 at 3,509,000.

In 1965, 1,712,132 (44.3%) of the estimated cases were registered for treatment, and 1,326,551 (37.8 %) in 1970. The decrease in registered patients is reportedly due to a high number of patients who were released from control during this period. In 1975, the figures available for 28 countries, revealed a modest increase of the registered cases.

No information is available about the regularity of treatment.

Prevalence pattern

In Africa, leprosy is mainly prevalent in the Western, Eastern and Central parts, while the prevalence rates are low in Northern and Southern Africa.

The WHO figures published in 1979 (2), show that the known prevalence rates vary from 21 o/oo (Mali) to a few hundred known patients in Tunis.

In the endemic areas, the known prevalence rates vary from 2.8 o/oo to 7.4 o/oo with an approximate average of 6 o/oo.

It is however difficult to compare prevalence rates from one country to another. The quality and reliability of the available information differ greatly. In some countries, patients who are released from control are still included in the number of registered patients, while in others, they are excluded.

Leprosy prevalence rates in Africa

Region	Number of countries	Number of registered patients	Prevalence rate (o/oo)
West Africa	15	838,792	7.4
Central Africa	8	227,036	4.7
East Africa	15	328,124	2.8
North Africa	4	33,173	0.6
South Africa	3	16,600	0.6
Total	45	1,443,724	5.7

It may be worth mentioning that in Africa, high prevalence rates are not found along the coast, contrary to the Philippines, where leprosy is often concentrated in the coastal areas.

Although WHO observed a moderate increase in the number of registered cases, the latest updating of the Atlas of Leprosy (3), prepared by Damien Foundation, shows a decrease in the countries for which more or less reliable data are available, except for those countries where leprosy activities just started.

In nine countries (Benin, Burundi, Cameroon, Congo, Upper Volta, Rwanda, Senegal, Tanzania, Togo) the number of registered cases has diminished from 232,923 at the end of 1977 to 210,823 at the end of 1979.

In Upper Volta, the number of patients registered at the end of 1977 was 66,471 (11.1 o/oo), and after the survey of Blanc, it was reduced to 39,144 (6.5 o/oo) at the end of 1979, i.e. a reduction of 41.1 %. (Unfortunately, Blanc's report is not available in Senegal.)

In three other countries, one in West Africa (Ivory Coast), one in East Africa (Malawi), and one in Central Africa (C.A.R.), the number of registered patients decreased from 121,881 (8.6 o/oo) to 107,401 (7.4 o/oo) in one year time.

In Senegal, 28,234 leprosy patients were registered in 1958, shortly after the introduction of Dapsone on a mass scale. It rose to 45,164 in 1964, remaining at a plateau for five-six years, to decrease onwards to reach 17,547 by the end of 1979.

Number of leprosy patients registered in Senegal:

A similar pattern is observed in all countries coordinated by OCCGE and OCEAC, the countries which constituted in the colonial days the French West Africa and French Equatorial Africa. In general the decrease of the prevalence rate started to drop considerably during the late sixties and early seventies, after about 15 years of intensive Dapsone treatment.

Evolution of the prevalence rate in countries coordinated by OCCGE and OCEAC

	Number of registered leprosy patients	Prevalence rate (o/oo)
OCCGE		
1971	457,657	17.3
1974	419,833	15.9
1976	356,039	12.3
OCEAC		
1965	191,901	17.7
1971	126,208	11.7
1976	85,378	6.7

The treatment has certainly played a considerable role in the reduction of the prevalence rate, but other factors must also be taken into consideration. The reported regularity for treatment in the OCCGE-countries varied between 21.5 % and 41.8 % between 1958 and 1963.

In Senegal the regularity for treatment in 1979 was reported to be 56.5 %. Projects in other continents, e.g. India have shown that Dapsone monotherapy, administered at less than 75 % regularity produces a very slow reduction rate in the BI of multibacillary cases.

The reduction of the prevalence rate is not due to a decrease in the incidence or case-detection rate. In countries about which information is available, the number of newly detected cases remains fairly stable over the years and its variations are mainly due to an intensification or to a slackening of the leprosy control activities. In Senegal, e.g. the number of newly detected cases turns around 1,500 per year since 1973 (4, 5).

Number of newly registered cases in Senegal

Year	Number of newly registered cases
1973	1,727
1974	1,629
1975	1,829
1976	1,522
1977	1,311
1978	no information
1979	1,464

The reduction of the prevalence rate is partly due to a steadily increasing number of paucibacillary patients who are released from control. In some countries it approaches or even exceeds the number of newly detected cases. In 1979, 1,130 cases were detected in Senegal and 898 were released from control. In Togo, the number of new cases was 852 in 1980, while 632 were released from control (6).

Another reason influencing the decrease in the prevalence rate, and which gives rise to concern, is the number of patients out of control. In 1975, Languillon reported the disappearance of 1,601 patients, representing 5.2 % of the number of patients registered at the beginning of the year. During 1979, 1,297 or 7.0 % of the registered patients were lost from control. Similar situations were observed in Ethiopia (7), and in Togo 708 (6.1 %) patients disappeared during 1980. Correct information is not always available, because the number of patients who are out of control is often combined with the number of patients who died, or whose diagnosis proved to be incorrect or who were entered twice in the registers, etc.....

Some of those patients may simply have moved to another place in order to avoid stigmatization, and may be newly registered in another place. Some of them may heal spontaneously, others may have died, but during visits to some villages, it appeared that quite a number of them were totally unknown in the village where they had been registered.

Clinical pattern

CLASSIFICATION

Leprosy in Africa is characterized by a low proportion of lepromatous cases, especially in the Western and Central parts.

In the nineteen fifties, the prevalence rate in the hot, humid tropical forest of Zaïre was around 50 o/oo, with only 2-5 % lepromatous cases. Surrounding this area, at an altitude of 800 - 1,000 meter, the prevalence rate decreased to about 20 o/oo, but the proportion of multibacillary cases increased to 10 %. Still more to the Eastern and South Eastern parts, the prevalence rates continue to decline, with a proportional increase of lepromatous cases.

Since the introduction of Dapsone treatment on a large scale, in the control of leprosy, a relatively high proportion of paucibacillary cases have been released from control, while multibacillary cases continue to be treated for life. As a result, there is a proportional increase of the multibacillary cases. In 1973, 9.0 % of the patients registered in Senegal were lepromatous and this percentage has increased up to 20.1 % in 1979 (5).

Evolution of leprosy case-load and proportion of lepromatous cases

Senegal:

Year	Number of registered cases	Proportion of lepromatous cases
1973	34,200	9.0
1974	29,709	10.5
1975	27,800	11.8
1976	23,200	17.7
1977	21,808	18.8
1978	18,638	-
1979	17,547	20.1

Similar patterns are observed in Togo (8), Sierra Leone (9), etc.... and the average proportion of lepromatous cases in West Africa is 14.5 %. A comparable proportion is expected, but the available data show it to be 23.4 %.

 Classification of leprosy in Africa

Region	Number of countries	L	T	I	NC
West Africa	10	14.5	67.6	17.8	0.04
Central Africa	6	23.4	45.1	16.9	14.5
East Africa	8	29.9	47.5	21.6	1.0
North Africa	3	20.2	77.5	2.2	—

These data are however of limited value, because most of them do not refer to the whole country, but only to some local leprosy control projects.

In East Africa the lepromatous proportion is higher (29.9 %) and is more comparable to the Asian situation, at least in Ethiopia. In other East African countries, such as Tanzania, the lepromatous proportion (14.8 %) is similar to the West African situation.

An explanation for the low lepromatous proportion and high prevalence rate is not easy to provide. A relationship between the metabolism of *Mycobacterium leprae* and the pigment metabolism exists. It may also be easier to diagnose early tuberculoid cases in a dark skin than in a light skin, and because of the presence of Trypanosomiasis in the leprosy endemic countries, populations may be more accustomed to regular total surveys.

DISABILITIES

There is very little reliable information about the prevalence of disabilities in leprosy in Africa, but the results of the WHO-LAT surveys show a considerable difference in the frequency of disabilities. Two surveys were carried out in Africa: one in Nigeria (Katsina), and one in Cameroon: the proportion of disabilities of grade 2 to 5 was 23 and 36 %, respectively.

The four surveys in Asia gave the following data: Thailand 65 %, Burma 64 % in Schwebo and 55 % in Myingyang, and 51 % in the Philippines.

Both in Africa and in Asia, fewer female patients showed disabilities.

Disabilities and sex

	Male	Female	Total
Katsina	41.0	21.4	23.4
Cameroon	37.8	32.5	35.6
Thailand	71.4	54.2	64.7
Burma (S)	68.0	58.2	63.8
(M)	59.3	48.9	55.2
Philippines	59.1	41.9	51.2
Argentine	70.9	53.1	63.6

The fact that there are more non-lepromatous cases in Africa, may be one possible explanation. Although disabilities occur earlier in paucibacillary leprosy, their frequency is much less than in lepromatous cases, most of whom develop disabilities in the course of the disease.

DAPSONE RESISTANCE

The importance and magnitude of Dapsone resistance has been emphasized by Pearson in Ethiopia. The problem is very serious in this country, probably and at least partly due to the fact that Dapsone resistant cases are searched for, and because of the locally available facilities for the confirmation of the suspected cases. The yearly incidence rate is claimed to be 3 % per year.

In 1969, Browne reported a case in Nigeria; Pattyn, Rollier reported 5 cases in Morocco in 1975 (10).

The first case in Bamako (Mali) was confirmed by Pattyn and Baquillon in 1979.

After a survey of the bacteriologically positive cases living in the surrounding of the Institut Marchoux, and who had been treated with Dapsone for periods varying from 6 to 31 years, a prevalence rate of 7.6 % was found in 105 examined patients.

The survey was extended to the out-patients treated in the Secteur des Grandes Endemies of Bamako, and a prevalence of 3.98 % was found among 251 patients.

In Senegal, 30 cases of Dapsone resistance have been proven by mouse foot pad inoculation (Pattyn), i.e. nearly 1 % (0.9 %) of the 3,396 registered multibacillary cases.

Because of transport difficulties, all these patients have been sent to the Institut de Léprologie Appliquée at Dakar and represent therefore only a small part of the existing Dapsone resistant cases in the country. Some patients who had been sent for confirmation of clinically suspected Dapsone resistance, and already received Rifampicine or Lamprène, rendering the foot pad inoculation impossible.

It is likely that the proportion of Dapsone resistance in Senegal amounts at 3 - 5 % of the multibacillary cases, or between 100 or 200.

These prevalence rates are of the same order as those observed in the other parts of the World, such as Malaysia: 2.5 % (Meade 1973); Costa Rica: 6.8 % (Peters et al 1976); Israel: 3.7 % (Levy et al 1977).

Conclusions

The estimated prevalence rate for Africa is very high, but relatively little information is available at least on the spot.

Leprosy is mainly concentrated in the Western, Central and Eastern parts and is characterized by a low proportion of lepromatous cases and a relatively low proportion of disabilities.

The occurrence of Dapsone resistance is of the same magnitude as in other parts of the World and if no measures are taken, there is a great risk in producing primary resistant multibacillary and paucibacillary cases.

References

- A Guide to Leprosy Control. W.H.O., 1980.
- 2 WHO Wkly Epidem. Rec.; no. 3, 1979, pp 17-23.
- 3 Atlas of Leprosy. Damien Foundation, Rue Stévin 16, B-1040 Brussels.
- 4 Languillon J. L'Institut Marchoux. Acta Leprologica, no. 24, Janvier-Mars 1966.
- 5 Languillon J. Institut de Leprologie Appliquée de Dakar. Acta Leprologica, 69, Octobre-Décembre 1977.
- 6 Programme National de Lutte Contre la Lèpre - Rapport annuel 1980 - Ministère de la Santé Publique, Togo.
- 7 Cap and Banjaw Mulatu. Leprosy Control in Ethiopia. D.A.H.W., 1977.
- 8 Programme National de Lutte Contre la Lèpre, Service de la lèpre, Bureau National de la Coordination, Lome, Togo.
- 9 Nebout M. La Lutte Contre la Lèpre en Afrique Intertropicale - Fondations Raoul Follereau, 33 Rue de Dantzig, Paris XV.
- 10 Bacquillon, Ferracci, Saint André, Pattyn. Dapsone Resistant Leprosy in a Population of Bamako. Lepr. Rev., 51, 4, December 1980.