The impact of obesity on male fecundity: a Brazilian study

Victor T. Dubeux¹, Thaisa Renovato¹, Ana Clara Esteves¹, Lialyz André¹, Aylana de Oliveira¹, Ivan Araújo Penna¹

¹Division of Infertility and Reproductive Medicine, Department of Mother and Child Health. Fluminense Federal University, Niterói, RJ, Brazil.

ABSTRACT

Introduction: Obesity has become a major problem in most developed countries. Infertility ranks high among the issues brought by excessive weight gain, particularly as it may affect male sexual function. This study investigated a population of Brazilian infertile men in an attempt to establish a relationship between obesity and infertility.

Methods: A total of 153 infertile men had their anthropometric data collected and were divided into groups according to their body mass index and waist circumference measurements. Sperm parameters including sperm count, concentration, morphology, and motility were compared.

Results: Multivariate analysis failed to show a positive correlation between excessive weight gain or increased waist circumference, and sperm alterations in a population of infertile men.

Conclusions: The findings described in this study support the idea that there is no association between obesity and semen alterations in a population of infertile men.

Keywords: Obesity, sperm count, infertility.

INTRODUCTION

The World Health Organization (WHO) defines infertility as the inability to achieve pregnancy within 12 months of regular intercourses for couples in conception. Infertility is estimated to affect the lives of at least 10% of the global population (Cooper *et al.*, 2010). The male factor alone accounts for approximately 25-30% of the cases of infertility (Taylor *et al.*, 2003).

Meanwhile, most developed nations have reported increases in the numbers of overweight and obese men and women in their reproductive years, in addition to more cases of diseases related to obesity such as hypertension, diabetes, and vascular disorders (Hammoud *et al.*, 2008).

Weight gain may be associated to changes in the reproductive system, including hormonal alterations due to the impact on testosterone production (Pasquali *et al.*, 2007; Qin *et al.*, 2007; Pasquali, 2006). However, direct effects on semen parameters are yet to be proven (MacDonald *et al.*, 2007; Jensen *et al.*, 2008).

This study aimed to investigate the effects of excessive weight gain on the semen parameters of a population of Brazilian infertile men.

MATERIALS AND METHODS

All male individuals referred to our department between September 2014 and September 2015 due to complaints of infertility were assessed before joining the study. Study participants gave written consent before joining the study, and their identities were anonymized. Only individuals aged 18 years or older who failed to conceive for at least a year while off contraceptive methods, as per the 2010 WHO definition of infertility (Cooper *et al.*, 2010), were enrolled in the study. Subjects with genital alterations that could lead to subfertility – such as vasectomy, presence of varicocele, impalpable vas deferens, urethral strictures, ectopic or atrophic testis – and individuals on drugs and smokers were excluded from the study.

Anthropometric data were obtained on the same day semen samples were collected and processed. All data were measured twice and a peer made a third measurement whenever significant differences were observed. Height was measured in centimeters with the patients standing with their backs against a wall and no shoes on. Weight was measured in kilograms with the patients wearing only underwear and standing with their feet together on a scale; all patients were weighed on the same scale. The body mass index (BMI) was calculated as weight in kilograms divided by the squared height in meters.

Study participants were divided into groups based on their BMIs. Group 1 – Normal (BMI < 25); Group 2 – Overweight (25.0 < BMI < 30); and Group 3 – Obese (BMI > 30.0)

Waist Circumference (WC) was measured in centimeters at the end of a normal exhalation. The tape measure was positioned at the midpoint between the 12th rib and the iliac crest.

The patients were then divided into two groups, one with WC ≤ 102cm and one with WC > 102cm (cutoff-points for cardiologic risk according to the WHO).

Sterile containers were used to hold the semen samples collected in the morning period by masturbation after three to five days of abstinence. The samples were sent for analysis immediately after collection and were processed at our facilities by the same technician, according to the World Health Organization 2010 criteria for semen examination (volume 1.5 ml; sperm count 15x10⁶/ml; total sperm count 39x10⁶; motility 32% A+B; and morphology 4% Kruger).

Statistical analysis

A sample size of 153 patients was deemed sufficient to identify a Spearman's RHO of at least 0.25 between spermatozoa concentration and BMI, with a 0.95 confidence interval and a statistical power of 0.8.

Continuous variables were described in terms of median values (interquartile range) and categorical variables as counts (ratios). The associations between continuous variables were tested with Spearman's rank correlation coefficient. The associations between categorical variables were tested with Fisher's exact and Chi-square tests.

The Mann-Whitney and Kruskal-Wallis tests were used to assess the distribution of continuous variables among different levels of categorical variables. A significance level of 0.05 was adopted for all statistical tests; the data sets were processed on Stata® version 14.0 (2015. StataCorp. College Station, TX, USA).

RESULTS

Table 1 shows the distribution of patients based on WC and BMI measurements. Since only two patients were underweight, they were deemed normal and added to the group with a normal BMI.

Multifactorial analysis failed to show any impact from BMI or WC measurements upon the studied variables when obese and non-obese individuals were compared. The continuous variables shown in the box plots (Figures 1 and 2) had interquartile ranges and medians with insignificant

Figure 1. Box plot representation of sperm parameters according to WC.

Figure 2. Box plot representation of sperm parameters according to the BMI.

P-values in all categories for BMI and WC comparisons.

In regards to continuous variables, the patient distribution also revealed an absence of association between obesity and sperm parameters (Figures 3 and 4).

This study failed to show a statistically significant P-value in the relationship between obesity and sperm parameters.

DISCUSSION

To our knowledge, this has been the first study carried out in Brazil to look into the relationship between obesity and semen parameters.

There are two ways to assess a man's degree of obe-

sity. One is the BMI, an indicator of whole-body fat distribution calculated from one's height and weight. The BMI is generally used to assess the distribution of fat throughout the body surface, particularly the deposition of fat in peripheral tissues. The other is the WC, which reflects visceral fat deposits directly related to the risk of cardiovascular disease (Hammoud *et al.*, 2008).

The BMI distribution in our population was fairly even, but greater numbers of individuals with a WC $<$ 102 cm were observed. The WC was also considered a valid parameter of obesity in this study, since it has been predominantly used to assess the risk cardiovascular disease.

Several reports on the effects of obesity on infertility

Figure 3. Patient BMI distribution

Figure 4. Patient WC distribution

Table 1. Patient BMI and WC distributions

	Patients	Percentage
Waist Circumference		
<102cm	119	75.32%
>102cm	39	24.68%
Total	158	100%
Body Mass Index		
Normal	55	34.81%
Overweight	56	35.44%
Obese	47	29.75%
Total	158	100%

were published in the last decade (MacDonald *et al.*, 2010; Jensen *et al.*, 2008; Kupka *et al.*, 2011). It has been found that male weight gain might lead to hormonal disorders affecting sex hormone-binding globulin and decreasing testosterone levels. Since free testosterone and follicular stimulating hormone bear direct impact on spermatogenesis, and once both hormones are slightly affected by obesity, it is plausible to think that semen parameters may be altered in this subgroup of patients (MacDonald *et al.*, 2010; Håkonsen *et al.*, 2011; Teerds *et al.*, 2011).

The hormonal status of our patients was not assessed because, as mentioned above, obese men are at a higher risk of presenting these alterations. This analysis aimed to assess the influence of obesity on male fecundity in infertile Brazilian men.

Although some authors have found male overweight to have no impact on sperm motility (Jensen *et al.*, 2004; Aggerholm *et al.*, 2008; Chavarro *et al.*, 2010; Fejes *et al.*, 2006; Pauli *et al.*, 2008), morphology (Jensen *et al.*, 2004; Chavarro *et al.*, 2010; Fejes *et al.*, 2006; Bakos *et al.*, 2011), or semen volume (Qin *et al.*, 2007; Jensen *et al.*, 2004; Fejes *et al.*, 2006; Pauli *et al.*, 2008; Magnusdottir *et al.*, 2005), other studies have described an association with decreased sperm counts (Jensen *et al.*, 2005; Håkonsen *et al.*, 2011; Bakos *et al.*, 2011; Magnusdottir *et al.*, 2011; Braga *et al.*, 2012). The conflicting nature of the data published to date hinders the establishment of a clear correlation between having a high BMI and getting pregnant spontaneously or through assisted reproductive technology.

Another important fact is that some of these studies were performed with volunteers (Qin *et al.*, 2007; Jensen *et al.*, 2004; Aggerholm, 2008; Magnusdottir *et al.*, 2005; Gutorova *et al.*, 2014), while others were carried out in fertility clinics (Hammoud *et al.*, 2008; Chavarro *et al.*, 2010; Fejes *et al.*, 2006; Pauli *et al.*, 2008). This difference may affect the results, since individuals seeking care at a fertility clinic have already tried to conceive without success, while the volunteer group is likely to contain fertile individuals.

The effects of obesity on sperm alterations and male infertility are probably multifactorial. They involve several points, including the hypothalamic-pituitary-gonadal axis and the aromatization of steroids to estrogens in peripheral tissues leading to hyperestrogenic hypogonadotropic hypogonadism (Schneider *et al.*, 1979). Associations with altered function of Sertoli cells and decreased spermatogenesis (Winters *et al.*, 2006) have been described, in addition to increased scrotal temperature caused by inguinal and pelvic fat accumulation (and scrotal fat deposition) resulting in higher scrotal temperature and altered spermatogenesis (Shafik & Olfat, 1981).

Obesity has also been associated with erectile dysfunction and decreased fertility secondary to fewer sexual intercours between obese couples. It has been shown that fertility rates tend to improve after weight loss (Hammoud *et al.*, 2008; Qin *et al.*, 2007).

Environmental factors may also be involved in sperm quality, as obese men exposed to lower temperatures did not show sperm alterations (Gutorova *et al.*, 2014).

The systematic reviews and meta-analyses published on the matter have reported conflicting results (Pasquali, 2006; Sermondade *et al.*, 2013). Most of them challenge the composition of the studied groups, while only a few studies have described favorable outcome measures. Additionally, the BMI may be an imperfect measure of obesity due to the different body compositions seen in the population (Prentice & Jebb, 2001).

Even though the WHO has named sperm count, morphology, and motility as the main parameters used to assess male fertility, other factors such as DNA fragmen-

tation, which may improve after weight loss, may be involved. (Sermondade *et al.*, 2013; Alshahrani *et al.*, 2016).

The individuals enrolled in our study had a history of infertility, which made them a better representation of the population in which a relationship between the BMI and infertility could be established. Hormonal parameters were not analyzed, once this study was not designed to find how obesity might impact sperm parameters in the enrolled population, but rather how spermatogenesis was affected.

CONCLUSIONS

It is yet to be determined whether infertile men should be advised to lose weight in order to improve their sperm parameters. This study failed to show an association between obesity and male fecundity.

CONFLICT OF INTERESTS

No conflict of interest have been declared.

Corresponding author:

Victor Teixeira Dubeux
Division of Infertility and Reproductive Medicine
Department of Mother and Child Health.
Fluminense Federal University, Niterói, RJ, Brazil.
E-mail: vdubeux@globo.com

REFERENCES

- Aggerholm A, Thstrup AM, Toft G, Ramlau-Hansen C, Bonde JP. Is overweight a risk factor for reduced semen quality and altered serum sex hormone profile? *Fertil Steril*. 2008; 90:619-26.
- Alshahrani S, Ahmed AF, Gabr AH, Abal Hassan M, Ahmad G. The impact of body mass index on semen parameters in infertile men. *Andrologia*, 2016 (in Press)
- Bakos HW, Henshaw RC, Mitchell M, Lane M. Paternal body mass index is associated with decreased blastocyst development and reduced birth rates following assisted reproductive technology. *Fertil Steril* 2011; 95: 1700-4.
- Braga DP, Halpern G, Figueira Rde C, Setti AS, Iaconalli A Jr, Borges E Jr. Food intake and social habits in male patients and its relationship to intracytoplasmic sperm injection outcomes. *Fertil Steril* 2012; 97:53-9.
- Chavarro JE, Toth TL, Wright DL, Meeker JD, Hauser R. Body mass index in relation to semen quality, sperm DNA integrity, and serum reproductive hormone levels among men attending an infertility clinic. *Fertil Steril* 2010; 93: 2222-31.
- Cooper TG, Noonan E, von Eckardstein S, Auger J, Baker HW, Behre HM, Haugen TB, Kruger T, Wang C, Mbizvo MT, Vogelzang KM. World Health Organization reference values for human semen characteristics. *Hum Reprod Update*. 2010;16:231-45.
- Fejes I, Koloszar S, Zacacski Z, Daru J, Szöllösi J, Pál A. Effect on body weight on testosterone/estradiol ratio in oligozoospermic patients. *Arch Androl*. 2006; 52: 97-102
- Gutorova NV, Kleshchiov MA, Tipisova EV, Osadchuk LV. Effects of overweight and obesity on the spermogram values and levels of reproductive hormones in the male population of the European north of Russia. *Bull Exp Biol Med*, 2014;157:95-8
- Håkonsen LB, Thulstrup AM, Aggerholm A, Olsen J, Bonde JP, Andersen CY, Bungum M, Ernst EH, Hansen ML, Ernst

- EH, Ramlau-Hansen CH. Does weight loss improve semen quality and reproductive hormones? Results from a cohort of severely obese men. *Reprod Health* 2011; 8:24.
- Hammoud AO, Wilde N, Gibson M, Parks A, Carell DT, Meikle AW. Male obesity and alteration in sperm parameters. *Fertil Steril* 2008; 90:2222-5.
- Jensen TK, Anderson AM, Jørgensen N, Andersen AG, Carlsen E, Petersen JH, Skakkebæk NE. Body mass index in relation to semen quality and reproductive hormones along 1,558 Danish men. *Fertil Steril* 2004;82:863-70.
- Kupka MS, Gnath C, Buehler K, Dahncke W, Kruessel JS. Impact of female and male obesity on IVF/ICSI: results of 700.000 ART-cycles in Germany. *Gynecol Endocrinol*. 2011;27:144-9.
- MacDonald AA, Herbison GP, Showell M, Farquhar CM. The impact of body mass index on semen parameters and reproductive hormones in human males: a systematic review with meta-analysis. *Hum Reprod Update* 2010;16:293-311.
- Magnusdottir EV, Thorsteinsson T, Thorsteinsdottir S, Heilmisdottir M, Olafsdottir K. Persistent organochlorines, sedentary occupation, obesity and human male subfertility. *Hum Reprod* 2005; 20: 208-15.
- Pasquali R. Obesity and androgens: facts and perspectives. *Fertil Steril* 2006; 85:1319-40.
- Pasquali R, Patton L, Gambineri A. Obesity and infertility. *Curr Opin Endocrinol Diabetes Obes*. 2007;14:482-7
- Pauli EM, Legro RS, Demers LM, Kunselman AR, Dodson WC, Lee PA. Diminished paternity and gonadal function with increasing obesity on men. *Fertil Steril*. 2008; 90: 346-51.
- Prentice AM, Jebb SA. Beyond body mass index. *Obes Rev* 2001;2:141-7.
- Qin D, Yuan W, Zhou W, Cui Y, Wu J, Gao E. Do reproductive hormones explain the association between body mass index and semen quality? *Asian J Androl* 2007;9: 827-34.
- Schneider G, Kirschner MA, Berkowitz R, Ertel NH. Increased estrogen production in obese men. *J Clin Endocrinol Metab*. 1979;48:633-8.
- Sermondade N, Faure C, Fezeu L, Shayeb AG, Bonde JP, Jensen TK, Van Wely M, Cao J, Martini AC, Eskandar M, Chavarro JE, Koloszar S, Twigt JM, Ramlau-Hansen CH, Borges E Jr, Lotti F, Steegers-Theunissen RP, Zorn B, Polotsky AJ, La Vignera S, Eskenazi B, Tremellen K, Magnusdottir EV, Fejes I, Hercberg S, Lévy R, Czernichow S. BMI Relation to sperm count: an updated systematic review and collaborative meta-analysis. *Hum Reprod Update*. 2013;19: 221-31.
- Shafik A, Olfat S. Scrotal Lipomatosis. *Br J Urol*. 1981;53:50-4
- Taylor, A. ABC of subfertility: extent of the problem. *Br Med J* 2003; 327:434-6.
- Teerds KG, de Rooij DG, Keijzer J. Functional relationship between obesity and male reproduction: from humans to animal models. *Hum Reprod Update* 2011; 17: 667-83.
- Winters SJ, Wang C, AbdIramahaman E, Hadeed V, Dyky MA, Brufsky A. Inhibin-B levels in healthy young adult men and prepubertal boys: is obesity the cause for the contemporary decline in sperm count because of fewer Sertoli cells? *J Androl* 2006; 27:560-4.