Pairs of Normal Matrices With Property L

Helmut Wielandt¹

A short proof is given, under weaker assumptions, of the following theorem first proved by N. Wiegmann: If the eigenvalues α_i, β_i of two normal $n \times n$ matrices A, B may be numbered in such a way that the eigenvalues $\gamma_i(z)$ of $C(z) = A + zB$ are given by $\gamma_i(z) = \alpha_i + z\beta_i$ (for $i=1, \dots, n$ and all complex values of z), then $AB=BA$.

Two $n \times n$ matrices A, B are said² to have property L if their eigenvalues α_k, β_k ($k=1, \dots, n$) may be ordered in such a way that the eigenvalues of $\alpha A + \beta B$ are given by $\alpha\alpha_k + \beta\beta_k$ for all complex numbers α, β . Though every pair of commuting matrices has property L , the converse is not true.² However, if A and B have property L and are normal, then $AB=BA$ according to a theorem of Wiegmann.³ We wish to show that the assumptions of Wiegmann's theorem may be weakened considerably. We begin by proving the following lemma which has some interest in itself.

LEMMA: Let A and B be arbitrary $n \times n$ matrices. Then the set Z of all points z in the complex plane for which $A + zB$ is normal, is either the whole plane, a straight line, a circle, or it contains, at most, two points. If B is normal, neither the circle case nor the two-point case may enter. If Z is the entire plane then $AB=BA$.

PROOF: $A + zB$ is normal if, and only if,

$$(A + zB)(A^* + \bar{z}B^*) - (A^* + \bar{z}B^*)(A + zB) = 0. \quad (1)$$

If $z = x + iy$, (1) is equivalent to a set of $2n^2$ real linear equations in x, y , and $(x^2 + y^2)$. Each of these $2n^2$ equations defines either the whole plane, a circle, a straight line, or else, at most, one point. The intersection of these $2n^2$ point sets obviously is one of the six types described in the lemma. That all these types may indeed occur is shown by the following matrices $A + zB$:

$$(1+z), \quad \begin{pmatrix} 1 & z \\ z & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & z \\ 1 & 0 \end{pmatrix},$$

$$\begin{pmatrix} 0 & z & 0 \\ 0 & 0 & z \\ z & 0 & z-1 \end{pmatrix}, \quad \begin{pmatrix} 0 & z \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

In case B is normal, the terms of (1) containing $z\bar{z}$ cancel, hence Z is determined by linear equations in x and y only. So Z is neither a circle nor a pair of points.

In case Z is the entire plane (1) holds for every z , hence the coefficient of z vanishes, that is, $BA^* - A^*B = 0$. Since B commutes with A^* , it also commutes with every polynomial $f(A^*)$, hence, with A itself. (If A is normal, then $A = f(A^*)$ for some polynomial f . This well-known fact follows by transforming A into diagonal form by means of a unitary transformation.) The lemma being proved, we turn to the generalization of Wiegmann's theorem.

THEOREM: If the eigenvalues α_k, β_k of two normal matrices $A = (a_{ik}), B = (b_{ik})$ may be ordered in such a way that the eigenvalues $\gamma_k(z)$ of $A + zB$ satisfy the inequalities

$$\sum_{k=1}^n |\gamma_k(z)|^2 \geq \sum_{k=1}^n |\alpha_k + z\beta_k|^2 \quad (2)$$

for some values $z = z_\lambda$ ($\lambda = 1, \dots, l$) which are the vertices of a polygon containing 0 in its interior, then $AB=BA$. (This obviously contains Wiegmann's theorem since if A and B have property L then (2) holds even with equality sign for every z .)

PROOF: According to a theorem of Schur,⁴ we have

$$\sum_{i,k} |a_{ik} + zb_{ik}|^2 \geq \sum_k |\gamma_k(z)|^2 \quad (3)$$

for every z , where the equality holds if, and only if, $A + zB$ is normal. From (3) and (2) we have for $z = z_\lambda$

$$\sum_{i,k} |a_{ik} + zb_{ik}|^2 \geq \sum_k |\alpha_k + z\beta_k|^2 \quad (4)$$

which reduces to

$$\sum_{i,k} (a_{ik}\bar{z}b_{ik} + \bar{a}_{ik}zb_{ik}) - \sum_k (\alpha_k\bar{z}\beta_k + \bar{\alpha}_kz\beta_k) \geq 0 \quad (5)$$

since $\sum |a_{ik}|^2 = \sum |\alpha_k|^2$ and $\sum |b_{ik}|^2 = \sum |\beta_k|^2$ in view of the fact that A and B are normal. Since the left-hand side of (5) is linear and homogeneous in x and y , (5) defines either a half plane containing 0 on its boundary or the left-hand side of (5) vanishes identically. The former case cannot occur, for the half plane contains $z_1, \dots, z_l$, hence it contains 0 in its interior. So in (5), and hence in (4), equality holds for every z . Combining this with (2) and (3) we

¹ American University and University of Tübingen. This paper was prepared under a National Bureau of Standards contract with The American University.

² T. S. Motzkin and O. Taussky, Pairs of matrices with property L , Trans. Am. Math. Soc. **73**, 108-114 (1952).

³ N. Wiegmann, Pairs of normal matrices with property L , Proc. Am. Math. Soc. **4**, 35-36 (1953).

⁴ I. Schur, Über die charakteristischen Wurzeln einer linearen Substitution mit einer Anwendung auf die Theorie der Integralgleichungen, Math. Ann. **66**, 488-510 (1909); Theorem II. The theorem of Schur has recently been used in a related problem by H. Schneider, Theorems on normal matrices, Quart. J. Math. Oxford [2] **3**, 241-249 (1952).

conclude that in (3) equality holds for $z=z_\lambda$, hence $A+z_\lambda B$ is normal. Since $z_1, \dots, z_l$ are not collinear and B is normal, the lemma asserts that $A+zB$ is normal for every z , hence $AB=BA$.

REMARK: The theorem is best possible in the sense that there exist noncommuting normal matrices A, B such that inequality (2) holds for every z of a closed half plane containing 0 on its boundary (and for a fixed suitable ordering of the eigenvalues α_k, β_k). An example is given by

$$A = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & i & 0 \\ 0 & 0 & 0 & -i \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}$$

with the ordering $\alpha_k=1, -1, i, -i; \beta_k=i, -i, 0, 0$. Here we have $\gamma_k(z)=1, -1, (-1-z^2)^{\frac{1}{2}}, -(-1-z^2)^{\frac{1}{2}}$, hence $\sum |\gamma_k(z)|^2 \geq \sum |\alpha_k + z\beta_k|^2$ for every z with $y \geq 0$ (though $\gamma_k(z)=\alpha_k + z\beta_k, k=1, \dots, 4$ is valid for $z=0$ only).

WASHINGTON, April 17, 1953.