Upper Bounds for the Determinant of a Row Stochastic Matrix

Karl Goldberg

Institute for Basic Standards, National Bureau of Standards, Washington, D.C.

(February 16, 1966)

If $A = (a_{ij})$ is a row stochastic matrix of order n then $|\det A| \leq (1 - \sum_{j=1}^n \min_i a_{ij})^{n-1}$. If α is a 1 by n real vector such that every element of αA is nonnegative then $|\det A| \leq s(\alpha)/|g(\alpha)|$, where $s(\alpha)$ is the sum of the elements of α and $g(\alpha)$ is the element of α of greatest absolute value. The conditions for equality are determined in both cases.

Key Words: Determinants, matrices (row stochastic), upper bounds.

It is known that if A is a row stochastic matrix then $|\det A| \leq 1$ with equality if and only if A is a permutation matrix. We will use this result to extend itself.

The result is equivalent to the following statement about nonnegative matrices.¹ Let A be a nonnegative matrix of order n , let $r_A^{(i)}$ denote the sum of the elements in the i th row, let

$$r_A = (r_A^{(1)}, r_A^{(2)}, \dots, r_A^{(n)})$$

and

$$R_A = \text{diag} (r_A^{(1)}, r_A^{(2)}, \dots, r_A^{(n)}).$$

Note that $r_A = 0$ if and only if some row of A is all 0's. Otherwise $R_A^{-1}A$ is row stochastic. Thus $|\det A| \leq r_A$ with equality if and only if either $r_A = 0$ or $A = R_A P$ for some permutation matrix P .

We use the following procedure. Given a row stochastic matrix A , let B be a nonsingular matrix such that $AB \geq 0$ (i.e., AB is nonnegative). Then the above result implies that

$$|\det A| \leq \frac{r_{AB}}{|\det B|}$$

with equality if and only if either $r_{AB} = 0$ or $A = R_{AB} P B^{-1}$. The same procedure with $CA \geq 0$ yields a similar result. Since B (or C) need not be nonnegative we have room for improvement.

From this point on A will denote a row stochastic matrix of order n .

Lower case Greek letters will denote 1 by n real vectors including

$$\epsilon = (1 \ 1 \ \dots \ 1) \text{ and } \delta_i = (0 \ \dots \ 0 \ 1 \ 0 \ \dots \ 0) = (\delta_{ij}).$$

For a vector α , $s(\alpha)$ will denote the sum of its elements, $g(\alpha)$ will denote its element of greatest absolute value, and $M_i(\alpha)$ will denote the matrix

$$M_i(\alpha) = \begin{bmatrix} \delta_1 \\ \cdot \\ \cdot \\ \cdot \\ \delta_{i-1} \\ \alpha \\ \delta_{i+1} \\ \cdot \\ \cdot \\ \cdot \\ \delta_n \end{bmatrix}$$

i.e., the identity matrix with the i th row replaced by α . For future clarity we note now that

$$s(\alpha) = \alpha \epsilon^T,$$

while $\epsilon^T \alpha$ is the matrix

$$\epsilon^T \alpha = \begin{bmatrix} \alpha \\ \alpha \\ \cdot \\ \cdot \\ \cdot \\ \cdot \\ \alpha \end{bmatrix}$$

Finally $A \epsilon^T = \epsilon^T$. We shall prove:

THEOREM 1. Let A be a row stochastic matrix of order n , and let

$$c = \sum_{j=1}^n \min_i a_{ij}.$$

Then

$$|\det A| \leq (1 - c)^{n-1}$$

¹ A matrix is nonnegative if each element is real and nonnegative. It is row stochastic if, in addition, each row sum equals 1.

with equality if and only if

$$A = (1 - s(\beta))P + \epsilon^T \beta$$

for some vector $\beta \geq 0$ with $1 \geq s(\beta)$, and some permutation matrix P .

THEOREM 2. Let A be a row stochastic matrix and α a real, nonzero vector such that $\alpha A \geq 0$. Then

$$|\det A| \leq \frac{s(\alpha)}{|g(\alpha)|}$$

with equality if and only if either $s(\alpha) = 0$, or α has exactly one positive element (in the i -th position) and

$$A = M_i(\gamma)P$$

for some permutation matrix P , with

$$\gamma = \frac{1}{g(\alpha)} \{ (g(\alpha) + s(\alpha))\delta_i - \alpha \}.$$

To prove Theorem 1 let $c_j = \min_i a_{ij}$ and let

$$B = I - \sum_{j=1}^n c_j \epsilon^T \delta_j.$$

The matrix $\epsilon^T \delta_j$ has 1's in the j th column and 0's elsewhere, so that $AB = (a_{ij} - c_j)$. By the definition of the c_j , AB is nonnegative.

The row sums of AB are the elements of the vector

$$AB\epsilon^T = A\epsilon^T - \sum_{j=1}^n c_j \epsilon^T \delta_j \epsilon^T = (1 - c)\epsilon^T.$$

Thus $r_{AB}^{(i)} = 1 - c$ for all i , so that

$$r_{AB} = (1 - c)^n.$$

It is a simple exercise (left to the reader) to prove that

$$\det B = 1 - c$$

and

$$B^{-1} = I + (1 - c)^{-1} \sum_{j=1}^n c_j \epsilon^T \delta_j \quad c \neq 1.$$

If $c \neq 1$ then, from our previous discussion, the necessity case of Theorem 1 follows with

$$\beta = (c_1 \ c_2 \ \dots \ c_n).$$

In the case of sufficiency, if $A = (1 - s(\beta))P + \epsilon^T \beta$ then all the row sums of A equal 1 ($A\epsilon^T = \epsilon^T$), A is nonnegative if $\beta \geq 0$, and $1 \geq s(\beta)$, $\beta = (c_1 c_2 \ \dots \ c_n)$ with the c_j defined as above, and $|\det A| = (1 - s(\beta))^{n-1}$.

A short discussion of the value of c will complete our proof. Since A is row stochastic the sum of all

of its elements is n . Thus

$$n = \sum_{j=1}^n \sum_{i=1}^n a_{ij} \geq \sum_{j=1}^n \sum_{i=1}^n c_j = nc$$

with equality ($c = 1$) if and only if $a_{ij} = c_j$ for all i, j , in which case A is of rank 1. Then $\det A = 1$ for $n = 1$, and $\det A = 0$ with $n > 1$, and $A = \epsilon^T \beta$. This checks with the statement of Theorem 1 and completes our proof.

To prove Theorem 2 let $C = M_i(\alpha)$. The rows of CA are the rows of A except for the i th row which is αA . By hypothesis $\alpha A \geq 0$ so that CA is nonnegative.

The row sums of CA are 1 except for the i th row whose sum is

$$s(\alpha A) = (\alpha A)\epsilon^T = \alpha(A\epsilon^T) = \alpha\epsilon^T = s(\alpha).$$

Thus $r_{CA} = s(\alpha)$ and R_{CA} is the identity matrix transformed by replacing the 1 in the i th position on the diagonal by $s(\alpha)$. Note that, if $s(\alpha) \neq 0$, we have

$$R_{CA}^{-1} M_i(\alpha) = M_i(\alpha/s(\alpha)).$$

The determinant of C is the product of its diagonal elements. Let

$$\alpha = (a_1 a_2 \ \dots \ a_n).$$

Then $\det C = a_i$. By our previous discussion $|\det A| \leq r_{CA}/|\det C|$. Thus

$$|\det A| \leq \frac{s(\alpha)}{|a_i|}.$$

This inequality holds for all i ; the best of the inequalities is

$$|\det A| \leq \frac{s(\alpha)}{|g(\alpha)|}.$$

Suppose $g(\alpha) = a_i$. We have equality if and only if either $s(\alpha) = 0$, (whence $\alpha A = 0$, $\alpha \neq 0$, so that $\det A = 0$) or, if $s(\alpha) \neq 0$:

$$A = C^{-1} R_{CA} P = (R_{CA}^{-1} M_i(\alpha))^{-1} P = (M_i(\alpha/s(\alpha)))^{-1} P$$

for some permutation matrix P .

But $(M_i(\alpha/s(\alpha)))^{-1} = M_i(\gamma)$ with

$$\gamma = a_i^{-1} (-a_i \ \dots \ -a_{i-1} s(\alpha) \ -a_{i+1} \ \dots \ -a_n).$$

Since $A = M_i(\gamma)P$ is nonnegative we must have $a_j \leq 0$ for $j \neq i$. This completes the proof of Theorem 2.

(Paper 70B2-178)