Analysis of Interior-Point Paths

Volume 111

Number 2

March-April 2006

J. Stoer

Institut für Angewandte
Mathematik und Statistik,
Universität Würzburg,
Germany

Infeasible-interior-point paths are the main tools in interior-point methods for solving many kinds of optimization problems. These paths are usually parametrized by a penalty-parameter $r \downarrow 0$ and further parameters describing their off-centrality and infeasibility. Starting with an early result of C. Witzgall et al. [12] in linear programming, this paper gives an overview on results concerning the existence of these paths, their analyticity and the limiting behavior of their derivatives as $r \downarrow 0$, and this also for degenerate problems in the areas of linear programming, linear complementarity problems, and semidefinite programming.

Key words: infeasible interior-point paths; linear complementarity problems; linear programs; semidefinite linear complementarity problems; semidefinite linear programs.

Accepted: December 14, 2005

Available online: <http://www.nist.gov/jres>

1. Introduction

This paper deals with the analysis of certain paths that are the main tool in sequential unconstrained optimization techniques (SUMT) and interior point methods. These were first studied systematically by Fiacco and McCormick in [3] as a means to solve constrained minimization problems of the form

$$\inf\{f(u) \mid g_i(u) \leq 0, i = 1, 2, \dots, m\}. \quad (1)$$

In particular, they used Frisch's logarithmic penalty function (see Frisch [4], [5]) and studied paths $u(r)$, $r > 0$, of local minimizers of the penalized problem

$$\inf\{f(u) - r \sum_i \log(-g_i(u)) \mid g_i(u) < 0, i = 1, 2, \dots, m\}$$

as $r \downarrow 0$. They investigated the local existence and uniqueness of these paths, their differentiability proper-

ties for $r > 0$ and the convergence of both $u(r)$ and $u'(r)$ for $r \downarrow 0$, if $r > 0$ is small and $u(r)$ is close to a local optimal solution u^* of problem (1). They used the crucial assumption that u^* satisfies the standard regularity conditions, as the existence of Lagrange multipliers, the first order necessary and the second order sufficiency condition for a local minimum, and the strict complementarity condition. These assumptions are natural and they allow the use of the implicit function theorem to analyze the path. However they also exclude degenerate problems of practical importance: These occur e.g. already with convex and even linear or quadratic problems (1) that have more than one optimal solution u^* or which, together with their optimal Lagrange multipliers, violate strict complementarity.

Another limitation of paths like $u(r)$ is that they exist only for strictly feasible optimization problems (1). More general are *infeasible-interior-point paths* that use strictly feasible solutions of a family of suitably

perturbed optimization problems that shrink to the unperturbed problem (1) as the path parameter r tends to zero. Again this poses the question of the existence of such paths and their limiting behavior, also for degenerate unperturbed problems.

It is the purpose of this paper to give an overview over some results that are relevant in this context.

In a joint paper [12] with P. T. Boggs and P. D. Domich, C. Witzgall was one of the first to deal also with degenerate problems in the context of linear programs. Their pioneering results will be outlined in Sec. 2. Similar and even sharper results have been found for much more general classes of programs. We describe some of them that refer to linear complementarity problems in Sec. 3. Still more difficult are semidefinite linear programs and semidefinite linear complementarity problems. Results for them are described in the final Sec. 4.

2. Linear Programs

Consider the dual linear programs

$$(P) \quad \min \quad c^T x \quad (D) \quad \max \quad b^T u \\ x: \quad Ax = b, \quad y, u: \quad A^T u + y = c, \\ x \geq 0, \quad y \geq 0,$$

where $A = (a_1, \dots, a_n)$ is a real $m \times n$ -matrix of rank m with columns a_i . Let S_P and S_D be the *optimal solution sets* of (P) and (D).

Consider for (D) the Frisch logarithmic penalty function and its minimum $u(r)$ for $r > 0$,

$$u(r) := \arg \min_u \{-b^T u - r \sum_{i=1}^n \log(c_i - a_i^T u) \mid A^T u < c\}.$$

The path $\{u(r)\}_{r>0}$, if it exists, is called the *dual central path*; then $u = u(r)$ solves the equation

$$b = r \sum_{i=1}^n \frac{a_i}{c_i - a_i^T u} \quad (2)$$

A central result of [12] adapted to (P) and (D) is

Theorem 1 Assume

- (a) rank $A = m$,
- (b) (D) is strictly feasible,
- (c) S_D is nonempty and bounded.

Then the path $u(r)$ is well-defined for $r > 0$ and the following limits exist:

$$u^* := \lim_{r \downarrow 0} u(r) \in S_D, \quad u'(0) := \lim_{r \downarrow 0} \frac{u(r) - u^*}{r}.$$

The limit point u^* is the analytic center of S_D , and the function $u(r)$ is analytic in r for $r > 0$.

Let

$$y(r) := c - A^T u(r), \quad x_i(r) := \frac{r}{y_i(r)}, \quad i = 1, \dots, n.$$

Then $(x, y, u)(r)$, $r > 0$, the *primal-dual central path*, solves

$$(PD)_r \quad \begin{aligned} Ax &= b, \quad x > 0, \\ A^T u + y &= c, \quad y > 0, \\ x \circ y &= re, \end{aligned}$$

where e is the vector

$$e := (1, 1, \dots, 1)^T \in \mathbf{R}^n,$$

and $x \circ y$ is the componentwise product

$$x \circ y := (x_1 y_1, \dots, x_n y_n)^T.$$

By Eq. 2 the primal-dual path exists in the situation of Theorem 1. In general, the primal-dual path is well-defined for $r > 0$, provided (P) and (D) are *strictly feasible*; primal-dual interior point methods try to follow this path for $r \downarrow 0$. But there are some difficulties with this concept:

- (PD)_r is a nonlinear equation for each r , so one can at best approximate points on it.
- The path does not exist, if (P) and (D) are not strictly feasible, even when they are solvable.

These difficulties are avoided if one uses *infeasible (weighted) interior-point paths* instead, defined as the solution $z := (x, y, u) = z(r, \eta) = (x, y, u)(r, \eta)$ of the nonlinear system

$$(PD)_{r,\eta} \quad \begin{aligned} Ax &= b + r \bar{b}, \quad x > 0, \\ A^T u + y &= c + r \bar{c}, \quad y > 0, \\ x \circ y &= r \eta, \end{aligned}$$

where $r > 0$, $0 < \eta \in \mathbf{R}^n$ is a weight vector, and $\bar{b}$, $\bar{c}$ are suitable perturbation vectors. The infeasible interior-point *central path* belongs to the weight vector e .

As initialization one may choose $r_0 := 1$, any $x_0 > 0$, $y_0 > 0$, u_0 , and put

$$\bar{b} := Ax_0 - b, \quad \bar{c} := A^T u_0 + y_0 - c, \quad \eta_0 := x_0 \circ y_0.$$

Then $(x_0, y_0, u_0) = (x, y, u)(r_0, \eta_0)$ solves at least $(PD)_{r,\eta}$ for $r = r_0$ and $\eta = \eta_0$. The choice $x_0 = y_0 := e$ leads to $\eta_0 = e$, that is to a point on the central path.

The practical significance of these paths lies in their use in *infeasible-interior-point methods* (see e.g. Mizuno [10] for an early analysis of such methods in the context of linear programming). Typically, these methods start with an initial point

$$z_0 = z(r_0, \eta_0), \quad r_0 > 0, \quad \eta_0 > 0,$$

on the initial trajectory at (r_0, η_0) defined by the initialization. Then these methods construct further iterates

$$z_k = z(r_k, \eta_k), \quad r_k > 0, \quad 0 < \eta_k \in \mathbf{R}^n, \quad k = 1, 2, \dots,$$

where the $r_k > r_{k+1} > \dots > 0$ converge to 0 and the weights $\eta_k \approx e$ stay bounded and close to e . The construction of z_{k+1} from z_k use, as a rule, the tangent of $z(\cdot, \eta_k)$ at r_k , that is the derivative $z'_k := \partial z(r, \eta_k) / \partial r|_{r=r_k}$ (usually called the *affine scaling direction*). The quality of these methods depends on how quickly the r_k converge to 0. With most methods, the superlinear convergence of the r_k relies on the existence (and boundedness) of the derivatives $z'_k = z'(r_k, \eta_k)$ as $r_k \downarrow 0$.

Also with the paths $z(r, \eta)$ the same questions arise as before:

- their existence and uniqueness for $0 < r \leq r_0 = 1, \eta > 0$,
- their analytical properties,
- and their limiting properties for $r \downarrow 0$.

We will answer these questions in the context of more general complementarity problems in Sec. 3. Here, we only formulate linear programs as particular complementarity problems, which is well-known.

By the duality theorem of linear programming, feasible solutions $x, (y, u)$ of (P) and (D) satisfy

$$0 \leq c^T x - b^T u = (A^T u + y)^T x - (Ax)^T u = y^T x,$$

and they are optimal solutions of (P) and (D) , if they are feasible and $x^T y = 0$.

By elimination of u , define the linear manifold $\mathcal{M} \subset \mathbf{R}^{2n}$,

$$\mathcal{M} := \{(x, y) \mid \exists u : Ax = b, A^T u + y = c\},$$

and the associated linear subspace

$$\Phi := \{(x, y) \mid \exists u : Ax = 0, A^T u + y = 0\}.$$

Then

$$(x, y) \in \Phi \Rightarrow x^T y = 0,$$

and by the assumption $\text{rank}(A) = m$,

$$\dim \Phi = \dim \mathcal{M} = n.$$

The linear programs (P) and (D) are then equivalent to the *associated complementarity problem* of finding a solution (x, y) of

$$(LCP)_{LP} \quad \begin{aligned} (x, y) &\in \mathcal{M}, \\ x &\geq 0, y \geq 0, \\ x \circ y &= 0 \quad (\Leftrightarrow x^T y = 0). \end{aligned}$$

Under the assumptions of Theorem 1, this particular complementarity problem satisfies

- (a) $\dim \Phi = n$
- (b) $x^T y = 0$ for all $(x, y) \in \Phi$
- (c) $(LCP)_{LP}$ is solvable.

3. General Linear Complementarity Problems

A general (LCP) (in horizontal form) is to solve

$$(LCP) \quad \begin{aligned} Px + Qy &= q, \\ x &\geq 0, y \geq 0, \\ x \circ y &= 0 \quad (\Leftrightarrow x^T y = 0), \end{aligned}$$

where $q \in \mathbf{R}^n$ and P, Q are real $n \times n$ -matrices with $\text{rank}[P, Q] = n$. With respect to (LCP) we define:

1. $z = (x, y)$ is a (strictly) *feasible point* of (LCP) , iff $Px + Qy = q$ and $z \geq 0$ (resp. $z > 0$).
2. The *set of solutions* of (LCP) is denoted by S . A solution $(x, y) \in S$ is called *strictly complementary*, if $x + y > 0$.
3. $\Phi \subset \mathbf{R}^{2n}$ is the linear space

$$\Phi := \mathcal{N}[P, Q] = \{(x, y) \mid Px + Qy = 0\}.$$

4. (LCP) , resp. Φ , is called *monotone*, if

$$(x, y) \in \Phi \Rightarrow x^T y \geq 0.$$

The LCP s associated with linear programs are monotone; by Eq. 3 they even satisfy $x^T y = 0$ for all $(x, y) \in \Phi$.

For the presentation of the following results, we make the following rather weak standard assumption

- (V) 1) $\text{rank}[P, Q] = n$, i.e. $\dim \Phi = n$,
 2) (LCP) is monotone,
 3) $S \neq \emptyset$, (LCP) is solvable.

Note that (V) 3) does not require the unique solvability of (LCP). Degenerate problems with more than one solution are not excluded.

Still more general than monotone complementarity problems are *sufficient* linear complementarity problems. They were introduced by Cottle, Pang, and Venkateswaran in [2] and further characterized by Väliäho [11]. They are the most general class of linear complementarity problems of the form (LCP) with the following properties:

1. For all q , the solution set $S = S(q)$ of (LCP) is a convex (perhaps empty) set.
2. Each stationary solution of the (nonconvex) quadratic optimization problem

$$\inf \{x^T y \mid (x, y) \text{ is a feasible point of (LCP)}\}$$

associated with (LCP) is a solution of (LCP).

Again, infeasible (weighted) interior-point paths play an important role in methods for solving (LCP). They are defined as the solutions $(x, y) = (x, y)(r, \eta)$ of

$$(LCP)_{r,\eta} \quad \begin{aligned} Px + Qy &= q + r\bar{q}, (x, y) > 0, \\ x \circ y &= r\eta, \end{aligned}$$

where $r > 0$, $0 < \eta \in R^n$ is a weight, and $\bar{q} \in R^n$ is a perturbation vector.

For initialization, choose any $x_0 > 0$, $y_0 > 0$, $r_0 > 0$, and put

$$\bar{q} := (Px_0 + Qy_0 - q)/r_0, \quad \eta_0 := x_0 \circ y_0/r_0.$$

Then at least $(x_0, y_0) = (x, y)(r_0, \eta_0)$, and

$$\bar{q} \in M_{r_0} := \{(Px + Qy - q)/r_0 \mid x > 0, y > 0\}.$$

Interior-point methods for solving (LCP) try to follow such paths for $r \downarrow 0$ in much the same way as described for linear programs in Sec. 2.

As before, the same questions with respect to these paths $(x, y)(r, \eta)$ arise,

- their existence and uniqueness for $0 < r \leq r_0 = 1$, $\eta > 0$,

- their analytical properties as functions of r, η ,
- and their limiting properties for $r \downarrow 0$.

The following theorem (see Stoer and Wechs [9]) can be viewed as a generalization of Theorem 1, the result of Witzgall et al.:

Theorem 2 Assume (V), $r_0 > 0$, and $\bar{q} \in M_{r_0}$. Then the following holds:

- a) The path $z(r, \eta) := (x, y)(r, \eta)$ is a well-defined analytic function of (r, η) for all $(r, \eta) \in (0, r_0) \times R_{++}^n$, which converges for $r \downarrow 0$ to a solution of (LCP):

$$\exists z^*(\eta) = (x^*, y^*)(\eta) := \lim_{r \downarrow 0} (x, y)(r, \eta) \in S.$$

- b) If, in addition, (LCP) has a strictly complementary solution, then there is an $\varepsilon > 0$, so that the function $z(r, \eta)$ can be extended to an analytic function $\hat{z}(r, \eta)$ on the enlarged domain of all $(r, \eta) \in [-\varepsilon, r_0] \times R_{++}^n$.
- c) If (LCP) is solvable but has no strictly complementary solution, then there is an $\varepsilon > 0$, and an analytic function $\hat{z}(\rho, \eta)$ on the domain $(\rho, \eta) \in [-\varepsilon, \sqrt{r_0}] \times R_{++}^n$ so that $z(\rho^2, \eta) = \hat{z}(\rho, \eta)$ for all

$$(\rho, \eta) \in (0, \sqrt{r_0}] \times R_{++}^n.$$

(“(x, y)(r, η) is an analytic function of $\rho = \sqrt{r}$ even in $r = 0$.”)

3.1 Remarks

1. By a result of Goldman and Tucker [6], any solvable complementarity problem associated with linear programs has also a strictly complementary solution. So, part b) of the theorem applies in this case, which generalizes Theorem 1: the limits

$$z^{*(k)}(\eta) := \lim_{r \downarrow 0} \frac{\partial^k}{\partial r^k} z(r, \eta), \quad k = 0, 1, \dots,$$

of all derivatives exist, and are in fact the derivatives of an analytic function $\hat{z}(r, \eta)$ at $r = 0$ extending $z(r, \eta)$.

2. It is easy to find convex quadratic programs for which the corresponding complementarity problem is only solvable but has no strictly complementary solution, so case c) of the theorem can occur.
3. Theorem 2 remains true for sufficient complementarity problems (i.e. if “monotone” is replaced by “sufficient” in Assumption (V) (see [8])).

The proof of Theorem 2 is not easy in the case of degeneracies, when (LCP) has more than one solution or no strictly complementary solution. The mathematical reason is that the solutions (x, y, r, η) of $(LCP)_{r,\eta}$ with $x \geq 0, y \geq 0, r \geq 0$ and $\eta > 0$ then form a semialgebraic variety with a singularity in $r = 0$, so that the implicit function theorem cannot be applied directly (though indirectly, see [9], [8] for details).

4. Semidefinite Linear Programs and Complementarity Problems

In this section we show that the results of Theorems 1 and 2 even generalize to semidefinite linear programs and to semidefinite linear complementarity problems.

We use the following notation, which is standard:

$S^n := \{A \in \mathbf{R}^{n \times n} \mid A = A^T\}$ is the space of all real symmetric $n \times n$ -matrices.

S^n is provided with the scalar product $\langle A, B \rangle := \text{tr}(B^T A) = \text{tr}(BA)$.

$A \succeq 0$ (resp. $A \succ 0$) means that $A \in S^n$ is positive semidefinite (resp. positive definite), and S^n_{++} is the set of all positive definite matrices in S^n .

Then the standard dual pair of semidefinite linear programs has the form

$$(P) \quad \inf \quad \langle C, X \rangle \\ X: \quad \langle A_i, X \rangle = b_i, \quad i = 1, 2, \dots, m, \\ X \succeq 0.$$

$$(D) \quad \sup_{u, Y:} \quad b^T u \\ \sum_{i=1}^m u_i A_i + Y = C, \\ Y \succeq 0.$$

Here,

$$C, A_i \in S^n \text{ for } i = 1, \dots, m, \quad b = (b_1, \dots, b_m)^T \in \mathbf{R}^m.$$

The following results are known from the duality theory of semidefinite linear programs:

If (P) and (D) have strictly feasible solutions then they also have optimal solutions, and X and (u, Y) are optimal for (P) and (D) , respectively, iff

$$\begin{aligned} \langle A_i, X \rangle &= b_i, \quad i = 1, 2, \dots, m, \\ \sum_{i=1}^m u_i A_i + Y &= C, \\ XY &= 0, \quad X \succeq 0, Y \succeq 0. \end{aligned}$$

By elimination of u one can formulate this as a particular semidefinite linear complementarity problem, which in general has the following form: Solve

$$(SDLCP) \quad \begin{aligned} P(X) + Q(Y) &= q, \\ XY &= 0, \quad X, Y \succeq 0. \end{aligned}$$

Here, $q \in \mathbf{R}^{\bar{n}}$, $\bar{n} := n(n+1)/2 = \dim S^n$, and $P, Q: S^n \rightarrow \mathbf{R}^{\bar{n}}$ are linear operators of the form

$$\begin{aligned} P(X) &= (\langle P_1, X \rangle, \dots, \langle P_{\bar{n}}, X \rangle)^T, \quad P_i \in S^n, \\ Q(X) &= (\langle Q_1, X \rangle, \dots, \langle Q_{\bar{n}}, X \rangle)^T, \quad Q_i \in S^n. \end{aligned}$$

As in the previous section we define:

1. The nullspace $\mathcal{N}([P, Q])$ of $[P, Q]$ is denoted by

$$\Phi := \{(X, Y) \mid P(X) + Q(Y) = 0, X, Y \in S^n\}.$$

2. $(SDLCP)$, resp. Φ , is called *monotone*, if

$$(X, Y) \in \Phi \Rightarrow \langle X, Y \rangle \geq 0.$$

3. The set $\mathcal{F}$ of *feasible points* of $(SDLCP)$ is defined by

$$\mathcal{F} := \{(X, Y) \mid P(X) + Q(Y) = q, X, Y \succeq 0\}.$$

4. The set of *solutions* of $SDLCP$ is denoted by

$$S := \{(X, Y) \in \mathcal{F} \mid XY = 0\}.$$

5. A solution $(X, Y) \in S$ is called *strictly complementary* if $X + Y \succ 0$.

In the case of the dual semidefinite linear programs (P) , (D) , Φ is given by

$$\begin{aligned} \Phi &= \{(X, Y) \mid \langle A_i, X \rangle = 0, \text{ for } i = 1, \dots, m, Y = \\ &\quad - \sum_{i=1}^m u_i A_i \text{ for some } u\}, \end{aligned}$$

and Φ even satisfies

$$(X, Y) \in \Phi \Rightarrow \langle X, Y \rangle = 0,$$

so the corresponding (SDLCP) is monotone. It is not known whether in this case the corresponding (SDLCP) always has a strictly complementary solution if it is solvable.

The following standard hypothesis, which corresponds to the standard assumption of the previous section, is very weak

- (V) 1) $\{P(X) + Q(Y) \mid X, Y \in \mathcal{S}^n\} = \mathbf{R}^n$,
 2) (SDLCP) is monotone,
 3) $\mathcal{S} \neq \emptyset$, (SDLCP) is solvable.

The definition of infeasible weighted interior-point paths differs slightly from the previous section: For $0 < r \leq r_0$ and weight matrices $M \succ 0$, infeasible weighted interior point paths are now defined as solutions $(X, Y) = (X, Y)(r, M)$ of

$$\begin{aligned} P(X) + Q(Y) &= q + r\bar{q}, \quad X \succ 0, Y \succ 0 \\ (XY + YX)/2 &= rM. \end{aligned}$$

Here, the condition $XY = 0$ of (SDLCP) is weakened to $(XY + YX)/2 = rM$, which corresponds to the “AHO” approach proposed by Alizadeh, Haeberly and Overton in [1].

The perturbation vector $\bar{q}$ is chosen such that $\bar{q} \in \mathcal{M}_{r_0}$ for some $r_0 > 0$, where

$$\mathcal{M}_{r_0} := \left\{ \frac{P(X) + Q(Y) - q}{r} \mid X, Y, XY + YX \succ 0 \right\}.$$

A possible initialization is to choose $r_0 := 1$, $X_0 = Y_0 = I$ and $M_0 = I$ and $\bar{q} := P(X_0) + Q(Y_0) - q$.

Again the same questions arise for these paths $(X, Y)(r, M)$ as in the previous sections, namely their

- existence and uniqueness for $0 < r \leq r_0$, $M \succ 0$,
- analytical properties as functions of r and M ,
- and their limiting properties for $r \downarrow 0$.

The following is the main result on (SDLCP). It corresponds to Theorems 1 and 2, and shows the fruitfulness of the early results of C. Witzgall:

Theorem 3 Assume (V), $r_0 > 0$ and $\bar{q} \in \mathcal{M}_{r_0}$. Then:

1. The path function $(X, Y)(r, M)$ is a well-defined analytic function for all $(r, M) \in (0, r_0] \times \mathcal{S}_{++}^n$, which converges to a solution of (SDLCP) as $r \downarrow 0$,

$$\exists (X^*, Y^*)(M) := \lim_{r \downarrow 0} (X, Y)(r, M) \in \mathcal{S}.$$

2. If (SDLCP) even has a strictly complementary solution, then there is an $\varepsilon > 0$ so that the path function $(X, Y)(r, M)$ can be extended to an analytic function $(\hat{X}, \hat{Y})(r, M)$ defined on the enlarged domain $(r, M) \in (-\varepsilon, r_0] \times \mathcal{S}_{++}^n$.
 “ $(X, Y)(r, M)$ is analytic in (r, M) even at $r = 0$.”

The proof of this theorem is even more involved than that of Theorem 2, it can be found in Preiss and Stoer [7].

It is an open problem, whether (and how) the path function $(X, Y)(r, M)$ can be extended, if (SDLCP) satisfies only (V), but has no strictly complementary solutions.

5. References

- [1] F. Alizadeh, J.-P. Haeberly, and M. Overton, Primal-Dual Interior-Point Methods for Semidefinite Programming: Convergence Rates, Stability and Numerical results, SIAM J. Optim. **8**, 746-768 (1998).
- [2] R. W. Cottle, J.-S. Pang, and V. Venkateswaran, Sufficient Matrices and the Linear Complementarity Problem, Linear Algebra Appl. **114/115**, 231-249 (1989).
- [3] A. V. Fiacco and G. P. McCormick, Nonlinear Programming: Sequential Unconstrained Minimization Techniques, John Wiley and Sons, Inc. (1968).
- [4] K. R. Frisch, Principles of Linear Programming—With Particular References to the Double Gradient Form of the Logarithmic Potential Method, Memorandum of October 18, 1954, University Institute of Economics, Oslo.
- [5] K. R. Frisch, The Logarithmic Potential Method of Convex Programming, Memorandum of May 13, 1955, University Institute of Economics, Oslo.
- [6] A. J. Goldman and A. W. Tucker, Theory of Linear Programming, in Annals of Mathematical Studies **38**, Linear Inequalities and Related Systems, H. W. Kuhn and A. W. Tucker, eds., North Holland, Amsterdam (1956) pp. 53-97.
- [7] M. Preiß and J. Stoer, Analysis of Infeasible-Interior-Point Paths Arising with Semidefinite Linear Complementarity Problems, Math. Programming Ser. A **99**, 499-520 (2004).
- [8] J. Stoer and M. Wechs, Infeasible-Interior-Point Paths for sufficient Linear Complementarity Problems and Analyticity, Math. Programming **83**, 407-423 (1998).
- [9] J. Stoer and M. Wechs, On the Analyticity Properties of Infeasible-Interior-Point Paths for Monotone Linear Complementarity Problems, Numer. Math. **81**, 631-645 (1999).
- [10] S. Mizuno, Polynomiality of Infeasible-Interior-Point Algorithms for Linear Programming, Mathematical Programming **67**, 109-119 (1994).
- [11] H. Väliäho, P_* -Matrices are just Sufficient, Linear Algebra Appl. **239**, 103-108 (1996).
- [12] C. Witzgall, P. T. Boggs, and P. D. Domich, On the Convergence Behavior of Trajectories for Linear Programming, in Contemporary Mathematics **114**, Mathematical Developments Arising from Linear Programming Algorithms, J. C. Lagarias and M. J. Todd, eds., American Mathematical Society, Providence, RI (1991) pp. 161-187.

***About the author:** Josef Stoer is a Professor Emeritus at the University of Würzburg, Institute of Applied Mathematics and Statistics.*