ALL MAXIMAL IDEMPOTENT SUBMONOIDS OF GENERALIZED COHYPERSUBSTITUTIONS OF TYPE $\tau = (2)$

NAGORNCHAT CHANSURIYA

*Faculty of Science, Energy and Environment
King Mongkut's University of Technology North Bangkok (Rayong Campus)
Rayong 21120, Thailand*

e-mail: nagornchat.c@sciee.kmutnb.ac.th

Abstract

A generalized cohypersubstitution of type τ is a mapping σ which maps every n_i -ary cooperation symbol f_i to the cotermin $\sigma(f)$ of type $\tau = (n_i)_{i \in I}$. Denote by $\text{Cohyp}_G(\tau)$ the set of all generalized cohypersubstitutions of type τ . Define the binary operation $\circ_{CG}$ on $\text{Cohyp}_G(\tau)$ by $\sigma_1 \circ_{CG} \sigma_2 := \hat{\sigma}_1 \circ \sigma_2$ for all $\sigma_1, \sigma_2 \in \text{Cohyp}_G(\tau)$ and $\sigma_{id}(f_i) := f_i$ for all $i \in I$. Then $\text{Cohyp}_G(\tau) := \{\text{Cohyp}_G(\tau), \circ_{CG}, \sigma_{id}\}$ is a monoid. In [5], the monoid $\text{Cohyp}_G(2)$ was studied. They characterized and presented the idempotent and regular elements of this monoid. In this present paper, we consider the set of all idempotent elements of the monoid $\text{Cohyp}_G(2)$ and determine all maximal idempotent submonoids of this monoid.

Keywords: generalized cohypersubstitutions, idempotent submonoids, maximal submonoids.

2010 Mathematics Subject Classification: 20B10, 20M05, 20M10.

REFERENCES

- [1] K. Denecke and K. Saengsura, *Separation of clones of cooperations by cohyperi-
den-
tities*, Discrete Math. **309** (2009) 772–783.
doi:10.1016/j.dise.2008.01.043
- [2] K. Denecke and S.L. Wismath, *Universal Algebra and Coalgebra* (World Scientific
Publishing Co. Pte. Ltd., Singapore, 2009).
- [3] S. Jermjitpornchai and N. Saengsura, *Generalized Cohypersubstitutions of Type
 $\tau = (n_i)_{i \in I}$* , Internat. J. Pure and Appl. Math. **86** (2013) 745–755.
doi:10.12732/ijpam.v86i4.12

- [4] P. Kunama and S. Leeratanavelee, *All Maximal Completely Regular submonoids of $Hyp_G(2)$* , Discuss. Math. Gen. Alg. and Appl. **37** (2017) 105–114.
doi:10.7151/dmgaa.1263
- [5] N. Saengsura and S. Jermjitpornchai, *Idempotent and Regular Generalized Cohyper-substitutions of Type $\tau = (2)$* , Internat. J. Pure and Appl. Math. **86** (2013) 757–766.
doi:10.12732/ijpam.v86i4.13
- [6] W. Wongpinit and S. Leeratanavelee, *All Maximal idempotent submonoids of $Hyp_G(2)$* , Acta Univ. Sapientiae Math. **7** (2015) 106–113.
doi:10.1515/ausm-2015-0007
- [7] W. Wongpinit and S. Leeratanavelee, *All Maximal idempotent submonoids of $Hyp_G(n)$* , Surveys in Mathematics and Its Applications **10** (2015) 41–48.

Received 4 December 2019

Revised 6 May 2020

Accepted 6 May 2020