GRAPHS WITH 4-RAINBOW INDEX 3 AND $n - 1$

XUELIANG LI¹, INGO SCHIERMEYER²

KANG YANG¹ AND YAN ZHAO¹

¹ Center for Combinatorics and LPMC-TJKLC
Nankai University
Tianjin 300071, China

² Institut für Diskrete Mathematik und Algebra
Technische Universität Bergakademie Freiberg
09596 Freiberg, Germany

e-mail: lxl@nankai.edu.cn
Ingo.Schiermeyer@tu-freiberg.de
yangkang@mail.nankai.edu.cn
zhaoyan2010@mail.nankai.edu.cn

Abstract

Let G be a nontrivial connected graph with an edge-coloring $c : E(G) \rightarrow \{1, 2, \dots, q\}$, $q \in \mathbb{N}$, where adjacent edges may be colored the same. A tree T in G is called a *rainbow tree* if no two edges of T receive the same color. For a vertex set $S \subseteq V(G)$, a tree that connects S in G is called an *S -tree*. The minimum number of colors that are needed in an edge-coloring of G such that there is a rainbow S -tree for every set S of k vertices of $V(G)$ is called the *k -rainbow index* of G , denoted by $rx_k(G)$. Notice that a lower bound and an upper bound of the k -rainbow index of a graph with order n is $k - 1$ and $n - 1$, respectively. Chartrand *et al.* got that the k -rainbow index of a tree with order n is $n - 1$ and the k -rainbow index of a unicyclic graph with order n is $n - 1$ or $n - 2$. Li and Sun raised the open problem of characterizing the graphs of order n with $rx_k(G) = n - 1$ for $k \geq 3$. In early papers we characterized the graphs of order n with 3-rainbow index 2 and $n - 1$. In this paper, we focus on $k = 4$, and characterize the graphs of order n with 4-rainbow index 3 and $n - 1$, respectively.

Keywords: rainbow S -tree, k -rainbow index.

2010 Mathematics Subject Classification: 05C05, 05C15, 05C75.

REFERENCES

- [1] J.A. Bondy and U.S.R. Murty, Graph Theory (GTM 244, Springer, 2008).
- [2] Q. Cai, X. Li and J. Song, *Solutions to conjectures on the (k, ℓ) -rainbow index of complete graphs*, Networks **62** (2013) 220–224.
doi:10.1002/net.21513
- [3] Y. Caro, A. Lev, Y. Roditty, Zs. Tuza and R. Yuster, *On rainbow connection*, Electron. J. Combin. **15** (2008) R57.
- [4] G. Chartrand, G.L. Johns, K.A. McKeon and P. Zhang, *Rainbow connection in graphs*, Math. Bohem. **133** (2008) 85–98.
- [5] G. Chartrand, G.L. Johns, K.A. McKeon and P. Zhang, *The rainbow connectivity of a graph*, Networks **54** (2009) 75–81.
doi:10.1002/net.20296
- [6] G. Chartrand, S.F. Kappor, L. Lesniak and D.R. Lick, *Generalized connectivity in graphs*, Bull. Bombay Math. Colloq **2** (1984) 1–6.
- [7] G. Chartrand, F. Okamoto and P. Zhang, *Rainbow trees in graphs and generalized connectivity*, Networks **55** (2010) 360–367.
doi:10.1002/net.20399
- [8] L. Chen, X. Li, K. Yang and Y. Zhao, *The 3-rainbow index of a graph*, Discuss. Math. Graph Theory **35** (2015) 81–94.
doi:10.7151/dmgt.1780
- [9] P. Erdős and A. Gyárfás, *A variant of the classical Ramsey problem*, Combinatorica **17** (1997) 459–467.
doi:10.1007/BF01195000
- [10] X. Li and Y. Sun, Rainbow Connections of Graphs (Springer Briefs in Math., Springer, New York, 2012).
- [11] X. Li, Y. Shi and Y. Sun, *Rainbow connections of graphs: A survey*, Graphs Combin. **29** (2013) 1–38.
doi:10.1007/s00373-012-1243-2
- [12] X. Li, I. Schiermeyer, K. Yang and Y. Zhao, *Graphs with 3-rainbow index $n - 1$ and $n - 2$* , Discuss. Math. Graph Theory **35** (2015) 105–120.
doi:10.7151/dmgt.1783

Received 14 January 2014

Revised 22 May 2014

Accepted 16 June 2014