NOTE

A NOTE ON TOTAL GRAPHS

S.F. FOROUHANDEH¹, N. JAFARI RAD¹

B.H. VAQARI MOTLAGH¹, H.P. PATIL²

AND

R. PANDIYA RAJ²

¹ *Department of Mathematics*
Shahrood University of Technology
Shahrood, Iran

² *Department of Mathematics*
Pondicherry Central University
Puducherry - India

e-mail: n.jafarirad@gmail.com
hpppondy@gmail.com

Abstract

Erratum: Identification and corrections of the existing mistakes in the paper *On the total graph of Mycielski graphs, central graphs and their covering numbers*, Discuss. Math. Graph Theory **33** (2013) 361–371.

Keywords: total graph, central graph, middle graph, Mycielski graph.

2010 Mathematics Subject Classification: 05C76, 05C69.

1. RESULTS

In this paper, we correct the Theorems 1, 4, 8 and 11, and their corollaries of [1]. There was omitted $t(G)$, i.e., the number of triangles in G or $L(G)$ in Theorem 1 of [1]. The total graph $T(G)$ contains triangles in G , $L(G)$ and in the incidence graph. All triangles are numbered in the published paper [1] beside triangles in G or $L(G)$. First, we give corrected version of Theorem 1 of [1] as follows by adding the number of omitted triangles $t(G)$, and its proof is in similar lines as before.

Theorem 1. For any (p, q) graph G ,

$$t[T(G)] = 2t(G) + \frac{1}{2} \sum_{i=1}^p \left[d_G^2(v_i) + 2m_i \binom{d_G(v_i)}{3} \right],$$

where $m_i = 1$ if $d_G(v_i) \geq 3$; otherwise $m_i = 0$.

Due to the change in the statement of Theorem 1 of [1], the remaining Theorems 4, 8, 11 and their corollaries of [1] are corrected as follows.

Corollary 2. (a) $t[T(C_3)] = 8$ and $t[T(C_n)] = 2n$ if $n > 3$.

(b) For $n \geq 1$, $t[T(K_n)] = \frac{1}{6} [(n^2 - n)(n^2 - 1)]$.

Corollary 3. For $1 \leq i \leq n$ and $n \geq 2$, $t[T(\square_{i=1}^n C_{m_i})] = \frac{2Mn}{3}(2n^2 + 1)$ where $M = m_1 m_2 \cdots m_n$, $m_i > 3$.

Theorem 4. Let G be any (p, q) -graph having $t(G)$ triangles and $\delta(G) \geq 2$. Then

$$t[T(\mu(G))] = 8t(G) + \frac{1}{2} \sum_{i=1}^p [3d_G^3(v_i) + d_G^2(v_i)] + \left(\frac{18q + 5p + p^3}{6} \right).$$

Corollary 5. For $n > 3$, $t(T[\mu(C_n)]) = \left(\frac{n^3 + 107n}{6} \right)$.

Corollary 6. For $n \geq 3$, $t(T[\mu(K_n)]) = \frac{1}{6}(9n^4 - 15n^3 + 6n^2 + 6n)$.

Theorem 7. For any (p, q) -graph G ,

$$t[M(G)] = t(G) + \frac{1}{2} \sum_{i=1}^p \left[d_G^2(v_i) + 2m_i \binom{d_G(v_i)}{3} \right] - q,$$

where $m_i = 1$ if $d_G(v_i) \geq 3$; otherwise $m_i = 0$.

Corollary 8. For any (p, q) -graph G ,

$$t[M(\mu(G))] = 4t(G) + \frac{1}{2} \sum_{i=1}^p [3d_G^3(v_i) + d_G^2(v_i)] + \frac{p(p^2 - 1)}{6}.$$

Theorem 9. For any (p, q) -graph G with $p \geq 4$,

$$t[T(C(G))] = 2m + \frac{1}{6}(p^4 - 3p^3 + 5p^2 - 3p + 12q),$$

where $m = t(C(G))$.

Corollary 10. For $m, n \geq 3$,

$$t[T(C(K_{m,n}))] = t[T(K_{m+n})] - mn(m + n - 4).$$

Acknowledgement

The authors gratefully acknowledge the referee for his valuable suggestions.

REFERENCES

- [1] H.P. Patil and R. Pandiya Raj, *On the total graph of Mycielski graphs, central graphs and their covering numbers*, Discuss. Math. Graph Theory **33** (2013) 361–71.
doi:10.7151/dmgt.1670

Received 3 May 2014

Revised 4 July 2014

Accepted 3 September 2014