CHROMATIC SUMS FOR COLORINGS AVOIDING MONOCHROMATIC SUBGRAPHS¹

EWA KUBICKA

University of Louisville, USA

e-mail: ewa@louisville.edu

GRZEGORZ KUBICKI

University of Louisville, USA

University of Opole, Poland

e-mail: gkubicki@louisville.edu

AND

KATHLEEN A. McKEON

Connecticut College, USA

e-mail: kamck@conncoll.edu

Abstract

Given graphs G and H , a vertex coloring $c : V(G) \rightarrow \mathbb{N}$ is an H -free coloring of G if no color class contains a subgraph isomorphic to H . The H -free chromatic number of G , $\chi(H, G)$, is the minimum number of colors in an H -free coloring of G . The H -free chromatic sum of G , $\Sigma(H, G)$, is the minimum value achieved by summing the vertex colors of each H -free coloring of G . We provide a general bound for $\Sigma(H, G)$, discuss the computational complexity of finding this parameter for different choices of H , and prove an exact formulas for some graphs G . For every integer k and for every graph H , we construct families of graphs, G_k with the property that k more colors than $\chi(H, G)$ are required to realize $\Sigma(H, G)$ for H -free colorings. More complexity results and constructions of graphs requiring extra colors are given for planar and outerplanar graphs.

Keywords: coloring, sum of colors, forbidden subgraphs.

2010 Mathematics Subject Classification: 05C15.

¹An extended abstract was presented at the conference: Erdős Centennial, July 1-5, 2013, Budapest [20].

REFERENCES

- [1] M. Albertson, R. Jamison, S. Hedetniemi and S. Locke, *The subchromatic number of a graph*, Discrete Math. **74** (1989) 33–49.
doi:10.1016/0012-365X(89)90196-9
- [2] J. Andrews and M. Jacobson, *On a generalization of chromatic number*, Congr. Numer. **47** (1985) 33–48.
- [3] D. Archdeacon, *A note on defective colorings of graphs in surfaces*, J. Graph Theory **11** (1987) 517–519.
doi:10.1002/jgt.3190110408
- [4] M. Borowiecki, I. Broere, M. Frick, P. Mihók and G. Semanišin, *A survey of hereditary properties of graphs*, Discuss. Math. Graph Theory **17** (1997) 5–50.
doi:10.7151/dmgt.1037
- [5] I. Broere and M. Mynhardt, *Generalized colorings of outer-planar and planar graphs*, Graph Theory with Applications to Algorithms and Computer Science, Wiley, New York (1985) 151–161.
- [6] I. Broere, S. Dorfling and E. Jonck, *Generalized chromatic numbers and hereditary properties of graphs*, Discuss. Math. Graph Theory **22** (2002) 259–270.
doi:10.7151/dmgt.1174
- [7] H. Broersma, F.V. Fomin, J. Kratochvíl and G.J. Woeginger, *Planar graph coloring avoiding monochromatic subgraphs: trees, and paths make it difficult*, Algorithmica **44** (2006) 343–361.
doi:10.1007/s00453-005-1176-8
- [8] M.I. Burstein, *The bi-colorability of planar hypergraphs*, Sakhartsh. SSR Mecn. Akad. Moambe **78** (1975) 293–296.
- [9] G. Chartrand, D. Geller and S. Hedetniemi, *A generalization of the chromatic number*, Proc. Cambridge Philos. Soc. **64** (1968) 265–271.
doi:10.1017/S0305004100042808
- [10] G. Chartrand and H. Kronk, *The point-arboricity of planar graphs*, J. London Math. Soc. **44** (1969) 612–616.
doi:10.1112/jlms/s1-44.1.612
- [11] G. Chartrand, H. Kronk and C. Wall, *The point-arboricity of a graph*, Israel J. Math. **6** (1968) 169–175.
doi:10.1007/BF02760181
- [12] L.J. Cowen, R.H. Cowen and D.R. Woodall, *Defective colorings of graphs in surfaces; partitions into subgraphs of bounded valency*, J. Graph Theory **10** (1986) 187–195.
doi:10.1002/jgt.3190100207
- [13] L.J. Cowen, W. Goddard and C.E. Jesurum, *Defective colorings revisited*, J. Graph Theory **24** (1997) 205–219.
doi:10.1002/(SICI)1097-0118(199703)24:3<205::AID-JGT2>3.0.CO;2-T

- [14] K. Dargen and K. Fraughnaugh, *Conditional chromatic numbers with forbidden cycles*, Linear Algebra Appl. **217** (1985) 53–66.
doi:10.1016/0024-3795(94)00139-5
- [15] P. Erdős, E. Kubicka and A.J. Schwenk, *Graphs that require many colors to achieve their chromatic sum*, Proc. Twentieth Southeastern Conference on Combinatorics, Graph Theory, and Computing **71** (1990) 17–28.
- [16] W. Goddard, *Acyclic colorings of planar graphs*, Discrete Math. **91** (1991) 91–94.
doi:10.1016/0012-365X(91)90166-Y
- [17] F. Harary and K. Fraughnaugh (Jones), *Conditional colorability II: bipartite variations*, Congr. Numer. **50** (1985) 205–218.
- [18] F. Harary and K. Fraughnaugh (Jones), *Degree conditional bipartition numbers in graphs*, Congr. Numer. **55** (1986) 39–50.
- [19] E. Kubicka, *The chromatic sums and efficient tree algorithms*, Ph.D. Thesis, Western Michigan University (1989).
- [20] E. Kubicka, G. Kubicki and K. McKeon, *Chromatic sums for colorings avoiding monochromatic subgraphs*, Electron. Notes Discrete Math. **43** (2013) 247–254.
doi:10.1016/j.endm.2013.07.041
- [21] E. Kubicka and A. Schwenk, *Introduction to chromatic sums*, Congr. Numer. **71** (1990) 7–28.
- [22] K. McKeon, *Generalized chromatic numbers of graphs with bipartite complements*, Congr. Numer. **197** (2009) 97–105.
- [23] K. McKeon and H. Pham, *Generalized chromatic numbers of graphs with prescribed complements*, Congr. Numer. **191** (2008) 71–79.
- [24] K.S. Poh, *On the linear vertex-arboricity of a planar graph*, J. Graph Theory **14** (1990) 73–75.
doi:10.1002/jgt.3190140108
- [25] C. Thomassen, *Decomposing a planar graph into degenerate graphs*, J. Combin. Theory Ser. B **65** (1995) 305–314.
doi:10.1006/jctb.1995.1057

Received 27 February 2013

Revised 10 November 2014

Accepted 10 November 2014