γ -CYCLES IN ARC-COLORED DIGRAPHS

HORTENSIA GALEANA-SÁNCHEZ¹

GUADALUPE GAYTÁN-GÓMEZ

*Instituto de Matemáticas
Universidad Nacional Autónoma de México
Ciudad Universitaria, México, D.F. 04510, México*

e-mail: hgaleana@matem.unam.mx
ggg_19808@hotmail.com

AND

ROCÍO ROJAS-MONROY

*Facultad de Ciencias
Universidad Autónoma del Estado de México
Instituto Literario No. 100, Centro 50000
Toluca, Edo. de México, México*

e-mail: mrrm@uaemex.mx

Abstract

We call a digraph D an m -colored digraph if the arcs of D are colored with m colors. A directed path (or a directed cycle) is called monochromatic if all of its arcs are colored alike. A subdigraph H in D is called rainbow if all of its arcs have different colors. A set $N \subseteq V(D)$ is said to be a kernel by monochromatic paths of D if it satisfies the two following conditions:

- (i) for every pair of different vertices $u, v \in N$ there is no monochromatic path in D between them, and
- (ii) for every vertex $x \in V(D) - N$ there is a vertex $y \in N$ such that there is an xy -monochromatic path in D .

A γ -cycle in D is a sequence of different vertices $\gamma = (u_0, u_1, \dots, u_n, u_0)$ such that for every $i \in \{0, 1, \dots, n\}$:

- (i) there is a $u_i u_{i+1}$ -monochromatic path, and
- (ii) there is no $u_{i+1} u_i$ -monochromatic path.

The addition over the indices of the vertices of γ is taken modulo $(n+1)$. If D is an m -colored digraph, then the closure of D , denoted by $\mathfrak{C}(D)$, is the m -colored multidigraph defined as follows: $V(\mathfrak{C}(D)) = V(D)$, $A(\mathfrak{C}(D)) =$

¹This work was partially supported by CONACYT 219840-2013 and PAPIIT IN106613.

$A(D) \cup \{(u, v) \text{ with color } i \mid \text{there exists a } uv\text{-monochromatic path colored } i \text{ contained in } D\}$.

In this work, we prove the following result. Let D be a finite m -colored digraph which satisfies that there is a partition $C = C_1 \cup C_2$ of the set of colors of D such that:

- (1) $D[\widehat{C}_i]$ (the subdigraph spanned by the arcs with colors in C_i) contains no γ -cycles for $i \in \{1, 2\}$;
- (2) If $\mathfrak{C}(D)$ contains a rainbow $C_3 = (x_0, z, w, x_0)$ involving colors of C_1 and C_2 , then $(x_0, w) \in A(\mathfrak{C}(D))$ or $(z, x_0) \in A(\mathfrak{C}(D))$;
- (3) If $\mathfrak{C}(D)$ contains a rainbow $P_3 = (u, z, w, x_0)$ involving colors of C_1 and C_2 , then at least one of the following pairs of vertices is an arc in $\mathfrak{C}(D)$: (u, w) , (w, u) , (x_0, u) , (u, x_0) , (x_0, w) , (z, u) , (z, x_0) .

Then D has a kernel by monochromatic paths.

This theorem can be applied to all those digraphs that contain no γ -cycles. Generalizations of many previous results are obtained as a direct consequence of this theorem.

Keywords: digraph, kernel, kernel by monochromatic paths, γ -cycle.

2010 Mathematics Subject Classification: 05C20.

REFERENCES

- [1] C. Berge, *Graphs* (North-Holland, Amsterdam, 1985).
- [2] P. Duchet, *Graphes Noyau-Parfaits*, Ann. Discrete Math. **9** (1980) 93–101.
doi:10.1016/S0167-5060(08)70041-4
- [3] P. Duchet, *Classical perfect graphs: An introduction with emphasis on triangulated and interval graphs*, Ann. Discrete Math. **21** (1984) 67–96.
- [4] P. Duchet and H. Meyniel, *A note on kernel-critical graphs*, Discrete Math. **33** (1981) 103–105.
doi:10.1016/0012-365X(81)90264-8
- [5] H. Galeana-Sánchez, *On monochromatic paths and monochromatic cycles in edge colored tournaments*, Discrete Math. **156** (1996) 103–112.
doi:10.1016/0012-365X(95)00036-V
- [6] H. Galeana-Sánchez, *Kernels in edge-coloured digraphs*, Discrete Math. **184** (1998) 87–99.
doi:10.1016/S0012-365X(97)00162-3
- [7] H. Galeana-Sánchez and J.J. García-Ruvalcaba, *Kernels in the closure of coloured digraphs*, Discuss. Math. Graph Theory **20** (2000) 243–254.
doi:10.7151/dmgt.1123
- [8] H. Galeana-Sánchez, G. Gaytán-Gómez and R. Rojas-Monroy, *Monochromatic cycles and monochromatic paths in arc-colored digraphs*, Discuss. Math. Graph Theory **31** (2011) 283–292.
doi:10.7151/dmgt.1545

- [9] H. Galeana-Sánchez and V. Neumann-Lara, *On kernels and semikernels of digraphs*, Discrete Math. **48** (1984) 67–76.
doi:10.1016/0012-365X(84)90131-6
- [10] H. Galeana-Sánchez and V. Neumann-Lara, *On kernel-perfect critical digraphs*, Discrete Math. **59** (1986) 257–265.
doi:10.1016/0012-365X(86)90172-X
- [11] H. Galeana-Sánchez and R. Rojas-Monroy, *A counterexample to a conjecture on edge-coloured tournaments*, Discrete Math. **282** (2004) 275–276.
doi:10.1016/j.disc.2003.11.015
- [12] H. Galeana-Sánchez and R. Rojas-Monroy, *On monochromatic paths and monochromatic 4-cycles in edge coloured bipartite tournaments*, Discrete Math. **285** (2004) 313–318.
doi:10.1016/j.disc.2004.03.005
- [13] H. Galeana-Sánchez and R. Rojas-Monroy, *Monochromatic paths and at most 2-coloured arc sets in edge-coloured tournaments*, Graphs Combin. **21** (2005) 307–317.
doi:10.1007/s00373-005-0618-z
- [14] H. Galeana-Sánchez and R. Rojas-Monroy, *Monochromatic paths and quasi-monochromatic cycles in edge-coloured bipartite tournaments*, Discuss. Math. Graph Theory **28** (2008) 285–306.
doi:10.7151/dmgt.1406
- [15] H. Galeana-Sánchez and R. Rojas-Monroy, *Independent domination by monochromatic paths in arc coloured bipartite tournaments*, AKCE Int. J. Graphs Comb. **6** (2009) 267–285.
- [16] H. Galeana-Sánchez, R. Rojas-Monroy and B. Zavala, *Monochromatic paths and monochromatic sets of arcs in 3-quasitransitive digraphs*, Discuss. Math. Graph Theory **29** (2009) 337–347.
doi:10.7151/dmgt.1450
- [17] J. von Neumann and O. Morgenstern, *Theory of Games and Economic Behavior* (Princeton University Press, Princeton, 1944).
- [18] M. Richardson, *Solutions of irreflexive relations*, Ann. of Math. **58** (1953) 573–590.
doi:10.2307/1969755
- [19] M. Richardson, *Extension theorems for solutions of irreflexive relations*, Proc. Natl. Acad. Sci. USA **39** (1953) 649–655.
doi:10.1073/pnas.39.7.649
- [20] R. Rojas-Monroy and J.I. Villarreal-Valdés, *Kernels in infinite diraphs*, AKCE Int. J. Graphs Comb. **7** (2010) 103–111.
- [21] Shen Minggang, *On monochromatic paths in m -coloured tournaments*, J. Combin. Theory Ser. B **45** (1988) 108–111.
doi:10.1016/0095-8956(88)90059-7

- [22] B. Sands, N. Sauer and R. Woodrow, *On monochromatic paths edge-coloured digraphs*, J. Combin. Theory Ser. B **33** (1982) 271–275.
doi:10.1016/0095-8956(82)90047-8
- [23] I. Włoch, *On imp-sets and kernels by monochromatic paths of the duplication*, Ars Combin. **83** (2007) 93–99.
- [24] I. Włoch, *On kernels by monochromatic paths in the corona of digraphs*, Cent. Eur. J. Math. **6** (2008) 537–542.
doi:10.2478/s11533-008-0044-6

Received 9 June 2014
Revised 1 May 2015
Accepted 1 May 2015