A MAXIMUM RESONANT SET OF POLYOMINO GRAPHS¹

HEPING ZHANG² AND XIANGQIAN ZHOU

*School of Mathematics and Statistics
Lanzhou University
Lanzhou, Gansu 730000, P.R. China*

e-mail: zhanghp@lzu.edu.cn
zhouxiangqian0502@126.com

Abstract

A polyomino graph P is a connected finite subgraph of the infinite plane grid such that each finite face is surrounded by a regular square of side length one and each edge belongs to at least one square. A dimer covering of P corresponds to a perfect matching. Different dimer coverings can interact via an alternating cycle (or square) with respect to them. A set of disjoint squares of P is a resonant set if P has a perfect matching M so that each one of those squares is M -alternating. In this paper, we show that if K is a maximum resonant set of P , then $P - K$ has a unique perfect matching. We further prove that the maximum forcing number of a polyomino graph is equal to the cardinality of a maximum resonant set. This confirms a conjecture of Xu *et al.* [26]. We also show that if K is a maximal alternating set of P , then $P - K$ has a unique perfect matching.

Keywords: polyomino graph, dimer problem, perfect matching, resonant set, forcing number, alternating set.

2010 Mathematics Subject Classification: 05C70, 05C90, 92E10.

REFERENCES

- [1] H. Abeledo and G.W. Atkinson, *Unimodularity of the Clar number problem*, Linear Algebra Appl. **420** (2007) 441–448.
doi:10.1016/j.laa.2006.07.026
- [2] C. Berge, C.C. Chen, V. Chvátal and C.S. Seow, *Combinatorial properties of polyominoes*, Combinatorica **1** (1981) 217–224.
doi:10.1007/BF02579327

¹This work is supported by NSFC (grant no. 11371180).

²Corresponding author.

- [3] Z. Che and Z. Chen, *Forcing on perfect matchings — A survey*, MATCH Commun. Math. Comput. Chem. **66** (2011) 93–136.
- [4] E. Clar, *The Aromatic Sextet* (Wiley, London, 1972).
- [5] M.E. Fisher, *Statistical mechanics of dimers on a plane lattice*, Phys. Rev. **124** (1961) 1664–1672.
doi:10.1103/PhysRev.124.1664
- [6] E.J. Cockayne, *Chessboard domination problems*, Discrete Math. **86** (1990) 13–20.
doi:10.1016/0012-365X(90)90344-H
- [7] C.M. Grinstead, B. Hahne and D. Van Stone, *On the queen domination problem*, Discrete Math. **86** (1990) 21–26.
doi:10.1016/0012-365X(90)90345-I
- [8] I. Gutman, S.J. Cyvin, *Advances in the Theory of Benzenoid Hydrocarbons* (Springer, Berlin, 1990).
doi:10.1007/3-540-51505-4
- [9] F. Harary, D.J. Klein and T.P. Živković, *Graphical properties of polyhexes: Perfect matching vector and forcing*, J. Math. Chem. **6** (1991) 295–306.
doi:10.1007/BF01192587
- [10] W.C. Herndon, *Resonance energies of aromatic hydrocarbons: Quantitative test of resonance theory*, J. Am. Chem. Soc. **95** (1973) 2404–2406.
doi:10.1021/ja00788a073
- [11] P.W. Kasteleyn, *The statistics of dimers on a lattice: I. The number of dimer arrangements on a quadratic lattice*, Physica **27** (1961) 1209–1225.
doi:10.1016/0031-8914(61)90063-5
- [12] X. Ke, *A lower bound on the number of elementary components of essentially disconnected generalized polyomino graphs*, J. Math. Chem. **50** (2012) 131–140.
doi:10.1007/s10910-011-9900-x
- [13] D.J. Klein and M. Randić, *Innate degree of freedom of a graph*, J. Comput. Chem. **8** (1987) 516–521.
doi:10.1002/jcc.540080432
- [14] W. Li and H. Zhang, *Dimer statistics of honeycomb lattices on Klein bottle, Möbius strip and cylinder*, Phys. A **391** (2012) 3833–3848.
doi:10.1016/j.physa.2012.03.004
- [15] S. Liu and J. Ou, *On maximal resonance of polyomino graphs*, J. Math. Chem. **51** (2013) 603–619.
doi:10.1007/s10910-012-0104-9
- [16] F. Lu and L. Zhang, *Dimers on two types of lattices on the Klein bottle*, J. Phys. A **45** (2012) #49.
doi:10.1088/1751-8113/45/49/494012
- [17] L. Lovász and M.D. Plummer, *Matching Theory* (Annals of Discrete Mathematics, Vol. **29**, North-Holland, Amsterdam, 1986).

- [18] A. Motoyama and H. Hosoya, *King and domino polynomials for polyomino graphs*, J. Math. Phys. **18** (1977) 1485–1490.
doi:10.1063/1.523411
- [19] L. Pachter and P. Kim, *Forcing matchings on square grids*, Discrete Math. **190** (1998) 287–294.
doi:10.1016/S0012-365X(97)00266-5
- [20] M. Randić, *Conjugated circuits and resonance energies of benzenoid hydrocarbons*, Chem. Phys. Lett. **38** (1976) 68–70.
doi:10.1016/0009-2614(76)80257-6
- [21] M. Randić, *Aromaticity and conjugation*, J. Am. Chem. Soc. **99** (1977) 444–450.
doi:10.1021/ja00444a022
- [22] H. Sachs, *Perfect matchings in hexagonal systems*, Combinatorica **4** (1980) 89–99.
doi:10.1007/BF02579161
- [23] H. Sachs and H. Zernitz, *Remark on the dimer problem*, Discrete Appl. Math. **51** (1994) 171–179.
doi:10.1016/0166-218X(94)90106-6
- [24] K. Salem and H. Abeledo, *A maximal alternating set of a hexagonal system*, MATCH Commun. Math. Comput. Chem. **55** (2006) 159–176.
- [25] S. Wei and X. Ke, *Elementary components of essentially disconnected polyomino graphs*, J. Math. Chem. **47** (2010) 496–504.
doi:10.1007/s10910-009-9589-2
- [26] L. Xu, H. Bian and F. Zhang, *Maximum forcing number of hexagonal systems*, MATCH Commun. Math. Comput. Chem. **70** (2013) 493–500.
- [27] W. Yan, Y.-N. Yeh and F. Zhang, *Dimer problem on the cylinder and torus*, Phys. A **387** (2008) 6069–6078.
doi:10.1016/j.physa.2008.06.042
- [28] F. Zhang, X. Guo and R. Chen, *The connectivity of Z-transformation graphs of perfect matchings of hexagonal systems*, Acta Math. Appl. Sin. **4** (1988) 131–135.
doi:10.1007/bf02006061
- [29] H. Zhang, *The connectivity of Z-transformation graphs of perfect matchings of polyominoes*, Discrete Math. **158** (1996) 257–272.
doi:10.1016/0012-365X(95)00048-2
- [30] H. Zhang and F. Zhang, *Perfect matchings of polyomino graphs*, Graphs Combin. **13** (1997) 295–304.
doi:10.1007/BF03353008
- [31] H. Zhang and F. Zhang, *Plane elementary bipartite graphs*, Discrete Appl. Math. **105** (2000) 291–311.
doi:10.1016/S0166-218X(00)00204-3
- [32] M. Zheng and R. Chen, *A maximal cover of hexagonal systems*, Graphs Combin. **1** (1985) 295–298.
doi:10.1007/BF02582955

Received 9 February 2015

Revised 29 June 2015

Accepted 29 June 2015