THE STEINER WIENER INDEX OF A GRAPH

XUELIANG LI¹²

*Center for Combinatorics and LPMC
Nankai University, Tianjin 300071, China*

e-mail: lxl@nankai.edu.cn

YAPING MAO³

*Department of Mathematics
Qinghai Normal University
Qinghai 810008, China*

e-mail: maoyaping@ymail.com

AND

IVAN GUTMAN

*Faculty of Science P.O. Box 60, 34000 Kragujevac, Serbia
and State University of Novi Pazar, Novi Pazar, Serbia*

e-mail: gutman@kg.ac.rs

Abstract

The Wiener index $W(G)$ of a connected graph G , introduced by Wiener in 1947, is defined as $W(G) = \sum_{u,v \in V(G)} d(u,v)$ where $d_G(u,v)$ is the distance between vertices u and v of G . The Steiner distance in a graph, introduced by Chartrand *et al.* in 1989, is a natural generalization of the concept of classical graph distance. For a connected graph G of order at least 2 and $S \subseteq V(G)$, the *Steiner distance* $d(S)$ of the vertices of S is the minimum size of a connected subgraph whose vertex set is S . We now introduce the concept of the Steiner Wiener index of a graph. The *Steiner k -Wiener index* $SW_k(G)$ of G is defined by $SW_k(G) = \sum_{\substack{S \subseteq V(G) \\ |S|=k}} d(S)$. Expressions for SW_k for some special graphs are obtained. We also give sharp upper and lower bounds of SW_k of a connected graph, and establish some of its properties in

¹Supported by NSFC No.11371205 and 11531011.

²Corresponding author.

³Supported by NSFC No.11161037 and 11551001, and SFQP No.2014-ZJ-907.

the case of trees. An application in chemistry of the Steiner Wiener index is reported in our another paper.

Keywords: distance, Steiner distance, Wiener index, Steiner Wiener k -index.

2010 Mathematics Subject Classification: 05C05, 05C12, 05C35.

REFERENCES

- [1] P. Ali, P. Dankelmann and S. Mukwembi, *Upper bounds on the Steiner diameter of a graph*, Discrete Appl. Math. **160** (2012) 1845–1850.
doi:10.1016/j.dam.2012.03.031

- [2] Y. Alizadeh, V. Andova, S. Klavžar and R. Škrekovski, *Wiener dimension: Fundamental properties and $(5, 0)$ -nanotubical fullerenes*, MATCH Commun. Math. Comput. Chem. **72** (2014) 279–294.
- [3] J.A. Bondy and U.S.R. Murty, *Graph Theory* (Springer, 2008).
- [4] F. Buckley and F. Harary, *Distance in Graphs* (Addison-Wesley, Redwood, 1990).
- [5] J. Cáceresa, A. Márquez and M.L. Puertasa, *Steiner distance and convexity in graphs*, European J. Combin. **29** (2008) 726–736.
doi:10.1016/j.ejc.2007.03.007
- [6] G. Chartrand, O.R. Oellermann, S. Tian and H.B. Zou, *Steiner distance in graphs*, Časopis Pest. Mat. **114** (1989) 399–410.
- [7] C.M. da Fonseca, M. Ghebleh, A. Kanso and D. Stevanović, *Counterexamples to a conjecture on Wiener index of common neighborhood graphs*, MATCH Commun. Math. Comput. Chem. **72** (2014) 333–338.
- [8] P. Dankelmann, O.R. Oellermann and H.C. Swart, *The average Steiner distance of a graph*, J. Graph Theory **22** (1996) 15–22.
doi:10.1002/(SICI)1097-0118(199605)22:1(15::AID-JGT3)3.0.CO;2-O
- [9] P. Dankelmann, H.C. Swart and O.R. Oellermann, *On the average Steiner distance of graphs with prescribed properties*, Discrete Appl. Math. **79** (1997) 91–103.
doi:10.1016/S0166-218X(97)00035-8
- [10] A. Dobrynin, R. Entringer and I. Gutman, *Wiener index of trees: theory and application*, Acta Appl. Math. **66** (2001) 211–249.
doi:10.1023/A:1010767517079
- [11] R.C. Entringer, D.E. Jackson and D.A. Snyder, *Distance in graphs*, Czechoslovak Math. J. **26** (1976) 283–296.
- [12] M.R. Garey and D.S. Johnson, *Computers and Intractability—A Guide to the Theory of NP-Completeness* (Freeman, San Francisco, 1979) 208–209.
- [13] W. Goddard and O.R. Oellermann, *Distance in graphs*, in: *Structural Analysis of Complex Networks*, M. Dehmer (Ed.), (Birkhäuser, Dordrecht, 2011) 49–72.
- [14] I. Gutman, B. Furtula and X. Li, *Multicenter Wiener indices and their applications*, J. Serb. Chem. Soc. **80** (2015) 1009–1017.
doi:10.2298/JSC150126015G
- [15] I. Gutman, S. Klavžar and B. Mohar, *Fifty years of the Wiener index*, MATCH Commun. Math. Comput. Chem. **35** (1997) 1–159.
- [16] I. Gutman and O.E. Polansky, *Mathematical Concepts in Organic Chemistry* (Springer, Berlin, 1986).
- [17] Y.L. Jin and X.D. Zhang, *On two conjectures of the Wiener index*, MATCH Commun. Math. Comput. Chem. **70** (2013) 583–589.
- [18] S. Klavžar and M.J. Nadjafi-Arani, *Wiener index in weighted graphs via unification of Θ^* -classes*, European J. Combin. **36** (2014) 71–76.
doi:10.1016/j.ejc.2013.04.008

- [19] M. Knor and R. Škrekovski, *Wiener index of generalized 4-stars and of their quadratic line graphs*, Australas. J. Combin. **58** (2014) 119–126.
doi:10.1002/jgt.3190140510
- [20] O.R. Oellermann and S. Tian, *Steiner centers in graphs*, J. Graph Theory **14** (1990) 585–597.
doi:/10.1002/jgt.3190140510
- [21] D.H. Rouvray, *Harry in the limelight: The life and times of Harry Wiener*, in: Topology in Chemistry—Discrete Mathematics of Molecules, D.H. Rouvray, R.B. King (Eds.), (Horwood, Chichester, 2002) 1–15.
- [22] D.H. Rouvray, *The rich legacy of half century of the Wiener index*, in: Topology in Chemistry—Discrete Mathematics of Molecules, D.H. Rouvray and R.B. King (Eds.), (Horwood, Chichester, 2002) 16–37.
- [23] H. Wiener, *Structural determination of paraffin boiling points*, J. Amer. Chem. Soc. **69** (1947) 17–20.
doi:10.1021/ja01193a005
- [24] K. Xu, M. Liu, K.C. Das, I. Gutman and B. Furtula, *A survey on graphs extremal with respect to distance-based topological indices*, MATCH Commun. Math. Comput. Chem. **71** (2014) 461–508.

Received 27 December 2014

Revised 13 August 2015

Accepted 13 August 2015