REGULARITY AND PLANARITY OF TOKEN GRAPHS

WALTER CARBALLOSA

Department of Mathematics and Statistics
Florida International University, Miami, FL

e-mail: waltercarb@gmail.com

RUY FABILA-MONROY

Departamento de Matemáticas
CINVESTAV, Mexico

e-mail: ruyfabila@math.cinvestav.edu.mx

JESÚS LEAÑOS

Faculty of Natural Sciences and Mathematics
University of Maribor, Slovenia

e-mail: jesus.leanos@gmail.com

AND

LUIS MANUEL RIVERA

Unidad Académica de Matemáticas
Universidad Autónoma de Zacatecas, Zac., Mexico

e-mail: luismanuel.rivera@gmail.com

Abstract

Let $G = (V, E)$ be a graph of order n and let $1 \leq k < n$ be an integer. The k -token graph of G is the graph whose vertices are all the k -subsets of V , two of which are adjacent whenever their symmetric difference is a pair of adjacent vertices in G . In this paper we characterize precisely, for each value of k , which graphs have a regular k -token graph and which connected graphs have a planar k -token graph.

Keywords: token graph, Johnson graph, regularity, planarity.

2010 Mathematics Subject Classification: 05C10, 05C69.

REFERENCES

- [1] J.M. Boyer and J.W. Myrvold, *On the cutting edge: simplified $O(n)$ planarity by edge addition*, J. Graph Algorithms Appl. **8** (2004) 241–273.
doi:10.7155/jgaa.00091
- [2] M. Behzad and S.E. Mahmoudian, *On topological invariants of the product of graphs*, Canad. Math. Bull. **12** (1969) 175–166.
doi:10.4153/CMB-1969-017-3
- [3] T. Etzion and S. Bitan, *On the chromatic number, colorings, and codes of the Johnson graph*, Discrete Appl. Math. **70** (1996) 163–175.
doi:10.1016/0166-218X(96)00104-7
- [4] R. Fabila-Monroy, D. Flores-Peñaloza, C. Huemer, F. Hurtado, J. Urrutia and D.R. Wood, *Token graphs*, Graphs Combin. **28** (2012) 365–380.
doi:10.1007/s00373-011-1055-9
- [5] R. Graham and N. Sloane, *Lower bounds for constant weight codes*, IEEE Trans. Inform. Theory **26** (1980) 37–43.
doi:10.1109/TIT.1980.1056141
- [6] J. Guo, K. Wang and F. Li, *Metric dimension of some distance-regular graphs*, J. Comb. Optim. **26** (2013) 190–197.
doi:10.1007/s10878-012-9459-x
- [7] F. Harary, Graph Theory (Addison-Wesley, Reading, MA, 1969).
- [8] K. Kuratowski, *Sur le problème des courbes gauches en topologie*, Fund. Math. **15** (1930) 271–283.
- [9] R.A. Liebler and C.E. Praeger, *Neighbour-transitive codes in Johnson graphs*, Des. Codes Cryptogr. **73** (2014) 1–25.
doi:10.1007/s10623-014-9982-0
- [10] B.D. McKay and A. Piperno, *Practical graph isomorphism II*, J. Symbolic Comput. **60** (2013) 94–112.
doi:10.1016/j.jsc.2013.09.003
- [11] M. Neunhöffer, and C.E. Praeger, *Sporadic neighbour-transitive codes in Johnson graphs*, Des. Codes Cryptogr. **72** (2014) 141–152.
doi:10.1007/s10623-013-9853-0
- [12] W.A. Stein, *et al.*, Sage Mathematics Software (Version 6.8), The Sage Development Team, 2015.
<http://www.sagemath.org>.
- [13] K. Wagner, *Über eine Eigenschaft der ebenen Komplexe*, Math. Ann. **114** (1937) 570–590.
doi:10.1007/BF01594196

Received 14 December 2015

Revised 26 May 2016

Accepted 26 May 2016