RAMSEY PROPERTIES OF RANDOM GRAPHS AND FOLKMAN NUMBERS

VOJTĚCH RÖDL¹

Emory University, Atlanta, USA

e-mail: rodl@mathcs.emory.edu

ANDRZEJ RUCIŃSKI²

A. Mickiewicz University, Poznań, Poland

e-mail: rucinski@amu.edu.pl

AND

MATHIAS SCHACHT³

Universität Hamburg, Hamburg, Germany

e-mail: schacht@math.uni-hamburg.de

Abstract

For two graphs, G and F , and an integer $r \geq 2$ we write $G \rightarrow (F)_r$ if every r -coloring of the edges of G results in a monochromatic copy of F . In 1995, the first two authors established a threshold edge probability for the Ramsey property $G(n, p) \rightarrow (F)_r$, where $G(n, p)$ is a random graph obtained by including each edge of the complete graph on n vertices, independently, with probability p . The original proof was based on the regularity lemma of Szemerédi and this led to tower-type dependencies between the involved parameters. Here, for $r = 2$, we provide a self-contained proof of a quantitative version of the Ramsey threshold theorem with only double exponential dependencies between the constants. As a corollary we obtain a double exponential upper bound on the 2-color Folkman numbers. By a different proof technique, a similar result was obtained independently by Conlon and Gowers.

Keywords: Ramsey property, random graph, Folkman number .

2010 Mathematics Subject Classification: 05C80, 05C55 .

¹Research supported by NSF grant DMS 080070.

²Research supported by the Polish NSC grant 2014/15/B/ST1/01688. Part of research performed at Emory University, Atlanta.

³Research supported by the Heisenberg-Programme of the Deutsche Forschungsgemeinschaft.

REFERENCES

- [1] J. Balogh, R. Morris and W. Samotij, *Independent sets in hypergraphs*, J. Amer. Math. Soc. **28** (2015) 669–709.
doi:10.1090/S0894-0347-2014-00816-X
- [2] D. Conlon and T. Gowers, *An upper bound for Folkman numbers*, preprint.
- [3] E. Friedgut, V. Rödl and M. Schacht, *Ramsey properties of random discrete structures*, Random Structures Algorithms **37** (2010) 407–436.
doi:10.1002/rsa.20352
- [4] S. Janson, T. Łuczak and A. Ruciński, Random Graphs (John Wiley and Sons, New York, 2000).
doi:10.1002/9781118032718
- [5] R. Nenadov and A. Steger, *A short proof of the random Ramsey theorem*, Combin. Probab. Comput. **25** (2016) 130–144.
doi:10.1017/S0963548314000832
- [6] V. Rödl and A. Ruciński, *Threshold functions for Ramsey properties*, J. Amer. Math. Soc. **84** (1995) 917–942.
doi:10.2307/2152833
- [7] V. Rödl, A. Ruciński and M. Schacht, *Ramsey properties of random k -partite, k -uniform hypergraphs*, SIAM J. Discrete Math. **21** (2007) 442–460.
doi:10.1137/060657492
- [8] V. Rödl, A. Ruciński and M. Schacht, *An exponential-type upper bound for Folkman numbers*, Combinatorica, in press.
doi:10.1007/s00493-015-3298-1
- [9] D. Saxton and A. Thomason, *Hypergraph containers*, Invent. Math. **201** (2015) 925–992.
doi:10.1007/s00222-014-0562-8
- [10] E. Szemerédi, *Regular partitions of graphs*, in: Problèmes Combinatoires et Théorie des Graphes (Colloq. Internat. CNRS, Univ. Orsay, Orsay, 1976), Colloq. Internat. CNRS **260** (CNRS, Paris, 1978) 399–401.

Received 11 September 2015

Revised 28 June 2016

Accepted 29 June 2016