BOUNDS ON THE LOCATING ROMAN DOMINATION NUMBER IN TREES

NADER JAFARI RAD AND HADI RAHBANI

*Department of Mathematics
Shahrood University of Technology
Shahrood, Iran*

e-mail: n.jafarirad@gmail.com

Abstract

A Roman dominating function (or just RDF) on a graph $G = (V, E)$ is a function $f : V \rightarrow \{0, 1, 2\}$ satisfying the condition that every vertex u for which $f(u) = 0$ is adjacent to at least one vertex v for which $f(v) = 2$. The weight of an RDF f is the value $f(V(G)) = \sum_{u \in V(G)} f(u)$. An RDF f can be represented as $f = (V_0, V_1, V_2)$, where $V_i = \{v \in V : f(v) = i\}$ for $i = 0, 1, 2$. An RDF $f = (V_0, V_1, V_2)$ is called a locating Roman dominating function (or just LRDF) if $N(u) \cap V_2 \neq N(v) \cap V_2$ for any pair u, v of distinct vertices of V_0 . The locating Roman domination number $\gamma_R^L(G)$ is the minimum weight of an LRDF of G . In this paper, we study the locating Roman domination number in trees. We obtain lower and upper bounds for the locating Roman domination number of a tree in terms of its order and the number of leaves and support vertices, and characterize trees achieving equality for the bounds.

Keywords: Roman domination number, locating domination number, locating Roman domination number, tree.

2010 Mathematics Subject Classification: 05C69.

REFERENCES

- [1] M. Adabi, E. Ebrahimi Targhi, N. Jafari Rad and M. Saied Moradi, *Properties of independent Roman domination in graphs*, Australas. J. Combin. **52** (2012) 11–18.
- [2] H.A. Ahangar, M.A. Henning, V. Samodivkin and I.G. Yero, *Total Roman domination in graphs*, manuscript (2014).
- [3] H.A. Ahangar, M.A. Henning, C. Löwenstein, Y. Zhao and V. Samodivkin, *Signed Roman domination in graphs*, J. Comb. Optim. **27** (2014) 241–255.
doi:10.1007/s10878-012-9500-0

- [4] M. Blidia, M. Chellali, F. Maffray, J. Moncel and A. Semri, *Locating-domination and identifying codes in trees*, Australas. J. Combin. **39** (2007) 219–232.
- [5] M. Blidia, O. Favaron and R. Lounes, *Locating-domination, 2-domination and independence in trees*, Australas. J. Combin. **42** (2008) 309–319.
- [6] M. Chellali, *On locating and differentiating-total domination in trees*, Discuss. Math. Graph Theory **28** (2008) 383–392.
doi:10.7151/dmgt.1414
- [7] M. Chellali, T.W. Haynes, S.T. Hedetniemi and A.A. McRae, *Roman 2-domination*, Discrete Appl. Math. **204** (2016) 22–28.
doi:10.1016/j.dam.2015.11.013
- [8] M. Chellali and N. Jafari Rad, *Locating-total domination critical graphs*, Australas. J. Combin. **45** (2009) 227–234.
- [9] X.G. Chen and M.Y. Sohn, *Bounds on the locating-total domination number of a tree*, Discrete Appl. Math. **159** (2011) 769–773.
doi:10.1016/j.dam.2010.12.025
- [10] E.J. Cockayne, Paul A. Dreyer Jr., S.M. Hedetniemi and S.T. Hedetniemi, *Roman domination in graphs*, Discrete Math. **278** (2004) 11–22.
doi:10.1016/j.disc.2003.06.004
- [11] F. Foucaud, M.A. Henning, C. Löwenstein and T. Sass, *Locating-dominating sets in twin-free graphs*, Discrete Appl. Math. **200** (2016) 52–58.
doi:10.1016/j.dam.2015.06.038
- [12] T.W. Haynes, M.A. Henning and J. Howard, *Locating and total dominating sets in trees*, Discrete Appl. Math. **154** (2006) 1293–1300.
doi:10.1016/j.dam.2006.01.002
- [13] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, *Fundamentals of Domination in Graphs* (Marcel Dekker, Inc., New York, 1998).
- [14] M.A. Henning and N. Jafari Rad, *Locating-total domination in graphs*, Discrete Appl. Math. **160** (2012) 1986–1993.
doi:10.1016/j.dam.2012.04.004
- [15] M.A. Henning and C. Löwenstein, *Locating-total domination in claw-free cubic graphs*, Discrete Math. **312** (2012) 3107–3116.
doi:10.1016/j.disc.2012.06.024
- [16] N. Jafari Rad, H. Rahbani and L. Volkmann, *Locating Roman domination in graphs*, manuscript (2015).
- [17] K. Kammerling and L. Volkmann, *Roman k -domination in graphs*, J. Korean Math. Soc. **46** (2009) 1309–1318.
doi:10.4134/JKMS.2009.46.6.1309
- [18] J.L. Sewell and P.J. Slater, *A sharp lower bound for locating-dominating sets in trees*, Australas. J. Combin. **60** (2014) 136–149.

- [19] C.S. ReVelle and K.E. Rosing, *Defendens imperium Romanum: a classical problem in military strategy*, Amer Math. Monthly **107** (2000) 585–594.
doi:10.2307/2589113
- [20] P. Roushini Leely Pushpam and T.N.M. Nalini Mai, *Edge Roman domination in graphs*, J. Combin. Math. Combin. Comput. **69** (2009) 175–182.
- [21] P.J. Slater, *Dominating and location in acyclic graphs*, Networks **17** (1987) 55–64.
doi:10.1002/net.3230170105
- [22] P.J. Slater, *Dominating and reference sets in graphs*, J. Math. Phys. Sci. **22** (1988) 445–455.
- [23] S.J. Seo and P.J. Slater, *Open neighborhood locating-dominating sets*, Australas. J. Combin. **46** (2010) 109–120.
- [24] I. Stewart, *Defend the Roman Empire!*, Sci. Amer. **281** (1999) 136–139.
doi:10.1038/scientificamerican1299-136

Received 7 January 2016
 Revised 21 September 2016
 Accepted 21 September 2016