UPPER BOUNDS FOR THE STRONG CHROMATIC INDEX OF HALIN GRAPHS

ZIYU HU

Department of Mathematical Sciences
Florida Atlantic University

e-mail: azuth.hu@gmail.com

KO-WEI LIH

Institute of Mathematics, Academia Sinica

e-mail: makwlih@sinica.edu.tw

AND

DAPHNE DER-FEN LIU¹

Department of Mathematics
California State University Los Angeles

e-mail: dliu@calstatela.edu

Abstract

Keywords: strong edge-coloring, strong chromatic index, Halin graphs.

2010 Mathematics Subject Classification: 05C15.

The strong chromatic index of a graph G , denoted by $\chi'_s(G)$, is the minimum number of vertex induced matchings needed to partition the edge set of G . Let T be a tree without vertices of degree 2 and have at least one vertex of degree greater than 2. We construct a Halin graph G by drawing T on the plane and then drawing a cycle C connecting all its leaves in such a way that C forms the boundary of the unbounded face. We call T the characteristic tree of G . Let G denote a Halin graph with maximum degree Δ and characteristic tree T . We prove that $\chi'_s(G) \leq 2\Delta + 1$ when $\Delta \geq 4$. In addition, we show that if $\Delta = 4$ and G is not a wheel, then $\chi'_s(G) \leq \chi'_s(T) + 2$. A similar result for $\Delta = 3$ was established by Lih and Liu [21].

¹Corresponding author. Supported in part by grants NASA MIRO NX15AQ06A and NSF DMS 1600778.

REFERENCES

- [1] L.D. Andersen, *The strong chromatic index of a cubic graph is at most 10*, Discrete Math. **108** (1992) 231–252.
doi:10.1016/0012-365X(92)90678-9
- [2] J. Bensmail, A. Harutyunyan, H. Hocquard and P. Valicov, *Strong edge-coloring of sparse planar graphs*, Discrete Appl. Math. **179** (2014) 229–234.
doi:10.1016/j.dam.2014.07.006
- [3] O.V. Borodin and A.O. Ivanova, *Precise upper bound for the strong edge chromatic number of sparse planar graphs*, Discuss. Math. Graph Theory **33** (2013) 759–770.
doi:10.7151/dmgt.1708
- [4] H. Bruhn and F. Joos, *A stronger bound for the strong chromatic index*, Electron. Notes Discrete Math. **49** (2015) 277–284.
doi:10.1016/j.endm.2015.06.038
- [5] G.J. Chang and G.-H. Duh, *On the precise value of the strong chromatic-index of a planar graph with a large girth*.
arXiv: 1508.03052
- [6] G.J. Chang and D. Liu, *Strong edge-coloring for cubic Halin graphs*, Discrete Math. **312** (2012) 1468–1475.
doi:10.1016/j.disc.2012.01.014
- [7] G.J. Chang, M. Montassier, A. Pêcher and A. Raspaud, *Strong chromatic index of planar graphs with large girth*, Discuss. Math. Graph Theory **34** (2014) 723–733.
doi:10.7151/dmgt.1763
- [8] D.W. Cranston, *Strong edge-coloring of graphs with maximum degree 4 using 22 colors*, Discrete Math. **306** (2006) 2772–2778.
doi:10.1016/j.disc.2006.03.053
- [9] P. DeOrsey, J. Diemunsch, M. Ferrara, N. Graber, S.G. Hartke, S. Jahanbekam, B. Lidicky, L.L. Nelsen, D. Stolee and E. Sullivan, *On the strong chromatic index of sparse graphs*.
arXiv: 1508.03515
- [10] P. Erdős, *Problems and results in combinatorial analysis and graph theory*, Discrete Math. **72** (1988) 81–92.
doi:10.1016/0012-365X(88)90196-3
- [11] P. Erdős and J. Nešetřil, *Problem*, in: Irregularities of Partitions, G. Halász and V.T. Sós (Eds.) (Springer, Berlin, 1989) 162–163.
- [12] R.J. Faudree, R.H. Schelp, A. Gyárfás and Zs. Tuza, *The strong chromatic index of graphs*, Ars Combin. **29B** (1990) 205–211.
- [13] J.L. Fouquet and J. Jolivet, *Strong edge-coloring of graphs and applications to multi- k -gons*, Ars Combin. **16A** (1983) 141–150.
- [14] J.L. Fouquet and J. Jolivet, *Strong edge-coloring of cubic planar graphs*, Progress in Graph Theory in: J.A. Bondy and U.S.R. Murty (Ed(s)), (Academic Press, Toronto, 1984) 247–264.

- [15] M.C. Golumbic and M. Lewenstein, *New results on induced matchings*, Discrete Appl. Math. **101** (2000) 157–165.
doi:10.1016/S0166-218X(99)00194-8
- [16] H. Hocquard, M. Montassier, A. Raspaud and P. Valicov, *On strong edge-colouring of subcubic graphs*, Discrete Appl. Math. **161** (2013) 2467–2479.
doi:10.1016/j.dam.2013.05.021
- [17] P. Horák, *The strong chromatic index of graphs with maximum degree four*, in: R. Bodendiek and K. Wagner (Ed(s)), Contemporary Methods in Graph Theory (BI Wissenschaft-Verlag, Mannheim, 1990) 399–403.
- [18] P. Horák, H. Qing and W.T. Trotter, *Induced matchings in cubic graphs*, J. Graph Theory **17** (1993) 151–160.
doi:10.1002/jgt.3190170204
- [19] D. Hudák, B. Lužar, R. Soták and R. Škrekovski, *Strong edge-coloring of planar graphs*, Discrete Math. **324** (2014) 41–49.
doi:10.1016/j.disc.2014.02.002
- [20] H.-H. Lai, K.-W. Lih and P.-Y. Tsai, *The strong chromatic index of Halin graphs*, Discrete Math. **312** (2012) 1536–1541.
doi:10.1016/j.disc.2011.09.016
- [21] K.-W. Lih and D.D.-F. Liu, *On the strong chromatic index of cubic Halin graphs*, Appl. Math. Lett. **25** (2012) 898–901.
doi:10.1016/j.aml.2011.10.046
- [22] M. Molloy and B. Reed, *A bound on the strong chromatic index of a graph*, J. Combin. Theory Ser. B **69** (1997) 103–109.
doi:10.1006/jctb.1997.1724
- [23] W.C. Shiu, P.C.B. Lam and W.K. Tam, *On strong chromatic index of Halin graphs*, J. Combin. Math. Combin. Comput. **57** (2006) 211–222.
- [24] W.C. Shiu and W.K. Tam, *The strong chromatic index of complete cubic Halin graphs*, Appl. Math. Lett. **22** (2009) 754–758.
doi:10.1016/j.aml.2008.08.019
- [25] J. Wu and W. Lin, *The strong chromatic index of a class of graphs*, Discrete Math. **308** (2008) 6254–6261.
doi:10.1016/j.disc.2007.11.051

Received 17 September 2015

Revised 30 August 2016

Accepted 30 August 2016