ARC-DISJOINT HAMILTONIAN CYCLES IN ROUND DECOMPOSABLE LOCALLY SEMICOMPLETE DIGRAPHS

RUIJUAN LI¹ AND TINGTING HAN

School of Mathematical Sciences
Shanxi University
030006 Taiyuan, P.R. China

e-mail: ruijuanli@sxu.edu.cn
tingtinghan19@163.com

Abstract

Let $D = (V, A)$ be a digraph; if there is at least one arc between every pair of distinct vertices of D , then D is a semicomplete digraph. A digraph D is locally semicomplete if for every vertex x , the out-neighbours of x induce a semicomplete digraph and the in-neighbours of x induce a semicomplete digraph. A locally semicomplete digraph without 2-cycle is a local tournament. In 2012, Bang-Jensen and Huang [J. Combin Theory Ser. B 102 (2012) 701–714] concluded that every 2-arc-strong locally semicomplete digraph which is not the second power of an even cycle has two arc-disjoint strong spanning subdigraphs, and proposed the conjecture that every 3-strong local tournament has two arc-disjoint Hamiltonian cycles. According to Bang-Jensen, Guo, Gutin and Volkmann, locally semicomplete digraphs have three subclasses: the round decomposable; the non-round decomposable which are not semicomplete; the non-round decomposable which are semicomplete. In this paper, we prove that every 3-strong round decomposable locally semicomplete digraph has two arc-disjoint Hamiltonian cycles, which implies that the conjecture holds for the round decomposable local tournaments. Also, we characterize the 2-strong round decomposable local tournaments each of which contains a Hamiltonian path P and a Hamiltonian cycle arc-disjoint from P .

Keywords: locally semicomplete digraph, local tournament, round decomposable, arc-disjoint, Hamiltonian cycle, Hamiltonian path.

2010 Mathematics Subject Classification: 05C20.

¹Research is supported partially by NNSFC under no. 11401353, NSF of Shanxi Province, China under no. 2016011005 and TYAL of Shanxi.

REFERENCES

- [1] J. Bang-Jensen and G. Gutin, *Digraphs: Theory, Algorithms and Applications* (Springer Monographs in Mathematics, Springer-Verlag, London, 2009).
doi:10.1007/978-1-84800-998-1
- [2] J. Bang-Jensen, *Locally semicomplete digraph: A generalization of tournaments*, J. Graph Theory **14** (1990) 371–390.
doi:10.1002/jgt.3190140310
- [3] J. Bang-Jensen, Y. Guo, G. Gutin and L. Volkmann, *A classification of locally semicomplete digraphs*, Discrete Math. **167-168** (1997) 101–114.
doi:10.1016/S0012-365X(96)00219-1
- [4] J. Bang-Jensen and J. Huang, *Decomposing locally semicomplete digraphs into strong spanning subdigraphs*, J. Combin. Theory Ser. B **102** (2012) 701–714.
doi:10.1016/j.jctb.2011.09.001
- [5] Y. Guo, *Locally Semicomplete Digraphs* (Ph.D. Thesis, RWTH Aachen University, 1995).
- [6] Y. Guo, *Strongly Hamiltonian-connected locally semicomplect digraphs*, J. Graph Theory **21** (1996) 65–73.
doi:10.1002/(SICI)1097-0118(199605)22:1<65::AID-JGT9>3.0.CO;2-J
- [7] F. Harary and L. Moser, *The theory of round robin tournaments*, Amer. Math. Monthly **73** (1966) 231–246.
doi:10.2307/2315334
- [8] J. Huang, *On the structure of local tournaments*, J. Combin. Theory Ser. B **63** (1995) 200–221.
doi:10.1006/jctb.1995.1016
- [9] S. Li, W. Meng, Y. Guo and G. Xu, *A local tournament contains a vertex whose out-arc are pseudo-girth-pancyclic*, J. Graph Theory **62** (2009) 346–361.
doi:10.1002/jgt.20404
- [10] D. Meierling, *Local tournaments with the minimum number of Hamiltonian cycles or cycles of length three*, Discrete Math. **310** (2010) 1940–1948.
doi:10.1016/j.disc.2010.03.003
- [11] C. Thomassen, *Edge-disjoint Hamiltonian paths and cycles in tournaments*, Proc. Lond. Math. Soc. (3) **45** (1982) 151–168.
doi:10.1112/plms/s3-45.1.151
- [12] R. Wang, A. Yang and S. Wang, *Kings in locally semicomplete digraphs*, J. Graph Theory **63** (2010) 279–283.
doi:10.1002/jgt.20426
- [13] X.H. Zhang, R.J. Li and S.J. Li, *H-force sets of locally semicomplete digraphs*, Discrete Appl. Math. **160** (2012) 2491–2496.
doi:10.1016/j.dam.2012.06.014

Received 17 August 2016

Revised 1 November 2016

Accepted 23 December 2016