IRREDUCIBLE NO-HOLE $L(2, 1)$ -COLORING OF EDGE-MULTIPLICITY-PATHS-REPLACEMENT GRAPH

NIBEDITA MANDAL AND PRATIMA PANIGRAHI

*Department of Mathematics
Indian Institute of Technology Kharagpur, India*

e-mail: nibedita.mandal.iitkgp@gmail.com
pratima@maths.iitkgp.ernet.in

Abstract

An $L(2, 1)$ -coloring (or *labeling*) of a simple connected graph G is a mapping $f : V(G) \rightarrow \mathbb{Z}^+ \cup \{0\}$ such that $|f(u) - f(v)| \geq 2$ for all edges uv of G , and $|f(u) - f(v)| \geq 1$ if u and v are at distance two in G . The *span of an $L(2, 1)$ -coloring f* , denoted by $\text{span}(f)$, of G is $\max\{f(v) : v \in V(G)\}$. The *span of G* , denoted by $\lambda(G)$, is the minimum span of all possible $L(2, 1)$ -colorings of G . For an $L(2, 1)$ -coloring f of a graph G with span k , an integer l is a *hole* in f if $l \in (0, k)$ and there is no vertex v in G such that $f(v) = l$. An $L(2, 1)$ -coloring is a *no-hole coloring* if there is no hole in it, and is an *irreducible coloring* if color of none of the vertices in the graph can be decreased and yield another $L(2, 1)$ -coloring of the same graph. An *irreducible no-hole coloring*, in short *inh-coloring*, of G is an $L(2, 1)$ -coloring of G which is both irreducible and no-hole. For an inh-colorable graph G , the *inh-span* of G , denoted by $\lambda_{inh}(G)$, is defined as $\lambda_{inh}(G) = \min\{\text{span}(f) : f \text{ is an inh-coloring of } G\}$. Given a function $h : E(G) \rightarrow \mathbb{N} - \{1\}$, and a positive integer $r \geq 2$, the *edge-multiplicity-paths-replacement graph $G(rP_h)$* of G is the graph obtained by replacing every edge uv of G with r paths of length $h(uv)$ each. In this paper we show that $G(rP_h)$ is inh-colorable except possibly the cases $h(e) \geq 2$ with equality for at least one but not for all edges e and (i) $\Delta(G) = 2$, $r = 2$ or (ii) $\Delta(G) \geq 3$, $2 \leq r \leq 4$. We find the exact value of $\lambda_{inh}(G(rP_h))$ in several cases and give upper bounds of the same in the remaining. Moreover, we find the value of $\lambda(G(rP_h))$ in most of the cases which were left by Lü and Sun in [*$L(2, 1)$ -labelings of the edge-multiplicity-paths-replacement of a graph*, J. Comb. Optim. 31 (2016) 396–404].

Keywords: $L(2, 1)$ -coloring, no-hole coloring, irreducible coloring, subdivision graph, edge-multiplicity-paths-replacement graph.

2010 Mathematics Subject Classification: 05C15.

REFERENCES

- [1] F.-H. Chang, M.-L. Chia, D. Kuo, S.-C. Liaw and M.-H. Tsai, *$L(2, 1)$ -labelings of subdivisions of graphs*, Discrete Math. **338** (2015) 248–255.
doi:10.1016/j.disc.2014.09.006
- [2] G.J. Chang and C. Lu, *Distance-two labelings of graphs*, European J. Combin. **24** (2003) 53–58.
doi:10.1016/S0195-6698(02)00134-8
- [3] P.C. Fishburn and F.S. Roberts, *No-hole $L(2, 1)$ -colorings*, Discrete Appl. Math. **130** (2003) 513–519.
doi:10.1016/S0166-218X(03)00329-9
- [4] J.R. Griggs and R.K. Yeh, *Labelling graphs with a condition at distance 2*, SIAM J. Discrete Math. **5** (1992) 586–595.
doi:10.1137/0405048
- [5] F. Havet and M.-L. Yu, *$(p, 1)$ -total labelling of graphs*, Technical Report 4650, INRIA (2002).
- [6] F. Havet and M.-L. Yu, *$(p, 1)$ -total labelling of graphs*, Discrete Math. **308** (2008) 496–513.
doi:10.1016/j.disc.2007.03.034
- [7] J. Jacob, R. Laskar and J. Villalpando, *On the irreducible no-hole $L(2, 1)$ coloring of bipartite graphs and Cartesian products*, J. Combin. Math. Combin. Comput. **78** (2011) 49–64.
- [8] N. Karst, J. Oehrlin, D.S. Troxell and J. Zhu, *$L(d, 1)$ -labelings of the edge-path-replacement by factorization of graphs*, J. Comb. Optim. **30** (2015) 34–41.
doi:10.1007/s10878-013-9632-x
- [9] R.C. Laskar, G.L. Matthews, B. Novick and J. Villalpando, *On irreducible no-hole $L(2, 1)$ -coloring of trees*, Networks **53** (2009) 206–211.
doi:10.1002/net.20286
- [10] R.C. Laskar and J.J. Villalpando, *Irreducibility of $L(2, 1)$ -coloring and inh-colorability of unicyclic and hex graphs*, Util. Math. **69** (2006) 65–83.
- [11] D. Lü, *$L(2, 1)$ -labelings of the edge-path-replacement of a graph*, J. Comb. Optim. **26** (2013) 385–392.
doi:10.1007/s10878-012-9470-2
- [12] D. Lü and J. Sun, *$L(2, 1)$ -labelings of the edge-multiplicity-paths-replacement of a graph*, J. Comb. Optim. **31** (2016) 396–404.
doi:10.1007/s10878-014-9761-x
- [13] N. Mandal and P. Panigrahi, *On irreducible no-hole $L(2, 1)$ -coloring of subdivision of graphs*, J. Comb. Optim. **33** (2017) 1421–1442.
doi:10.1007/s10878-016-0047-3
- [14] D.B. West, Introduction to Graph Theory (New Delhi, Prentice-Hall, 2003).

- [15] M.A. Whittlesey, J.P. Georges and D.W. Mauro, *On the λ -number of Q_n and related graphs*, SIAM J. Discrete Math. **8** (1995) 499–506.
doi:10.1137/S0895480192242821

Received 26 August 2016
Revised 10 January 2017
Accepted 10 January 2017