TURÁN FUNCTION AND H -DECOMPOSITION PROBLEM FOR GEM GRAPHS

HENRY LIU

*School of Mathematics and Statistics
Central South University
Changsha 410083, China*

e-mail: henry-liu@csu.edu.cn

AND

TERESA SOUSA

*Escola Naval and Centro de Investigação Naval
Escola Naval - Alfeite
2810-001 Almada, Portugal*

and

*Centro de Matemática e Aplicações
Faculdade de Ciências e Tecnologia
Universidade Nova de Lisboa
Campus de Caparica
2829-516 Caparica, Portugal*

e-mail: teresa.maria.sousa@marinha.pt

Abstract

Given a graph H , the *Turán function* $\text{ex}(n, H)$ is the maximum number of edges in a graph on n vertices not containing H as a subgraph. For two graphs G and H , an H -*decomposition* of G is a partition of the edge set of G such that each part is either a single edge or forms a graph isomorphic to H . Let $\phi(n, H)$ be the smallest number ϕ such that any graph G of order n admits an H -decomposition with at most ϕ parts. Pikhurko and Sousa conjectured that $\phi(n, H) = \text{ex}(n, H)$ for $\chi(H) \geq 3$ and all sufficiently large n . Their conjecture has been verified by Özkahya and Person for all edge-critical graphs H . In this article, we consider the *gem graphs* gem_4 and gem_5 . The graph gem_4 consists of the path P_4 with four vertices a, b, c, d and edges ab, bc, cd plus a universal vertex u adjacent to a, b, c, d , and the graph gem_5 is similarly defined with the path P_5 on five vertices. We determine

the Turán functions $\text{ex}(n, \text{gem}_4)$ and $\text{ex}(n, \text{gem}_5)$, and verify the conjecture of Pikhurko and Sousa when H is the graph gem_4 and gem_5 .

Keywords: gem graph, Turán function, extremal graph, graph decomposition.

2010 Mathematics Subject Classification: 05C35, 05C70.

REFERENCES

- [1] P. Allen, J. Böttcher and Y. Person, *An improved error term for minimum H -decompositions of graphs*, J. Combin. Theory Ser. B **108** (2014) 92–101.
doi:10.1016/j.jctb.2014.03.001
- [2] B. Bollobás, *On complete subgraphs of different orders*, Math. Proc. Cambridge Philos. Soc. **79** (1976) 19–24.
doi:10.1017/S0305004100052063
- [3] D. Dor and M. Tarsi, *Graph decomposition is NP-complete: a complete proof of Holyer’s conjecture*, SIAM J. Comput. **26** (1997) 1166–1187.
doi:10.1137/S0097539792229507
- [4] P. Erdős, A.W. Goodman and L. Pósa, *The representation of a graph by set intersections*, Canad. J. Math. **18** (1966) 106–112.
doi:10.4153/CJM-1966-014-3
- [5] R. Faudree and R. Schelp, *Path Ramsey numbers in multicolorings*, J. Combin. Theory Ser. B **19** (1975) 150–160.
doi:10.1016/0095-8956(75)90080-5
- [6] H. Liu and T. Sousa, *Decompositions of graphs into fans and single edges*, J. Graph Theory **85** (2017) 400–411.
doi:/10.1002/jgt.22069
- [7] L. Özkahya and Y. Person, *Minimum H -decompositions of graphs: edge-critical case*, J. Combin. Theory Ser. B **102** (2012) 715–725.
doi:10.1016/j.jctb.2011.10.004
- [8] O. Pikhurko, *A note on the Turán function of even cycles*, Proc. Amer. Math. Soc. **140** (2012) 3687–3692.
doi:10.1090/S0002-9939-2012-11274-2
- [9] O. Pikhurko and T. Sousa, *Minimum H -decompositions of graphs*, J. Combin. Theory Ser. B **97** (2007) 1041–1055.
doi:10.1016/j.jctb.2007.03.002
- [10] T. Sousa, *Decompositions of graphs into 5-cycles and other small graphs*, Electron. J. Combin. **12** (2005) #R49.
- [11] T. Sousa, *Decompositions of graphs into a given clique-extension*, Ars Combin. **100** (2011) 465–472.

- [12] T. Sousa, *Decompositions of graphs into cycles of length seven and single edges*, Ars Combin. **119** (2015) 321–329.
- [13] P. Turán, *On an extremal problem in graph theory*, Mat. Fiz. Lapok **48** (1941) 436–452.

Received 10 October 2016

Revised 26 January 2017

Accepted 2 February 2017