ANTI-RAMSEY NUMBER OF HANOI GRAPHS

IZOLDA GORGOL AND ANNA LECHOWSKA

*Department of Applied Mathematics
Lublin University of Technology*

e-mail: i.gorgol@pollub.pl
annalech2112@gmail.com

Abstract

Let $\text{ar}(G, H)$ be the largest number of colors such that there exists an edge coloring of G with $\text{ar}(G, H)$ colors such that each subgraph isomorphic to H has at least two edges in the same color. We call $\text{ar}(G, H)$ the *anti-Ramsey number* for a pair of graphs (G, H) . This notion was introduced by Erdős, Simonovits and Sós in 1973 and studied in numerous papers.

Hanoi graphs were introduced by Scorer, Grundy and Smith in 1944 as the model of the well known Tower of Hanoi puzzle.

In the paper we study the anti-Ramsey number of Hanoi graphs and consider them both as the graph G and H . Among others we present the exact value of the anti-Ramsey number in case when both graphs are constructed for the same number of pegs.

Keywords: anti-Ramsey number, rainbow number, Hanoi graph.

2010 Mathematics Subject Classification: 05C30, 05C35, 05C15.

REFERENCES

- [1] N. Alon, *On the conjecture of Erdős, Simonovits and Sós concerning anti-Ramsey theorems*, J. Graph Theory **7** (1983) 91–94.
doi:10.1002/jgt.3190070112
- [2] D. Arett and S. Doree, *Coloring and counting on the Tower of Hanoi graphs*, Math. Mag. **83** (2010) 202–208.
doi:10.4169/002557010X494841
- [3] M. Axenovich, H. Harborth, A. Kemnitz, M. Möller and I. Schiermeyer, *Rainbows in the hypercube*, Graphs Combin. **23** (2007) 123–133.
doi:10.1007/s00373-007-0691-6
- [4] M. Axenovich and T. Jiang, *Anti-Ramsey numbers for small complete bipartite graphs*, Ars Combin. **73** (2004) 311–318.

- [5] M. Axenovich, T. Jiang and A. Kündgen, *Bipartite anti-Ramsey numbers of cycles*, J. Graph Theory **47** (2004) 9–28.
doi:10.1002/jgt.20012
- [6] J.-P. Bode, D. Grimm and A. Kemnitz, *Hypertube-anti-Ramsey numbers of Q_5* , Ars Combin. **110** (2013) 105–111.
- [7] T. Bousch, *La quatrième tour de Hanoi*, Bull. Belg. Math. Soc. Simon Stevin **21** (2014) 895–912.
- [8] R. Diestel, Graph Theory (Springer-Verlag, New York, 1997).
- [9] P. Erdős, A. Simonovits, and V. Sós, *Anti-Ramsey theorems*, in: Infinite and finite sets, A. Hajnal, R. Rado and V. Sós (Ed(s)), (Colloq. Math. Soc. J. Bolyai, North-Holland, 1973) 633–643.
- [10] S. Fujita, A. Kaneko, I. Schiermeyer and K. Suzuki, *A Rainbow k -matching in the complete graph with r colors*, Electron. J. Combin. **16** (2009) #R51.
- [11] S. Fujita, C. Magnant and K. Ozeki, *Rainbow generalizations of Ramsey theory: A survey*, Graphs Combin. **26** (2010) 1–30.
doi:10.1007/s00373-010-0891-3
- [12] I. Gorgol, *Anti-Ramsey numbers in complete split graphs*, Discrete Math. **339** (2016) 1944–1949.
doi:10.1016/j.disc.2015.10.038
- [13] R. Haas and M. Young, *The anti-Ramsey number of perfect matching*, Discrete Math. **312** (2012) 933–937.
doi:10.1016/j.disc.2011.10.017
- [14] A.M. Hinz and D. Parisse, *Coloring Hanoi and Sierpiński graphs*, Discrete Math. **312** (2012) 1521–1535.
doi:10.1016/j.disc.2011.08.019
- [15] A.M. Hinz and D. Parisse, *On the planarity of Hanoi graphs*, Expo. Math. **20** (2002) 264–267.
doi:10.1016/S0723-0869(02)80023-8
- [16] A.M. Hinz, S. Klavžar, U. Milutinović and C. Petr, *The Tower of Hanoi – Myths and Maths* (Birkhäuser/Springer, Basel, 2013).
- [17] M. Hornák, S. Jendrol’, I. Schiermeyer and R. Soták, *Rainbow numbers for cycles in plane triangulations*, J. Graph Theory **78** (2015) 248–257.
doi:10.1002/jgt.21803
- [18] T. Jiang, *Edge-colorings with no large polychromatic stars*, Graphs Combin. **18** (2002) 303–308.
doi:10.1007/s003730200022
- [19] T. Jiang and D.B. West, *On the Erdős-Simonovits-Sós conjecture about the anti-Ramsey number of a cycle*, Combin. Probab. Comput. **12** (2003) 585–598.
doi:10.1017/S096354830300590X

- [20] T. Jiang and D.B. West, *Edge-colorings of complete graphs that avoid polychromatic trees*, Discrete Math. **274** (2004) 137–145.
doi:10.1016/j.disc.2003.09.002
- [21] S. Jendrol', I. Schiermeyer and J. Tu, *Rainbow numbers for matchings in plane triangulations*, Discrete Math. **331** (2014) 158–164.
doi:10.1016/j.disc.2014.05.012
- [22] S. Klavžar, U. Milutinović and C. Petr, *Combinatorics of topmost discs of multi-peg Tower of Hanoi problem*, Ars Combin. **59** (2001) 55–64.
- [23] X. Li, J. Tu and Z. Jin, *Bipartite rainbow numbers of matchings*, Discrete Math. **309** (2009) 2575–2578.
doi:10.1016/j.disc.2008.05.011
- [24] J.J. Montellano-Ballesteros and V. Neuman-Lara, *An anti-Ramsey theorem on cycles*, Graphs Combin. **21** (2005) 343–354.
doi:10.1007/s00373-005-0619-y
- [25] J.J. Montellano-Ballesteros, *On totally multicolored stars*, J. Graph Theory **51** (2006) 225–243.
doi:10.1002/jgt.20140
- [26] R.S. Scorer, P.M. Grundy and C.A.B. Smith, *Some binary games*, Math. Gaz. **28** (1944) 96–103.
doi:10.2307/3606393
- [27] I. Schiermeyer, *Rainbow numbers for matchings and complete graphs*, Discrete Math. **286** (2004) 157–162.
doi:10.1016/j.disc.2003.11.057

Received 21 November 2016

Revised 1 August 2017

Accepted 2 August 2017