NEIGHBOR SUM DISTINGUISHING TOTAL
CHROMATIC NUMBER OF PLANAR GRAPHS
WITHOUT 5-CYCLES¹

XUE ZHAO

AND

CHANG-QING XU²

*School of Science,
Hebei University of Technology
Tianjin 300401, P.R. China*

e-mail: zhaoxhxy@163.com
chqxu@hebut.edu.cn

Abstract

For a given graph $G = (V(G), E(G))$, a proper total coloring $\phi : V(G) \cup E(G) \rightarrow \{1, 2, \dots, k\}$ is neighbor sum distinguishing if $f(u) \neq f(v)$ for each edge $uv \in E(G)$, where $f(v) = \sum_{uv \in E(G)} \phi(uv) + \phi(v)$, $v \in V(G)$. The smallest integer k in such a coloring of G is the neighbor sum distinguishing total chromatic number, denoted by $\chi''_{\Sigma}(G)$. Piłśniak and Woźniak first introduced this coloring and conjectured that $\chi''_{\Sigma}(G) \leq \Delta(G) + 3$ for any graph with maximum degree $\Delta(G)$. In this paper, by using the discharging method, we prove that for any planar graph G without 5-cycles, $\chi''_{\Sigma}(G) \leq \max\{\Delta(G) + 2, 10\}$. The bound $\Delta(G) + 2$ is sharp. Furthermore, we get the exact value of $\chi''_{\Sigma}(G)$ if $\Delta(G) \geq 9$.

Keywords: neighbor sum distinguishing total coloring, discharging method, planar graph.

2010 Mathematics Subject Classification: 05C15.

REFERENCES

- [1] J.A. Bondy and U.S.R. Murty, *Graph Theory with Applications* (North-Holland, NewYork-Amsterdam-Oxford, 1982).

¹This work was supported by NSFC (No.11671232), HNSF(No.A2015202301) and HU STP (No.ZD2015106, QN2017044).

²Corresponding author.

- [2] X. Cheng, D. Huang, G. Wang and J. Wu, *Neighbor sum distinguishing total colorings of planar graphs with maximum degree Δ* , Discrete Appl. Math. **190–191** (2015) 34–41.
doi:10.1016/j.dam.2015.03.013
- [3] A. Dong and G. Wang, *Neighbor sum distinguishing total colorings of graphs with bounded maximum average degree*, Acta Math. Sin. (Engl. Ser.) **30** (2014) 703–709.
doi:10.1007/s10114-014-2454-7
- [4] L. Ding, G. Wang and G. Yan, *Neighbor sum distinguishing total colorings via the Combinatorial Nullstellensatz*, Sci. China Math. **57** (2014) 1875–1882.
doi:10.1007/s11425-014-4796-0
- [5] S. Ge, J. Li and C. Xu, *Neighbor sum distinguishing total chromatic number of planar graphs without 4-cycles*, Util. Math. **105** (2017) 259–265.
- [6] S. Ge, J. Li and C. Xu, *Neighbor sum distinguishing total coloring of planar graphs without 5-cycles*, Theoret. Comput. Sci. **689** (2017) 169–175.
doi:10.1016/j.tcs.2017.05.037
- [7] H. Li, L. Ding, B. Liu and G. Wang, *Neighbor sum distinguishing total colorings of planar graphs*, J. Comb. Optim. **30** (2015) 675–688.
doi:10.1007/s10878-013-9660-6
- [8] J. Li, S. Ge and C. Xu, *Neighbor sum distinguishing total colorings of planar graphs with girth at least 5*, Util. Math. **104** (2017) 115–121.
- [9] H. Li, B. Liu and G. Wang, *Neighbor sum distinguishing total colorings of K_4 -minor free graphs*, Front. Math. China **8** (2013) 1351–1366.
doi:10.1007/s11464-013-0322-x
- [10] Q. Ma, J. Wang and H. Zhao, *Neighbor sum distinguishing total colorings of planar graphs without short cycles*, Util. Math. **98** (2015) 349–359.
- [11] M. Piłśniak and M. Woźniak, *On the total-neighbor-distinguishing index by sums*, Graphs Combin. **31** (2015) 771–782.
doi:10.1007/s00373-013-1399-4
- [12] C. Qu, G. Wang, J. Wu and X. Yu, *On the neighbor sum distinguishing total coloring of planar graphs*, Theoret. Comput. Sci. **609** (2016) 162–170.
doi:10.1016/j.tcs.2015.09.017
- [13] C. Qu, G. Wang, G. Yan and X. Yu, *Neighbor sum distinguishing total choosability of planar graphs*, J. Comb. Optim. **32** (2016) 906–916.
doi:10.1007/s10878-015-9911-9
- [14] H. Song, W. Pan, X. Gong and C. Xu, *A note on the neighbor sum distinguishing total coloring of planar graphs*, Theoret. Comput. Sci. **640** (2016) 125–129.
doi:10.1016/j.tcs.2016.06.007
- [15] H. Song and C. Xu, *Neighbor sum distinguishing total chromatic number of K_4 -minor free graph*, Front. Math. China **12** (2017) 937–947.
doi:10.1007/s11464-017-0649-9

- [16] H. Song and C. Xu, *Neighbor sum distinguishing total coloring of planar graphs without 4-cycles*, J. Comb. Optim. **34** (2017) 1147–1158.
doi:10.1007/s10878-017-0137-x
- [17] J. Wang, J. Cai and Q. Ma, *Neighbor sum distinguishing total choosability of planar graphs without 4-cycles*, Discrete Appl. Math. **206** (2016) 215–219.
doi:10.1016/j.dam.2016.02.003
- [18] J. Wang, J. Cai and B. Qiu, *Neighbor sum distinguishing total choosability of planar graphs without adjacent triangles*, Theoret. Comput. Sci. **661** (2017) 1–7.
doi:10.1016/j.tcs.2016.11.003
- [19] J. Wang, Q. Ma and X. Han, *Neighbor sum distinguishing total colorings of triangle free planar graphs*, Acta Math. Sin. (Engl. Ser.) **31** (2015) 216–224.
doi:10.1007/s10114-015-4114-y
- [20] D. Yang, X. Yu, L. Sun, J. Wu and S. Zhou, *Neighbor sum distinguishing total chromatic number of planar graphs with maximum degree 10*, Appl. Math. Comput. **314** (2017) 456–468.
doi:10.1016/j.amc.2017.06.002
- [21] J. Yao, X. Yu, G. Wang and C. Xu, *Neighbor sum distinguishing total coloring of 2-degenerate graphs*, J. Comb. Optim. **34** (2017) 64–70.
doi:10.1007/s10878-016-0053-5
- [22] J. Yao, X. Yu, G. Wang and C. Xu, *Neighbor sum (set) distinguishing total choosability of d -degenerate graphs*, Graphs Combin. **32** (2016) 1611–1620.
doi:10.1007/s00373-015-1646-y

Received 5 December 2017

Revised 8 February 2018

Accepted 7 March 2018