ON FACTORABLE BIGRAPHIC PAIRS¹

JIAN-HUA YIN AND SHA-SHA LI

*Department of Mathematics
College of Information Science and Technology
Hainan University, Haikou 570228, P.R. China*

e-mail: yinhj@hainu.edu.cn

Abstract

Let $S = (a_1, \dots, a_m; b_1, \dots, b_n)$, where $a_1, \dots, a_m$ and $b_1, \dots, b_n$ are two sequences of nonnegative integers. We say that S is a *bigraphic pair* if there exists a simple bipartite graph G with partite sets $\{x_1, x_2, \dots, x_m\}$ and $\{y_1, y_2, \dots, y_n\}$ such that $d_G(x_i) = a_i$ for $1 \leq i \leq m$ and $d_G(y_j) = b_j$ for $1 \leq j \leq n$. In this case, we say that G is a *realization* of S . Analogous to Kundu's k -factor theorem, we show that if $(a_1, a_2, \dots, a_m; b_1, b_2, \dots, b_n)$ and $(a_1 - e_1, a_2 - e_2, \dots, a_m - e_m; b_1 - f_1, b_2 - f_2, \dots, b_n - f_n)$ are two bigraphic pairs satisfying $k \leq f_i \leq k + 1$, $1 \leq i \leq n$ (or $k \leq e_i \leq k + 1$, $1 \leq i \leq m$), for some $0 \leq k \leq m - 1$ (or $0 \leq k \leq n - 1$), then $(a_1, a_2, \dots, a_m; b_1, b_2, \dots, b_n)$ has a realization containing an $(e_1, e_2, \dots, e_m; f_1, f_2, \dots, f_n)$ -factor. For $m = n$, we also give a necessary and sufficient condition for an $(k^n; k^n)$ -factorable bigraphic pair to be connected $(k^n; k^n)$ -factorable when $k \geq 2$. This implies a characterization of bigraphic pairs with a realization containing a Hamiltonian cycle.

Keywords: degree sequence, bigraphic pair, Hamiltonian cycle.

2010 Mathematics Subject Classification: 05C07.

REFERENCES

- [1] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications (American Elsevier, New York, 1976).
- [2] Y. Chen, *A short proof of Kundu's k -factor theorem*, Discrete Math. **71** (1988) 177–179.
doi:10.1016/0012-365X(88)90070-2

¹Supported by National Natural Science Foundation of China (Grant No. 11561017) and Hainan Provincial Natural Science Foundation of China (Grant No. 2016CXTD004).

- [3] D. Gale, *A theorem on flows in networks*, Pacific J. Math. **7** (1957) 1073–1082.
doi:10.2140/pjm.1957.7.1073
- [4] D.J. Kleitman and D.L. Wang, *Algorithm for constructing graphs and digraphs with given valences and factors*, Discrete Math. **6** (1973) 79–88.
doi:10.1016/0012-365X(73)90037-X
- [5] S. Kundu, *The k -factor conjecture is true*, Discrete Math. **6** (1973) 367–376.
doi:10.1016/0012-365X(73)90068-X
- [6] S. Kundu, *Generalizations of the k -factor theorem*, Discrete Math. **9** (1974) 173–179.
doi:10.1016/0012-365X(74)90147-2
- [7] A.R. Rao and S.B. Rao, *On factorable degree sequences*, J. Combin. Theory Ser. B **13** (1972) 185–191.
doi:10.1016/0095-8956(72)90055-X
- [8] H.J. Ryser, *Combinatorial properties of matrices of zeros and ones*, Canad. J. Math. **9** (1957) 371–377.
doi:10.4153/CJM-1957-044-3
- [9] J.H. Yin, *A characterization for a graphic sequence to be potentially C_r -graphic*, Sci. China Math. **53** (2010) 2893–2905.
doi:10.1007/s11425-010-3124-6

Received 15 September 2017

Revised 5 May 2018

Accepted 5 May 2018