TOTAL 2-RAINBOW DOMINATION NUMBERS OF TREES

H. ABDOLLAHZADEH AHANGAR¹, J. AMJADI², M. CHELLALI³,
S. NAZARI-MOGHADDAM² AND S.M. SHEIKHOLESAMI²

¹*Department of Mathematics
Babol Noshirvani University of Technology
Babol, I.R. Iran*

e-mail: ha.ahangar@nit.ac.ir

²*Department of Mathematics
Azarbaijan Shahid Madani University
Tabriz, I.R. Iran*

e-mail: j-amjadi;s.nazari;s.m.sheikholeslami@azaruniv.ac.ir

³*LAMDA-RO Laboratory, Department of Mathematics
University of Blida
B.P. 270, Blida, Algeria*

e-mail: m_chellali@yahoo.com

Abstract

A 2-rainbow dominating function (2RDF) of a graph $G = (V(G), E(G))$ is a function f from the vertex set $V(G)$ to the set of all subsets of the set $\{1, 2\}$ such that for every vertex $v \in V(G)$ with $f(v) = \emptyset$ the condition $\bigcup_{u \in N(v)} f(u) = \{1, 2\}$ is fulfilled, where $N(v)$ is the open neighborhood of v . A total 2-rainbow dominating function f of a graph with no isolated vertices is a 2RDF with the additional condition that the subgraph of G induced by $\{v \in V(G) \mid f(v) \neq \emptyset\}$ has no isolated vertex. The total 2-rainbow domination number, $\gamma_{tr2}(G)$, is the minimum weight of a total 2-rainbow dominating function of G . In this paper, we establish some sharp upper and lower bounds on the total 2-rainbow domination number of a tree. Moreover, we show that the decision problem associated with $\gamma_{tr2}(G)$ is NP-complete for bipartite and chordal graphs.

Keywords: 2-rainbow dominating function, 2-rainbow domination number, total 2-rainbow dominating function, total 2-rainbow domination number.

2010 Mathematics Subject Classification: 05C69.

REFERENCES

- [1] H. Abdollahzadeh Ahangar, J. Amjadi, N. Jafari Rad and V. Samodivkin, *Total k -rainbow domination in graphs*, Comm. Combin. Optim. **3** (2018) 37–50.
doi:10.22049/CCO.2018.25719.1021
- [2] H. Abdollahzadeh Ahangar, M.A. Henning, V. Samodivkin and I.G. Yero, *Total Roman domination in graphs*, Appl. Anal. Discrete Math. **10** (2016) 501–517.
doi:10.2298/AADM160802017A
- [3] H. Abdollahzadeh Ahangar, M. Khaibari, N. Jafari Rad and S.M. Sheikholeslami, *Graphs with large total 2-rainbow domination number*, Iran. J. Sci. Technol. Trans. A Sci. **42** (2018) 841–846.
doi:10.1007/s40995-017-0465-9
- [4] G.J. Chang, J. Wu and X. Zhu, *Rainbow domination on trees*, Discrete Appl. Math. **158** (2010) 8–12.
doi:10.1016/j.dam.2009.08.010
- [5] M. Chellali and N. Jafari Rad, *On 2-Rainbow domination and Roman domination in graphs*, Australas. J. Combin. **56** (2013) 85–93.
- [6] B. Brešar, M.A. Henning and D.F. Rall, *Rainbow domination in graphs*, Taiwanese J. Math. **12** (2008) 201–213.
- [7] B. Brešar and T.K. Sumenjak, *On the 2-rainbow domination in graphs*, Discrete Appl. Math. **155** (2007) 2394–2400.
doi:10.1016/j.dam.2007.07.018
- [8] N. Dehgard, S.M. Sheikholeslami and L. Volkmann, *The rainbow domination subdivision numbers of graphs*, Mat. Vesnik **67** (2015) 102–114.
- [9] E. DeLaVina, C.E. Larson, R. Pepper and B. Waller, *On total domination and support vertices of a tree*, AKCE Int. J. Graphs Comb. **7** (2010) 85–95.
- [10] M. Falahat, S.M. Sheikholeslami and L. Volkmann, *New bounds on the rainbow domination subdivision number*, Filomat **28** (2014) 615–622.
doi:10.2298/FIL1403615F
- [11] C.-H. Liu and G.J. Chang, *Roman domination on strongly chordal graphs*, J. Comb. Optim. **26** (2013) 608–619.
doi:10.1007/s10878-012-9482-y
- [12] Y. Wu and N. Jafari Rad, *Bounds on the 2-Rainbow domination number of graphs*, Graphs Combin. **29** (2013) 1125–1133.
doi:10.1007/s00373-012-1158-y
- [13] Y. Wu and H. Xing, *Note on 2-rainbow domination and Roman domination in graphs*, Appl. Math. Lett. **23** (2010) 706–709.
doi:10.1016/j.aml.2010.02.012

Received 11 April 2018

Revised 22 October 2018

Accepted 3 November 2018