ON THE STAR CHROMATIC INDEX OF GENERALIZED PETERSEN GRAPHS

ENQIANG ZHU^{1,2} AND ZEHUI SHAO

*Institute of Computing Science and Technology
Guangzhou University, Guangzhou 510006, China*

e-mail: zhuenqiang@gzhu.edu.cn

Abstract

The star k -edge-coloring of graph G is a proper edge coloring using k colors such that no path or cycle of length four is bichromatic. The minimum number k for which G admits a star k -edge-coloring is called the star chromatic index of G , denoted by $\chi'_s(G)$. Let $\text{GCD}(n, k)$ be the greatest common divisor of n and k . In this paper, we give a necessary and sufficient condition of $\chi'_s(P(n, k)) = 4$ for a generalized Petersen graph $P(n, k)$ and show that “almost all” generalized Petersen graphs have a star 5-edge-colorings. Furthermore, for any two integers k and n ($\geq 2k + 1$) such that $\text{GCD}(n, k) \geq 3$, $P(n, k)$ has a star 5-edge-coloring, with the exception of the case that $\text{GCD}(n, k) = 3$, $k \neq \text{GCD}(n, k)$ and $\frac{n}{3} \equiv 1 \pmod{3}$.

Keywords: star edge-coloring, star chromatic index, generalized Petersen graph.

2010 Mathematics Subject Classification: 05C15.

REFERENCES

- [1] M.O. Albertson, G.G. Chappell, H.A. Kierstead, A. Kündgen and R. Ramamurthi, *Coloring with no 2-colored P_4 's*, Electron. J. Combin. **11** (2004) #R26.
- [2] L. Bezegová, B. Lužar, M. Mockovčiová, R. Soták and R. Škrekovski, *Star edge coloring of some classes of graphs*, J. Graph Theory **81** (2016) 73–82.
doi:10.1002/jgt.21862
- [3] J.A. Bondy and U.S.R. Murty, Graph Theory (Springer, New York, 2008).

¹This work was supported by the National Natural Science Foundation of China under Grants 61672051, 61872101, and the Natural Science Foundation of Guangdong Province under Grant 2018A0303130115.

²Corresponding author.

- [4] Y. Bu, D.W. Cranston, M. Montassier, A. Raspaud and W. Wang, *Star-coloring of sparse graphs*, J. Graph Theory **62** (2009) 201–209.
doi:10.1002/jgt.20392
- [5] M. Chen, A. Raspaud and W. Wang, *6-star-coloring of subcubic graphs*, J. Graph Theory **72** (2013) 128–145.
doi:10.1002/jgt.21636
- [6] Z. Dvořák, B. Mohar and R. Šámal, *Star chromatic index*, J. Graph Theory **72** (2013) 313–326.
doi:10.1002/jgt.21644
- [7] H.A. Kierstead, A. Kündgen and C. Timmons, *Star coloring bipartite planar graphs*, J. Graph Theory **60** (2009) 1–10.
doi:10.1002/jgt.20342
- [8] X.S. Liu and K. Deng, *An upper bound on the star chromatic index of graphs with $\Delta \geq 7$* , J. Lanzhou Univ. Nat. Sci. **44** (2008) 94–95.
- [9] M.E. Watkins, *A theorem on tait colorings with an application to the generalized Petersen graphs*, J. Combin. Theory **6** (1969) 152–164.
doi:10.1016/S0021-9800(69)80116-X

Received 24 October 2017
 Revised 27 November 2018
 Accepted 28 November 2018