SIGNED COMPLETE GRAPHS WITH MAXIMUM INDEX

SAIEED AKBARI

Department of Mathematical Science
Sharif University of Technology, Tehran, Iran

e-mail: s_akbari@sharif.edu

SOUDABEH DALVANDI, FARIDEH HEYDARI

AND

MOHAMMAD MAGHASEDI

Department of Mathematics
Karaj Branch, Islamic Azad University, Karaj, Iran

e-mail: s.dalvandi@kiau.ac.ir
f.heydari@kiau.ac.ir
maghasedi@kiau.ac.ir

In the Memorial of Professor Slobodan Simić

Abstract

Let $\Gamma = (G, \sigma)$ be a signed graph, where G is the underlying simple graph and $\sigma : E(G) \rightarrow \{-, +\}$ is the sign function on the edges of G . The adjacency matrix of a signed graph has -1 or $+1$ for adjacent vertices, depending on the sign of the edges. It was conjectured that if Γ is a signed complete graph of order n with k negative edges, $k < n - 1$ and Γ has maximum index, then negative edges form $K_{1,k}$. In this paper, we prove this conjecture if we confine ourselves to all signed complete graphs of order n whose negative edges form a tree of order $k + 1$. A $[1, 2]$ -subgraph of G is a graph whose components are paths and cycles. Let Γ be a signed complete graph whose negative edges form a $[1, 2]$ -subgraph. We show that the eigenvalues of Γ satisfy the following inequalities:

$$-5 \leq \lambda_n \leq \dots \leq \lambda_2 \leq 3.$$

Keywords: signed graph, complete graph, index.

2010 Mathematics Subject Classification: 05C22, 05C50.

REFERENCES

- [1] N. Abreu, D.M. Cardoso, F.A.M. Franca and C.T.M. Vinagre, *On main eigenvalues of certain graphs*, Comput. Appl. Math., to appear.
- [2] S. Akbari, S. Dalvandi, F. Heydari and M. Maghasedi, *On the eigenvalues of signed complete graphs*, Linear Multilinear Algebra **67** (2019) 433–441.
doi:10.1080/03081087.2017.1403548
- [3] S. Akbari, S. Dalvandi, F. Heydari and M. Maghasedi, *On the multiplicity of -1 and 1 in signed complete graphs*, Util. Math., to appear.
- [4] F. Belardo, P. Petecki and J. Wang, *On signed graphs whose second largest Laplacian eigenvalue does not exceed 3*, Linear Multilinear Algebra **64** (2016) 1785–1799.
doi:10.1080/03081087.2015.1120701
- [5] A.E. Brouwer and W.H. Haemers, Spectra of Graphs (Springer, 2011).
doi:10.1007/978-1-4614-1939-6
- [6] D. Cvetković, P. Rowlinson and S.K. Simić, An Introduction to the Theory of Graph Spectra (Cambridge University Press, Cambridge, 2010).
- [7] F. Harary, *On the notion of balance in a signed graph*, Michigan Math. J. **2** (1953) 143–146.
doi:10.1307/mmj/1028989917
- [8] F. Heider, *Attitude and cognitive organization*, J. Psychology **21** (1946) 107–112.
doi:10.1080/00223980.1946.9917275
- [9] T. Koledin and Z. Stanić, *Connected signed graphs of fixed order, size, and number of negative edges with maximal index*, Linear Multilinear Algebra **65** (2017) 2187–2198.
doi:10.1080/03081087.2016.1265480
- [10] H. Lin, R. Liu and J. Shu, *Some results on the largest and least eigenvalues of graphs*, Electron. J. Linear Algebra **27** (2014) 670–682.
doi:/10.13001/1081-3810.1928
- [11] S.M. Lv, L. Wei and H.X. Zhao, *On the Seidel integral complete multipartite graphs*, Acta Math. Appl. Sin. Engl. Ser. **28** (2012) 705–710.
doi:10.1007/s10255-012-0126-x
- [12] Z. Stanić, *Perturbations in a signed graph and its index*, Discuss. Math. Graph Theory **38** (2018) 841–852.
doi:10.7151/dmgt.2035

Received 4 June 2019
Revised 26 November 2019
Accepted 26 November 2019