COLORINGS OF PLANE GRAPHS WITHOUT LONG MONOCHROMATIC FACIAL PATHS

JÚLIUS CZAP

Department of Applied Mathematics and Business Informatics
Faculty of Economics, Technical University of Košice
Němcovej 32, 04001 Košice, Slovakia

e-mail: julius.czap@tuke.sk

IGOR FABRICI AND STANISLAV JENDROL'

Institute of Mathematics, P.J. Šafárik University
Jesenná 5, 04001 Košice, Slovakia

e-mail: igor.fabrici@upjs.sk
stanislav.jendrol@upjs.sk

Abstract

Let G be a plane graph. A facial path of G is a subpath of the boundary walk of a face of G . We prove that each plane graph admits a 3-coloring (a 2-coloring) such that every monochromatic facial path has at most 3 vertices (at most 4 vertices). These results are in a contrast with the results of Chartrand, Geller, Hedetniemi (1968) and Axenovich, Ueckerdt, Weiner (2017) which state that for any positive integer t there exists a 4-colorable (a 3-colorable) plane graph G_t such that in any its 3-coloring (2-coloring) there is a monochromatic path of length at least t . We also prove that every plane graph is 2-list-colorable in such a way that every monochromatic facial path has at most 4 vertices.

Keywords: plane graph, facial path, vertex-coloring.

2010 Mathematics Subject Classification: 05C10, 05C15.

REFERENCES

- [1] K. Appel and W. Haken, *Every planar map is four colorable*, Bull. Amer. Math. Soc. **82** (1976) 711–712.
doi:10.1090/s0002-9904-1976-14122-5

- [2] M. Axenovich, T. Ueckerdt and P. Weiner, *Splitting planar graphs of girth 6 into two linear forests with short paths*, J. Graph Theory **85** (2017) 601–618.
doi:10.1002/jgt.22093
- [3] J.A. Bondy and U.S.R. Murty, Graph Theory (Springer, 2008).
- [4] O.V. Borodin, *A new proof of the 6-color theorem*, J. Graph Theory **19** (1995) 507–521.
doi:10.1002/jgt.3190190406
- [5] O.V. Borodin, A. Kostochka and M. Yancey, *On 1-improper 2-coloring of sparse graphs*, Discrete Math. **313** (2013) 2638–2649.
doi:10.1016/j.disc.2013.07.014
- [6] H. Broersma, F.V. Fomin, J. Kratochvíl and G.J. Woeginger, *Planar graph coloring avoiding monochromatic subgraphs: trees and paths make it difficult*, Algorithmica **44** (2006) 343–361.
doi:10.1007/s00453-005-1176-8
- [7] G. Chartrand, D.P. Geller and S.T. Hedetniemi, *A generalization of the chromatic number*, Math. Proc. Cambridge Philos. Soc. **64** (1968) 265–271.
doi:10.1017/s0305004100042808
- [8] G. Chartrand, D.P. Geller and S. Hedetniemi, *Graphs with forbidden subgraphs*, J. Combin. Theory Ser. B **10** (1971) 12–41.
doi:10.1016/0095-8956(71)90065-7
- [9] L. Cowen, W. Goddard and C.E. Jesurum, *Defective coloring revisited*, J. Graph Theory **24** (1997) 205–219.
doi:10.1002/(sici)1097-0118(199703)24:3<205::aid-jgt2>3.0.co;2-t
- [10] J. Czap, S. Jendroľ and J. Valiska, *Worm colorings of planar graphs*, Discuss. Math. Graph Theory **37** (2017) 353–368.
doi:10.7151/dmgt.1921
- [11] W. Goddard, *Acyclic colorings of planar graphs*, Discrete Math. **91** (1991) 91–94.
doi:10.1016/0012-365x(91)90166-y
- [12] J.L. Gross and T.W. Tucker, Topological Graph Theory (Dover Publications, 2001).
- [13] H. Grötzsch, *Ein Dreifarbensatz für dreikreisfreie Netze auf der Kugel*, Wiss. Z. Martin-Luther-Universität, Halle-Wittenberg, Math.-Nat. Reihe **8** (1959) 109–120.
- [14] L. Lovász, *On decomposition of graphs*, Studia Sci. Math. Hungar. **1** (1966) 237–238.
- [15] K.S. Poh, *On the linear vertex-arboricity of a planar graph*, J. Graph Theory **14** (1990) 73–75.
doi:10.1002/jgt.3190140108
- [16] C. Thomassen, *2-list-coloring planar graphs without monochromatic triangles*, J. Combin. Theory Ser. B **98** (2008) 1337–1348.
doi:10.1016/j.jctb.2008.02.006

- [17] Y. Wang and L. Xu, *Improper choosability of planar graphs without 4-cycles*, SIAM J. Discrete Math. **27** (2013) 2029–2037.
doi:10.1137/120885140
- [18] Y. Wang and J. Xu, *Decomposing a planar graph without cycles of length 5 into a matching and a 3-colorable graph*, European J. Combin. **43** (2015) 98–123.
doi:10.1016/j.ejc.2014.08.020

Received 26 September 2019

Revised 10 March 2020

Accepted 10 March 2020