From a clinical case to a review of literature: Struma ovarii associated with ascites and high serum CA125

Corina-Elena Minciuna^{1,2}, Stefan Tudor^{1,2}, Vlad Herlea^{3,4}, Catalin Vasilescu^{1,2}

¹General Surgery Department, Fundeni Clinical Institute, Bucharest, Romania

²Carol Davila University of Medicine and Pharmacy, Bucharest, Romania

³Department of Pathology, Fundeni Clinical Institute, Bucharest, Romania

⁴Faculty of Medicine, "Titu Maiorescu" University, Bucharest, Romania

Abstract

Struma ovarii is a rare dermoid tumor of the ovary (1% of all ovarian tumors and 2,7% of ovarian teratomas), with more than 50% of the overall tissue being thyroid. Approximately one third of the cases present ascites. Pseudo-Meigs syndrome has been reported in fewer than 10 cases and 5-8% present thyroid hyperfunction. However, only 26 cases reported the association of ascites or pseudoMeigs' syndrome, elevated cancer antigen 125 and benign struma ovarii in the literature. The authors submit a case of benign struma ovarii that presented as a pelvic mass with ascites and highly elevated CA 125 serum levels and an extensive literature survey on this topic.

Keywords: Struma ovarii, CA125, ascites, ovarian tumor

Introduction

First described in 1899 by Boettlin¹, Struma ovarii represents a rare dermoid tumor of the ovary (1% of all ovarian tumors and 2,7% of ovarian teratomas²), with more than 50% of the overall tissue being thyroid³. Typically occurs between the ages of 40 -60 and rarely occurs before puberty.⁴ Approximately one third of the cases present ascites; pseudo-Meigs syndrome has been reported in fewer than 10 cases and 5-8% present thyroid hyperfunction.⁴ However, only 27 cases reported the association of ascites or pseudo Meigs' syndrome, elevated cancer antigen 125 and struma ovarii in the literature.

The authors submit a case of benign struma ovarii that presented with a pelvic mass, ascites and highly elevated CA 125 serum levels.

Case presentation

A 42-years old female patient with no prior medical history presented to Fundeni Clinical Institute surgical department with gross abdominal distension and diffuse abdominal pain that started 3 months ago and had gradually getting worse.

Physical examination reveals an overweight (BMI 29.69) otherwise healthy-looking patient with gross abdominal distension with dullness to percussion and no abdominal tenderness, hepato-splenomegaly, or palpable abdominal mass. She reports normal bowel movement and physiological mictions. External pelvic examination is normal, and the internal pelvic examination shows normal vaginal walls and cervix. Bimanual palpation exposes a mass that compresses the left lateral and posterior fornices and is fixed on the adjacent planes. Digital rectal examination reveals a left-anterior external compression but otherwise normal. No other pathological findings were found.

Corresponding author: Stefan Tudor, General Surgery Department, Fundeni Clinical Institute, 258 Fundeni Street, Bucharest, Romania, E-mail: stefantudor@gmail.com

Received: January 10, 2021; **Accepted:** February 20, 2021; **Published:** Mars 05, 2021

Citation: Minciuna Corina-Elena, Tudor St. Herlea, VI, Vasilescu C. From a clinical case to a review of literature: Struma ovarii associated with ascites and high serum CA125. Journal of Surgery [Jurnalul de chirurgie]. 2021; 17(1): 37 - 45
DOI: [10.22551/JSURG.2021.01.05](https://doi.org/10.22551/JSURG.2021.01.05)

Copyright: Corina-Elena Minciuna © 2021. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Management

Initial laboratory workup was normal, except CA125 that was 1700 UI/ml with normal range of 0-35 UI/ml.

An abdominal ultrasound showed grade III ascites and tumoral left ovarian cystic mass with heterogeneous tissue structure.

A CT scan of the thorax, abdomen and pelvis was performed revealing the left ovarian cystic mass with approximately 60 mm diameter, and heterogenous structure through the presence of tissular areas. The tumor had blurred margins and was in contact with the middle rectum. Grade III ascites was confirmed with no peritoneal carcinomatosis. The right ovary was of normal aspect and no additional tumors were detected.

A colonoscopy showed extrinsic compression on the middle rectum with normal mucosa, otherwise normal up to the caecum.

The case was discussed with the hospital tumor board and it was decided to carry out surgery in order to perform a biopsy and frozen section histopathological examination depending on which the surgery will be adapted. Surgery associated risks were presented to the patient and a written informed consent was obtained.

Intraoperatively, approximately 6 liters of clear ascites had been collected not before some samples being obtained in order to perform cytology and microbiology examination. At the abdominal cavity inspection, a large cystic left ovarian tumor was observed that had no rectal involvement and was adherent to the anterior wall of the recto-uterine pouch. No other macroscopic or palpatory pathological findings were noted. A left oophorectomy was performed, and the surgical specimen was sent to frozen section histopathological examination. The cytology of the ascites was negative for malignancy and the frozen section histopathological examination

revealed: the resected specimen - ovary was enlarged 7/6/6 centimeters with lobulated aspect on cut surface with cystic area and compact with soft green-brown tissue; microscopically there were cellular regions composed of numerous microfollicular acini, sertoliform tubules as solid areas in which the cells have eosinophilic cytoplasm. Those areas were separated by oedematous and fibromatous stroma (Fig.1). No further surgical intervention was performed.

Post-operative recovery was uneventful; the patient was discharged 5 days after.

Outcome

At one-month follow-up, the surgical wound was healed, and the patient had no ascites at the abdominal ultrasound. The laboratory workup was normal with CA125 level of 20 UI/ml. The immunohistochemistry was performed and the tumoral cells were positive for CK7 (Fig.2), TTF1 (Fig.3), PAX8, negative for CK20, Chromogranin, Synaptophysin, CDX2, and Ki67 (Fig.4) was 5-7% positive nuclear staining in tumoral cells.

The final histopathological result and the immunohistochemistry result concur the diagnosis of benign struma ovarii. Further laboratory workup showed a normal serum-level of thyroid hormones with a TSH (thyrotropin/ thyroid stimulating hormone) level of 6.85 μ UI/ml (normal range 0.27-4.2 μ UI/ml), being detected with subclinical hypothyroidism. The patient was referred to the endocrinologist and entered a follow-up program for early detection of overt hypothyroidism.

At six-month follow-up, the patient had no other symptoms, an abdominal ultrasound was performed, with no additional findings. No additional thyroid-hormone substitution was needed, given the normalization of the TSH level.

Figure 1. HE, x100, struma ovarii, with microfollicular and solid areas

Figure 2. Immunohistochemical stain, CK7 positive tumoral cells

Figure 3. Immunohistochemical stain, TTF1 positive tumoral cells

Figure 4. Immunohistochemical stain, Ki 67 5-7% nuclear positive in tumoral cells

Literature Review

A computerized advanced search was performed on PubMed (Public/ Publisher MEDLINE) electronic database using the following criteria: “struma ovarii” & “ascites” or “pseudo Meigs’ syndrome” & “CA125” on 25 February 2020 and identified 31 case reports/ literature reviews. Search was not limited by publication date but was restricted to English literature. Nevertheless, 4 case reports didn’t associate elevated CA125 with struma ovarii and ascites or pseudo Meigs syndrome and were excluded.

The results are reported in Table 1.

Table 1. Literature review of cases of struma ovarii with ascites or pseudo-Meigs’ syndrome and elevated CA 125 levels

Author	Year	No. of patients	Age	Site (left/ right)	Presenting Symptoms	Ascites Volume (ml)	Tumor size (cm)	CA125 (µg/ml)	Coexisting thyroid disease	Treatment
Jotkowitz and Gee ⁵	1993	1	79	NA	Abdominal distension	Several liters	NA	4670	Absent	TAH, BSO, omentectomy& peritoneal biopsies (frozen section performed)
Leung and Hammond ⁶	1993	2	60	R	Nonspecific symptoms	500	10	224	Absent	TAH, BSO and omentectomy (frozen section performed)
			77	L	Abdominal distension	4000	8x10	2860	Absent	TAH, BSO & left pelvic node sampling (frozen section performed)

Struma ovarii associated with ascites and high serum CA125

Bethune et al. ⁷	1996	1	62	R	Dyspnea, abdominal distension	3500	9x5x5	1621	Absent	TAH,BSO& infracolic omentectomy
Mancuso et al. ⁸	2001	1	31	L	Lower abdominal pain	300	10x9	689	Absent	LSO (frozen section performed)
Long et al. ⁹	2001	2	53	L	Abdominal distension	4100	15x11x7	540	Absent	TAH, BSO & infracolic omentectomy (frozen section performed)
			78	L	Abdominal distension, cachexia	NA	12,2x10x5,2	124.9	Absent	TAH& BSO(frozen section performed)
Huh et al. ¹⁰	2002	1	65	R	Abdominal distension, dyspnea	20000	5x4x4	402	Hypothyroidism	TAH, BSO, appendectomy and omental biopsy (frozen section performed)
Loizzi et al. ¹¹	2002	1	83	R	Abdominal pain, distension	3000	10x7x6,5	1570	Absent	TAH &BSO (frozen section performed)
Bokhari et al. ¹²	2003	1	51	R	Abdominal distension	NA	15x6,5x11	1160	Absent	TAH,BSO& apendectomy (frozen section performed)
Zannoni et al. ¹³	2004	1	66	L	<u>Abdominal distension</u>	<u>5500</u>	<u>9,5x5,6x7</u>	<u>1636</u>	<u>Absent</u>	<u>TAH,BSO, omentectomy, sampling of pelvic lymph nodes , peritoneal biopsy (frozen section performed)</u>
Rim et al. ¹⁴	2005	1	50	L	Abdominal distension	3000	4x4	878,67	Absent	TAH &BSO (frozen section performed)
Guida et al. ¹⁵	2005	1	42	R	Abdominal distension	4000	9,1x7,7	2548	Graves disease	TAH &BSO
Loizzi et al. ¹⁶	2005	1	65	R	Dyspnea, abdominal pain	Few liters	12x7	161	hyperthyroidism	RSO
Uehara and Sawada ¹⁷	2007	1	67	L	Abdominal distension, dyspnea	NA	7x7	2086	Absent	TAH &BSO

Obeidat et al. ¹⁸	2007	1	52	R	Abdominal distension, dyspnea	4000	10x15x8	149	Absent	TAH, BSO & omentectomy
Paladini et al. ²⁰	2008	1	42	R	Abdominal distension, cachexia	8000	11x7,3x8	2548	Basedow syndrom unresponsive to medication- total thyroidectomy=> persistent hyperthyroidism	RSO (frozen section performed)
Mui et al. ²¹	2009	1	56	L	Abdominal distension	8210	6x5x4	5218	Nontoxic multinodular goiter	TAH,BSO& omentectomy (frozen section performed)
Rana et al. ²²	2009	1	70	R&L	Abdominal distension, dyspnea	NA	L 7,5x5,5x4 R 2x1,5x1	284	Absent	TAH,BSO & partial omentectomy
Jiang et al. ²³	2010	1	46	R	Abdominal distension, fatigue	6000	20x18x15	1230	Absent	TAH&BSO (frozen section performed)
Obeidat et al. ²⁴	2012	1	55	L	Abdominal distension	6142 /day	14x15	872	Absent	LSO (frozen section performed)
Mostaghel et al. ²⁵	2012	1	72	R	Dyspnea	700	9,5x12	607.4	Absent	TAH,BSO& omentectomy (frozen section performed)
Anastasakis et al. ²⁶	2013	1	49	R	Abdominal distension	moderate	18x12x10	404.7	Graves disease	TAH & BSO
Diavatis and Papanikolaou ²⁷	2016	1	75	L	Abdominal pain	NA	6,34x5,91	264	Absent	TAH,BSO& omentectomy (frozen section performed)
Yadav et al. ²⁸	2017	1	55	L	Abdominal distension	680	6x5x3,5	258	Absent	LSO (frozen section performed)
Fujiwara et al. ²⁹	2018	1	50	L	Abdominal distension	3300	10x8x7	1237	Absent	TAH & BSO (frozen section performed)

Present case	2020	1	42	L	Abdominal distension	6000	4x6	1700	Absent	LSO (frozen section performed)
--------------	------	---	----	---	----------------------	------	-----	------	--------	--------------------------------

The data from these 28 cases including our case was analyzed using Microsoft Office Excel.

The median age (years) was 58.85 years with a minimum of 31 years and a maximum of 83 years. Also, one patient presented with bilateral struma ovarii and one case didn't mention the side. For the rest, 12 (46.15%) cases were localized on the right ovary and 14(53.84%) on the left ovary.

82,1% of the patients presented with abdominal distension, 25 % presented dyspnea, 10,7% with abdominal pain. Other symptoms were fatigue, cachexia and nonspecific. Coexisting thyroid disease was present in 6 patients (21,4%).

Only one case (Zanonni et al.¹³) of malignant struma ovarii was associated with ascites and high levels of CA125, all others were benign struma ovarii.

Even though frozen section histopathological examination was performed in 71,4% of the cases, extensive surgery was performed in 78,57% of cases.

Discussion

Struma ovarii is a rare ovarian tumor (1% of all ovarian tumors) that is often diagnosed on the histopathological report and sometimes further immunohistology is needed in order to deliver an accurate diagnosis, given the fact that on radiological examination the characteristic features are not straightforward interpreted. Frequently, it is misdiagnosed as ovarian neoplasm and unnecessarily extensive surgery is performed.

Meigs syndrome is characterized by the triad of benign solid ovarian tumor (most frequent fibromas, cystadenoma, granulosa cell tumor or thecoma), ascites and pleural effusion that resolves after the resection of the tumor. Pseudo-Meigs syndrome consists of the same triad with the difference that the benign tumor is other than those in Meigs syndrome.²⁴

Serum Cancer Antigen 125 is a tumoral marker originating from fetal coelomic epithelium: Mullerian epithelium, pleura, pericardium and peritoneum. Even though high levels of CA125 are associated with ovarian carcinoma, it can be raised in various other gynecological pathologies: endometriosis, acute pelvic inflammatory disease, fibromatosis,³⁰ and non-gynecological: acute peritonitis, acute/ chronic pancreatitis, hepatitis, cirrhosis, renal failure, autoimmune disease.³¹

Every single case in which the suspicion is towards malignancy should be addressed to the local/hospital tumor-board for a patient-centered treatment planning. In order to avoid meaningless

surgical procedures, the surgeon should have prior histopathological confirmation of the malignancy or intraoperative biopsy with frozen section histopathological result in order to further advance with the intervention. In this case, by doing the frozen section histopathological examination, the patient received the standard of care for this type of pathology. Even though this type of tumor and clinical presentation is rare, suspicion of malignancy should always be present. The surgical standard of care for benign struma ovarii is salpingo-oophorectomy of the affected side.

Furthermore, after the histopathological department establishes the diagnoses, the patient should be referred to an endocrinologist to assess the thyroid function. If malignant struma ovarii is discussed, the patient should be referred to a surgeon specialized in endocrine surgery in order to undergo a total thyroidectomy followed by radioiodine ablation therapy and subsequent oncological evaluation.

Conclusion

Struma ovarii is a rare tumor, with only 27 cases of struma ovarii associated with high levels of CA125 and ascites or Pseudo-Meigs' syndrome reported in the literature. In this case not only the patient had struma ovarii and high levels of CA 125 but was accompanied by grade III ascites, similar with ovarian carcinoma, making the therapeutical decision more difficult.

Abbreviation List

BSO -bilateral salpingo-oophorectomy

CA125 – cancer antigen 125

CK7- cytokeratin 7

HE- hematoxylin and eosin stain

HE4 – human epididymis protein 4

Ki67-cellular marker for proliferation

L -left

LSO – left salpingo-oophorectomy

NA- non assessed

R -right

RSO – right salpingo-oophorectomy

TAH – total abdominal hysterectomy

TTF1- thyroid transcription factor-1

Disclosure: The authors have nothing to disclose.

Funding: The authors declare no specific funding for this research from any funding agency in the public, commercial, or not-for-profit sectors.

Competing interests: The authors declare none.

Patient consent: The patient consent was obtained.

References

1. Roth LM, Talerma A. The enigma of struma ovarii. *Pathology*. 2007 Feb;39(1):139-46. PubMed PMID: 17365830.
2. Yoo SC, Chang KH, Lyu MO, Chang SJ, Ryu HS, Kim HS. Clinical characteristics of struma ovarii. *J Gynecol Oncol*. 2008 Jun;19(2):135-8. PubMed PMID: 19471561. Pubmed Central PMCID: PMC2676458.
3. Willemse PH, Oosterhuis JW, Aalders JG, Piers DA, Sleijfer DT, Vermey A, et al. Malignant struma ovarii treated by ovariectomy, thyroidectomy, and 131I administration. *Cancer*. 1987 Jul 15;60(2):178-82. PubMed PMID: 3297279.
4. Rubinsak L, Chelmow, D. Struma Ovarii. *Medscape* [updated Mar 8, 2019; cited 2019]; Available from: <https://emedicine.medscape.com/article/256937-overview#a6>.
5. Jotkowitz MW, Gee DC. Unique case of massive ascites, extreme elevation of serum CA 125 tumour marker. *Aust N Z J Obstet Gynaecol*. 1993 Nov;33(4):453-4. PubMed PMID: 8179576.
6. Leung YC, Hammond IG. Limitations of CA125 in the preoperative evaluation of a pelvic mass: struma ovarii and ascites. *Aust N Z J Obstet Gynaecol*. 1993 May;33(2):216-7. PubMed PMID: 8216132.
7. Bethune M, Quinn M, Rome R. Struma ovarii presenting as acute pseudo-Meigs syndrome with an elevated CA 125 level. *Aust N Z J Obstet Gynaecol*. 1996 Aug;36(3):372-3. PubMed PMID: 8883774.
8. Mancuso A, Triolo O, Leonardi I, De Vivo A. Struma ovarii: a rare benign pathology which may erroneously suggest malignancy. *Acta Obstet Gynecol Scand*. 2001 Nov;80(11):1075-6. PubMed PMID: 11703211.
9. Long CY, Chen YH, Chen SC, Lee JN, Su JH, Hsu SC. Pseudo-Meigs syndrome and elevated levels of tumor markers associated with benign ovarian tumors--two case reports. *Kaohsiung J Med Sci*. 2001 Nov;17(11):582-5. PubMed PMID: 11852467.
10. Huh JJ, Montz FJ, Bristow RE. Struma ovarii associated with pseudo-Meigs' syndrome and elevated serum CA 125. *Gynecol Oncol*. 2002 Aug;86(2):231-4. PubMed PMID: 12144834.
11. Loizzi V, Cappuccini F, Berman ML. An unusual presentation of struma ovarii mimicking a malignant process. *Obstet Gynecol*. 2002 Nov;100(5 Pt 2):1111-2. PubMed PMID: 12423825.
12. Bokhari A, Rosenfeld GS, Cracchiolo B, Heller DS. Cystic struma ovarii presenting with ascites and an elevated CA-125 level. A case report. *J Reprod Med*. 2003 Jan;48(1):52-6. PubMed PMID: 12611097.
13. Zannoni GF, Gallotta V, Legge F, Tarquini E, Scambia G, Ferrandina G. Pseudo-Meigs' syndrome associated with malignant struma ovarii: a case report. *Gynecol Oncol*. 2004 Jul;94(1):226-8. PubMed PMID: 15262148.
14. Rim SY, Kim SM, Choi HS. Struma ovarii showing clinical characteristics of ovarian malignancy. *Int J Gynecol Cancer*. 2005 Nov-Dec;15(6):1156-9. PubMed PMID: 16343201.
15. Guida M, Mandato VD, Di Spiezio Sardo A, Di Carlo C, Giordano E, Nappi C. Coexistence of Graves' disease and benign struma ovarii in a patient with marked ascites and elevated CA-125 levels. *J Endocrinol Invest*. 2005 Oct;28(9):827-30. PubMed PMID: 16370564.
16. Loizzi V, Cormio G, Resta L, Fattizzi N, Vicino M, Selvaggi L. Pseudo-Meigs syndrome and elevated CA125 associated with struma ovarii. *Gynecol Oncol*. 2005 Apr;97(1):282-4. PubMed PMID: 15790478.
17. Uehara T, Sawada M. Struma ovarii associated with Meigs syndrome. *Jpn J Clin Oncol*. 2007 Jan;37(1):73. PubMed PMID: 17272322.
18. Obeidat BR, Amarín ZO. Struma ovarii with pseudo-Meigs' syndrome and elevated CA125 levels. *J Obstet Gynaecol*. 2007 Jan;27(1):97-8. PubMed PMID: 17365481.
19. Mitrou S, Manek S, Kehoe S. Cystic struma ovarii presenting as pseudo-Meigs' syndrome with elevated CA125 levels. A case report and review of the literature. *Int J Gynecol Cancer*. 2008 Mar-Apr;18(2):372-5. PubMed PMID: 18334015.
20. Paladini D, Vassallo M, Sglavo G, Nappi C. Struma ovarii associated with hyperthyroidism, elevated CA 125 and pseudo-Meigs syndrome may mimic advanced ovarian cancer. *Ultrasound Obstet Gynecol*. 2008 Aug;32(2):237-8. PubMed PMID: 18618418.
21. Mui MP, Tam KF, Tam FK, Ngan HY. Coexistence of struma ovarii with marked ascites and elevated CA-125 levels: case report and literature review. *Arch Gynecol Obstet*. 2009 May;279(5):753-7. PubMed PMID: 18807056.
22. Rana V, Srinivas V, Bandyopadhyay S, Ghosh SK, Singh Y. Bilateral benign non functional struma ovarii with Pseudo-Meigs' syndrome. *Indian J Pathol Microbiol*. 2009 Jan-Mar;52(1):94-6. PubMed PMID: 19136795.
23. Jiang W, Lu X, Zhu ZL, Liu XS, Xu CJ. Struma ovarii associated with pseudo-Meigs' syndrome and elevated serum CA 125: a case report and review of the literature. *J Ovarian Res*. 2010 Jul 29;3:18. PubMed PMID: 20670426. Pubmed Central PMCID: PMC2923141.

24. obeidat BRp, T. J. ; Saidi, S.A. Struma ovarii associated with Pseudo-Meigs' syndrome and elevated serum CA 125: A case report and literature review. *Gynecology Obstetric*. 2012;2(129).
25. Mostaghel N, Enzevaei A, Zare K, Fallahian M. Struma ovarii associated with Pseudo-Meig's syndrome and high serum level of CA 125; a case report. *J Ovarian Res*. 2012 Mar 21;5:10. PubMed PMID: 22436494. Pubmed Central PMCID: PMC3350392.
26. Anastasilakis AD, Ruggeri RM, Polyzos SA, Makras P, Molyva D, Campenni A, et al. Coexistence of Graves' disease, papillary thyroid carcinoma and unilateral benign struma ovarii: case report and review of the literature. *Metabolism*. 2013 Oct;62(10):1350-6. PubMed PMID: 23806737.
27. Diavatis S, Papanikolaou A. Level of HE4 is Correlated with Diagnosis of Struma Ovarii: A Case Report. *Am J Case Rep*. 2016 Jul 6;17:459-61. PubMed PMID: 27381498. Pubmed Central PMCID: PMC4939854.
28. Yadav S, Tomar R, Verma N, Khurana N, Triathi R. Struma Ovarii with Pseudo-Meigs' Syndrome and Raised Cancer Antigen-125 Levels Masquerading as an Ovarian Carcinoma Case report and literature review. *Sultan Qaboos Univ Med J*. 2017 May;17(2):e229-e33. PubMed PMID: 28690899. Pubmed Central PMCID: PMC5488828.
29. Fujiwara S, Tsuyoshi H, Nishimura T, Takahashi N, Yoshida Y. Precise preoperative diagnosis of struma ovarii with pseudo-Meigs' syndrome mimicking ovarian cancer with the combination of (131)I scintigraphy and (18)F-FDG PET: case report and review of the literature. *J Ovarian Res*. 2018 Feb 2;11(1):11. PubMed PMID: 29391043. Pubmed Central PMCID: PMC5796437.
30. He RH, Yao WM, Wu LY, Mao YY. Highly elevated serum CA-125 levels in patients with non-malignant gynecological diseases. *Arch Gynecol Obstet*. 2011 Mar;283 Suppl 1:107-10. PubMed PMID: 21069368.
31. King GGL, John; . CA125. 2020; Available from: <https://emedicine.medscape.com/article/2087557-overview#a2>.