SOME SPECIES OF *CROTALARIA* IN AUSTRALIA

ALMA T. LEE

(Accepted 7.1977)

ABSTRACT

Lee, A. T. (National Herbarium of New South Wales, Royal Botanic Gardens, Sydney, Australia 2000) 1978. *Some species of Crotalaria in Australia.* *Telopea* 1 (5): 319–356.—The taxa currently or in the past known by the names *C. dissitiflora* Benth. and its vars. *rugosa* Benth., *eremaea* (F. Muell.) Benth., and *grandiflora* Benth., *C. eremaea* F. Muell., *C. benthamiana* E. Pritzel, *C. strehlowii* E. Pritzel (Neotype designated), *C. novae-hollandiae* DC. and its formae *lasiophylla* Benth. (Lectotype designated), *parviflora* Benth., and *oblongifolia* Benth., *C. oblongifolia* Hook., *C. crassipes* Benth., and *C. mitchellii* Benth. and its var. *tomentosa* Ewart are critically examined. Changes in circumscription and nomenclature are proposed and the following are recognized: *C. benthamiana* (Lectotype designated); *C. dissitiflora* with two subspecies; *C. eremaea* with two subspecies incorporating *C. strehlowii*; *C. mitchellii* with two subspecies, including subsp. *laevis* subsp. nov., and *C. smithiana* nom. et stat. nov. replacing *C. mitchellii* var. *tomentosa*; *C. novae-hollandiae* with two subspecies incorporating the three formae of Bentham together with *C. crassipes*, a Lectotype designated for f. *lasiophylla*; and *C. medicaginea* again recognized as the correct name for Australian plants now often identified as *C. trifolium*. Species naturalized in New South Wales are enumerated with some changes in nomenclature resulting from recent work on African species.

In preparing an account of the genus *Crotalaria* for the Flora of New South Wales, several species presented problems requiring consideration of extensive populations partly or even wholly outside this State. Thus *C. dissitiflora* and its var. *eremaea* have long been recorded for New South Wales, but the populations occurring in this State are now seen to be the southeastern extremities of two very extensive, polytypic species *C. dissitiflora* and *C. eremaea*; the large collections of both in several Australian herbaria are reviewed, more fully described, and their localities mapped. The Western Australian taxon variously known as *C. benthamiana* and *C. dissitiflora* var. *grandiflora* is re-examined and the relationships of the taxa are briefly discussed.

Similarly, the large collections of both *C. novae-hollandiae* and *C. mitchellii* which have been critically examined show these to be two allopatric species each with an extensive range, in northern and eastern Australia respectively; both are polytypic and of the two only *C. mitchellii* occurs in New South Wales. *C. novae-hollandiae* is considered to comprise two subspecies, incorporating but not closely corresponding with the forms distinguished by Bentham. Two subspecies are recognized in *C. mitchellii* also, and subsp. *laevis* subsp. nov. is described. *C. mitchellii* var. *tomentosa* is raised to specific rank as *C. smithiana*; it resembles *C. mitchellii* chiefly in having small flowers, but is quite distinct in its unifoliate (not simple) leaves, and also quite disjunct geographically.

Distribution maps are provided for all these taxa, which are discussed under the following headings:

1. The *C. dissitiflora* complex
 - i. *C. benthamiana*
 - ii. *C. dissitiflora*
 - a. subsp. *dissitiflora*
 - b. subsp. *rugosa*
 - iii. *C. eremaea*
 - a. subsp. *eremaea*
 - b. subsp. *strehlowii*
2. *C. mitchellii* and *C. smithiana*
 - i. *C. mitchellii*
 - a. subsp. *mitchellii*
 - b. subsp. *laevis*
 - ii. *C. smithiana*
3. *C. novae-hollandiae* (including *C. crassipes*)
 - C. novae-hollandiae*
 - a. subsp. *novae-hollandiae* (syn. *C. crassipes*)
 - b. subsp. *lasiophylla*
4. *C. medicaginea* or *C. trifoliatrum*
 - C. medicaginea*
5. Species naturalized or escaped from cultivation in New South Wales—a check list.

In the text below, all specimens cited as Types have been examined unless otherwise indicated; they are not cited again among specimens examined but localities are mapped where appropriate.

Terms used to describe shape of parts are those defined in Chart 1a of the Systematics Association Committee (1962), and their precise definition is indicated by the formula following in parentheses e.g. (SAC 7-9), also taken from that chart. Flower parts are measured as shown in Fig. 3.

Distribution maps have been compiled from specimens cited, including Types whether seen or not, but where crowding of symbols was confusing, (near) duplicate localities have not been mapped and so the number of citations does not always match the number of symbols for a given taxon. Although an accurate representation of localities has been sought, the scale of symbol and map and lack of precision in data, especially on early labels (some Types), make caution necessary in interpreting some of the localities shown.

The subdivisions or Districts of the Australian States used in the citation of specimens are taken from the following sources:—

Queensland: map of pastoral districts supplied by BRI.

New South Wales: map in Contr. New South Wales Natl. Herb. Fl. Ser. 1-18: 16 (1961).

South Australia: author's preface in J. M. Black, Fl. South Australia ed. 2, 1: 5 (1948), or labels on the sheets in AD.

Western Australia: map in Gardner, C. A. & H. W. Bennetts, *The Toxic Plants of Western Australia*: 206 (1956). West Australian Newspapers, Perth.

Northern Territory: map in Chippendale, *Check List of Northern Territory Plants*, in *Proc. Linn. Soc. New South Wales* 96: 209 (1971).

1. THE *C. DISSITIFLORA* COMPLEX

Until recently, large and very extensive populations of *Crotalaria* in Australia were covered by the name *C. dissitiflora* and, though other names were published and sometimes used for parts of these regional populations, the application of the names and relations within the group were imperfectly understood. It is now evident from the larger collections available that there are three parts of the total population which maintain their distinctness to a high degree. Most of the difficulties which arise in the assigning of a few herbarium specimens I believe to be due to deficiencies in the specimens themselves and lack or confusion of pertinent data, rather than to any real breakdown of isolating mechanisms.

The first part of the complex is (i) *C. benthamiana*, a monotypic species limited to the Fortescue District of Western Australia where it appears to maintain its distinctness by complete geographic isolation. It is close to one subspecies of *C. dissitiflora* and like the whole of that species shows a preference for heavy soils. The other major parts comprise the taxa usually known as (ii) *C. dissitiflora*, and (iii) *C. eremaea* and *C. strehlowii*, and the morphological differences between these two parts are supported by a largely allopatric distribution and apparently quite rigid preferences for different soils. These soil preferences have long been recognized in Queensland, and were incorporated by the late L. S. Smith in a working key to species of *Crotalaria* in Australia which he compiled for herbarium use. *C. dissitiflora* occurs, as is indicated by many collectors, only on heavy, often dark soils, while *C. eremaea* (here including *C. strehlowii*) is nearly always found on sand.

(ii) *C. dissitiflora*, as recognized here, comprises two geographic races. One is largely tropical, from the Ord District of Western Australia to Queensland, and the other subtropical in Queensland and New South Wales. Names are already published for these races, but not in the subspecific category which I consider is here so clearly indicated, and new combinations are therefore provided below. Subsp. *dissitiflora* is well documented as to the heavy soils on which it occurs; we have less information about the other subspecies, but those specimens which are annotated or from known soils indicate a similar preference for heavy soils, in the Northern Territory and Western Australia, as well as in Queensland.

Within *C. dissitiflora*, subsp. *dissitiflora* is distinguished from subsp. *rugosa* chiefly by the length and density of pubescence with a resultant appearance of the venation, and intergradation between the subspecies is shown in these characters (see *C. dissitiflora*, Distribution).

(iii) The third major part of the total population is very widespread in eremean parts of Western Australia, the Northern Territory and South Australia. As in *C. dissitiflora*, there are two components, here clearly distinguished over most of their extensive range and, predictably, already named in specific rank—as *C. eremaea*

(1859) and *C. strehlowii* (1918). There are, however, a number of herbarium specimens which do not fall comfortably into either of these components and which have mostly been found in the zone of contact between them. Nevertheless, great dissimilarity between the largely stabilised extremes of the variation permits a division on morphological characters which, while partly arbitrary, does correspond broadly with geographic distribution and features of habitat, and these last factors could be expected to operate towards further isolation of the two components. The arguments for recognition of the two in specific as against subspecific rank seem, in fact, to be almost equally tenable, but I propose to adopt the treatment which more clearly reflects the biological situation, i.e. to recognize a single species, *C. eremaea*, with two imperfectly discrete geographic subspecies.

Between these two subspecies the most reliable diagnostic character is leaflet form, which is largely correlated with density and extent of pubescence. There are also differences in flower size and form in the species, as there are in *C. dissitiflora*, but in *C. eremaea* the number of variants is greater and there is poorer correlation with other characters. The Central Australian plants (subsp. *strehlowii*) probably have the largest "straight" flowers, while the Simpson Desert populations (mostly subsp. *eremaea*) more often display the small "round" form of flower (Figs. 1, 2), though there are many exceptions to this generalization and the difference is influenced by age of flower and early pod development. While both subspecies of *C. eremaea* occur on sandy soils, the Type subspecies is most common in large communities of the Simpson Desert and its extensions; specimens of the other subspecies are more often annotated as from creek or river bed or bank, as well as from sandhills of Central Australia. Little is known yet of Western Australian occurrences except that most of the few collections are from the southern half of the eremean Carnegie District, where sandy soils of the Great Victoria and Gibson Deserts are predominant. Intergradation between the subspecies occurs commonly along their zone of contact, northwards from Spencer's Gulf.

It seems probable that *C. benthamiana*, *C. dissitiflora* and *C. eremaea* have evolved from a common stock, diverging under the influence of changing climatic and geographic conditions, and that the derivatives have become stabilized genetically in relation to the nature of the soil and to geographic isolation. In the Northern Territory, where *C. dissitiflora* subsp. *rugosa* is occasionally sympatric with *C. eremaea* subsp. *strehlowii*, there appears to be complete discontinuity. Further south in Western Australia, there is no known contact of either *C. benthamiana* or *C. dissitiflora* with *C. eremaea*, although paucity of collections from some 700 km of intervening desert may be obscuring a greater propinquity than is at present apparent. Thus, in spite of some confusion, I believe there is at least a very nearly complete discontinuity between these species, even between *C. dissitiflora* and *C. eremaea*, and indeed the great majority of the large collections of this whole complex can be confidently assigned to their species. But there have been difficulties in specific separation, and for some time a very few specimens, with unquestionable characters of *C. dissitiflora* from areas in which *C. eremaea* clearly predominates, threatened to make nonsense of the taxonomic arrangement which so many other collections had supported. Some of these difficulties arose in areas where different soils are encountered in relatively small patches or tongues, and *C. dissitiflora* was in fact on a tongue of heavy soil in a wider area of sand characteristically inhabited by *C. eremaea*. It is possible that three isolated collections from Central Australia,

in the very heart of the *C. eremaea* subsp. *strehlowii* population, may prove to represent small disjunct populations of *C. dissitiflora* persisting in patches, including gilgais, of heavy soils recently mapped for nearby areas (Stephens 1961, Jackson 1962, Litchfield 1969). They could possibly, though not probably, be relics of a few plants introduced for garden cultivation. Whatever the explanation, the following three specimens are *C. dissitiflora* subsp. *dissitiflora* on morphological grounds (the third with some reservations) and, though separated by vast distances across sandy areas from the main population, are at least in the right latitudes for this subspecies: *Chippendale NT 4080*, *Johnson NT 55*, *Latz 158*: see citations under *C. dissitiflora* subsp. *dissitiflora* from Northern Territory, and Map 1 (see p. 330).

The other serious problem arose in assigning three specimens from the "corner country" of Queensland and South Australia. Two of these were collected from brown loam soils (the third lacks information on soil) which are atypical for either species. The dominant *Crotalaria* in this area is *C. eremaea* subsp. *eremaea* but the morphology of these specimens is closest to that of *C. dissitiflora* subsp. *rugosa* which is not known so far south. The conviction persists however, that they are part of the *C. eremaea* population, though more knowledge concerning local soils (cf. Stephens, 1963) and their possible effects on morphology in this group may require revision of this opinion. The possibility of incomplete discontinuity in this zone must also remain for consideration. The three specimens are cited under *C. eremaea* subsp. *eremaea* marked *, and shown on Map 1: *Johnson 73/134*, *Johnson 73/139*, *Basedow 51*.

TABLE 1
DIFFERENCES BETWEEN *C. BENTHAMIANA*, *C. DISSITIFLORA* AND *C. EREMAEA*

	<i>C. dissitiflora</i> and <i>C. benthamiana</i>	<i>C. eremaea</i>
Plants	Usually branched (and regenerating) from deeply sited, $\pm$ congested, ascending underground parts; to c. 0.5 m in height.	Often branched from a main stem, apparently individual plants, but at least sometimes connected by (and regenerating from) extensive $\pm$ horizontal underground parts; usually 0.5–1 m in height.
Leaves	Rather thin, not wrinkling on drying; always with some pubescence below, glabrous to densely pubescent above, the upper surface at least and where visible through the pubescence, minutely dotted.	Rather thick, in texture or at least by virtue of dense pubescence, wrinkling on drying if surface visible; glabrous to densely tomentose with many intergrades; leaf surface where visible minutely pitted or wrinkled rather than dotted.
Leaflets	Generally narrow-elliptical to -ovate, the laterals like the terminal and not less than half its length.	Generally narrow-oblong to trullate, the laterals if present usually much reduced, sometimes almost circular, or at most half as long as the terminal.
Stipules	Present on at least some leaves, usually 1–5 mm long (rarely less than 1 mm), curved-spreading.	Usually absent, or less than 1.5 mm long, rarely to 4 mm and then not usually spreading.
Flowers	In <i>C. dissitiflora</i> : Almost always small and "round" (Fig. 1), 8–11 mm long, rarely c. 14 mm. In <i>C. benthamiana</i> : Long-beaked and "straight" (Fig. 2), 15–17 mm long.	"round" or "straight", 12–15 (–18) mm long.

KEY TO THE SPECIES OF THE *C. DISSITIFLORA* COMPLEX

(See also Table 1)

1. Leaves always trifoliolate, the lateral leaflets at least half as long as the terminal. Plants 20–30 (–40) cm high, on heavy soils.
2. Flowers large, 15–17 mm long, the keel with a straight beak extending forward and not incurved (Fig. 2). In the Fortescue District of Western Australia
..... *C. benthamiana*
- 2.* Flowers smaller, mostly 8–11 mm long, the keel rounded with a slightly incurved beak (Fig. 1). In the Ord District of Western Australia, tropical Northern Territory and Queensland, and subtropical Queensland and New South Wales. .
..... *C. dissitiflora*
- 1.* Leaves unifoliolate, or if trifoliolate the lateral leaflets usually much reduced or less than half as long as the terminal (but rarely, along S.A.-N.S.W. and -Q. borders and in W.A., about half as long as the terminal). Flowers various in form. Plants to c. 1 m high, on sandy soils. In eremean parts of southern Western Australia, Northern Territory, Queensland and New South Wales *C. eremaea*

1 i *C. benthamiana* Pritzel in Diels & Pritzel in Bot. Jahrb. Syst. 35: 267 (1904); Gardner, Enum. Pl. Austral. Occ.: 62 (1930); Beard, (Descr. Cat.) W. Austral. Pl.: 45 (1965) and ed. 2: 65 (1970). —*C. dissitiflora* var. (?) *grandiflora* Benth., Fl. Austral. 2: 184 (1864). TYPE: Typification of this taxon poses a problem of interpretation. Pritzel's protologue, while stating unequivocally that his name, *C. benthamiana*, was that of a *new species*, also explicitly embraced the taxon previously described by Benth. His publication could therefore legitimately be considered to comprise a *new name* and *new status* rather than a new species, and in this case the Type would be that of Benth. variety. Either argument could be upheld, but since Pritzel's cited specimen has not been available to me and is presumed destroyed in B, it is convenient to consider the circumscription of *C. benthamiana* to embrace all three of the specimens cited by the two authors as Syntypes, and from them to select a Lectotype. In this case there is no reason to doubt that all are conspecific. SYNTYPES: (i) Hammersley [Hamersley] Range [Western Australia-Fortescue], *M.B.* [i.e. *Maitland Brown*, collector with F. Gregory's Expedition to the North-West] (MEL 1010368, "seen by Benth."); also a print at NSW of CSIRO Neg. No. Kew 1121); (ii) not seen: Port Nichol and De Gray R [Western Australia-Fortescue], *Ridley's Expedition*; (iii) not seen, believed destroyed in B: in the tropical region from Nickol Bay to Spring Station [Western Australia-Fortescue], in dry gravelly hills, fruiting April, *E. Pritzel* (?B). LECTOTYPE, here designated: (i) above, cited as "Hammersley Range, *F. Gregory's Expedition*".

[*C. dissitiflora* sensu auctt., non Benth. sensu str.: Mueller (1862) in part, viz. the Hamersley Range Type (see also *C. dissitiflora* subsp. *rugosa*), in reference below; Beard, Wildfl. Northwest: 9, 28 (c. 1972). These authors here used *C. dissitiflora* for *C. benthamiana* which they regarded as a part of *C. dissitiflora*].

Soft-wooded perennials, apparently (from the few specimens available) identical in habit with *C. dissitiflora*, the leaves always trifoliolate (except the earliest), petiolate and stipulate, the stipules usually persistent and spreading. *Leaflets* of thin texture but quite densely clothed with loosely appressed shining hairs partly obscuring the minute dotting of both surfaces; lateral leaflets at least half as long

as the terminal, all elliptical (SAC 2-4) or obovate (SAC 46-48) to oblong (SAC 15-16), a few showing the weakly bullate effect of the pubescence and venation seen in *C. dissitiflora* subsp. *rugosa*. *Inflorescences* terminal on stems and branches, as in *C. dissitiflora*. *Flowers* clear yellow without markings, larger than in the related taxa, long and "straight" (Fig. 2) in form. *Calyx* broadly campanulate, with five subequal teeth slightly exceeding the tube and the very short receptacle, pubescent, with a pair of very small bracteoles inserted on the receptacle. *Standard* broad-ovate to almost circular. *Wings* without a distinct auricle. *Keel* somewhat angular with the beak extended forward and not incurved, the claw very short. *Ovary* pubescent; fruits and seeds not seen.

Terminal leaflets c. 2-4 cm long. *Stipules* 2-5 mm long. *Flowers* 15-17 mm long, pedicels 2-5 mm, much exceeding the bracts, c. 2 mm. *Calyx* c. 5 mm long, bracteoles 0.5-2 mm long. *Standard* 11-15 mm long, 13-16 mm broad, the claw c. 2 mm. *Wings* 12-13.5 mm long, the claw c. 2 mm. *Keel* 12-16 mm long, 5-5.5 mm broad, on a claw 1-1.5 mm long.

DISTRIBUTION (see Map 1): Known only from the Fortescue District of Western Australia, from near the coast to Wittenoom and Mt Tom Price inland; early collections are annotated "Tableland, Rocky" and "dry gravelly hills" while later records (J. S. Beard, pers. comm.) indicate "red [basalt] soil, a cracking clay giving rise to a very uneven 'crab-hole plain'".

SPECIMENS EXAMINED: WESTERN AUSTRALIA: **Northern Province-Fortescue:** Port Walcott, Harper (MEL); a sheet at PERTH ex MEL with this label is mixed, but the fragment marked β appears to be a duplicate of the collection at MEL, the others probably samples of other MEL collections of *C. dissitiflora*; Nickol R, Forrest 1878 (MEL); between Onslow and Roebourne at "Karratha" Station, Dampier, Godlonton 8-9.1968 (PERTH); "Karratha" Station, between Onslow and Roebourne, Fitzgerald 10.1968 (PERTH); Fortescue R, Carey 1878 (MEL); Wittenoom, on Port Hedland road, Garstone 7.1964 (PERTH); between Port Hedland and Mt Tom Price, Donovan 3.1970 (BRI).

The status of this taxon is debatable; vegetatively it is indistinguishable from *C. dissitiflora* subsp. *rugosa*, though in their flowers these two taxa are virtually constant and different from one another (except for rare collections from Queensland, marked * in citations of subsp. *rugosa*); they are, moreover, geographically disjunct. Since soil preference serves so notably to distinguish *C. dissitiflora* from *C. eremaea*, it is significant that *C. benthamiana* grows on heavy soils, though its flower form is not uncommon among the varying forms seen in *C. eremaea*. Nevertheless there are differences between the grey and black soils preferred by *C. dissitiflora* and the red basaltic soils of the Pilbara (a region corresponding in large part with the Fortescue District) described by Beard, and others recorded by Beadle (pers. comm. as "red-brown sandy clay with lime" from near Port Hedland), and Stephens (1963: heavy brown soils from the Fortescue R just north of the Hamersley Range). These differences, according to Beard, are associated with lower rainfall in the Pilbara. Thus *C. benthamiana* shows some affinities with *C. dissitiflora*, as well as some differences, and this is a situation recognized in other taxa separated into Pilbara and Kimberley occurrences. Here there appears to be almost complete morphological discontinuity in floral characters † supported by an allopatric distribution, while the subspecies of *C. dissitiflora* show a slighter degree of difference with probably considerable intergradation.

Fig 1

Fig 1. An extreme example of the "small, round flower", and keel (from a different flower).

Fig 2

Fig. 2. An extreme example of the "long, straight flower", and its keel.

Fig. 3. Diagrams showing method of measuring the petals: cl claw; standard and wing: l maximum length excluding claw; br maximum breadth; keel: l length & br breadth as shown; bk beak.

1 ii *C. dissitiflora* Benth. in Mitch., Trop. Australia: 386 (1848); Bentham, Fl. Austral. 2: 184 (1864) excl. var. *eremaea*; F. Mueller, Fragm. 3: 56 (1862), and Fragm. 9: 156 (1875) excl. var. *eremaea*; Moore & Betche, Handb. Fl. New South Wales: 145 (1893) excl. var. *eremaea*; F. M. Bailey, Queensland Fl. 2: 376 (1900) excl. var. *eremaea*; Maiden & Betche, Census New South Wales Pl.: 106 (1916) excl. var. *eremaea*; Ewart & Davies, Fl. N. Territory: 141 (1917) excl. "glabrous form . . . Hill No. 54"; Domin in Biblioth. Bot. 22 (89²): 735 (1925) excl. reference to South Australian occurrence; Polhill in Kew Bull. 22: 247 (1968) excl. synonymy. **HOLOTYPE:** Baloon [Balonne] R [Queensland-Maranoa], Mitchell, 17 Nov. 1846 (No. 459, K, print at NSW of CSIRO Neg. No. 1118). The Type sheet bears three specimens but, from details given in the text of Mitchell's journal, only that cited can be the Holotype. A second, No. 379, bears a collection date (somewhat ambiguous but possibly applying to both the numbers on its label) previous to that mentioned in Bentham's description, while the third is undated (Herb. Hookerianum) but possibly an Isotype. Two specimens, matched with No. 459 above by Mr L. Pedley while Australian Botanical Liaison Officer at Kew, have been examined on loan (marked * in citations).

See under the subspecies for synonymy.

† See *C. dissitiflora* subsp. *rugosa* in key, p. 331.

References in the literature under this name seem, so far as can be ascertained from their content, localities cited, or areas treated, rarely to apply to the taxon in the sense adopted here. Now that *C. dissitiflora* is seen as specifically distinct from *C. benthamiana* and *C. eremaea*, the name *C. dissitiflora* becomes "misapplied" when used for all or part of these taxa as follows:

[*C. dissitiflora* was misapplied by Black (1948) and Eichler (1965) to the polytypic species *C. eremaea* q.v.; by Tate (1890) to *C. eremaea* subsp. *eremaea* q.v. (or perhaps to *C. eremaea* as a whole); by Tate (1896), Ewart & Davies "glabrous form" (1917), Gardner (1930) and Beard (1970), to *C. eremaea* subsp. *strehlowii*, q.v.; and by Beard (c. 1972) to *C. benthamiana* q.v.].

Soft-wooded perennials of small stature, usually less than 30 cm high, with a system of deeply sited and rather congested, ascending underground structures (histologically roots by comparison with *C. eremaea*) from which regeneration occurs after adverse climatic periods or disturbance such as ploughing. *Stems* several from the base, erect or decumbent, at first densely pubescent with loosely appressed silvery hairs becoming sparse and brownish with age. *Leaves* always trifoliate (after the primary, and perhaps also early leaves on regenerating stems), petiolate and stipulate; stipules linear, characteristically spreading, persistent on at least some leaves, but small and sometimes obscured, or fallen from old, dried specimens. *Leaflets* of rather thin texture, elliptical to ovate or obovate (SAC 2-3, 37-39 or 46-48), the laterals at least half as long as the terminal, variously pubescent, the upper or both surfaces minutely dotted (when visible through the pubescence). *Inflorescences* terminal on stems and branches, racemes of about 10-20 flowers on axes usually a little longer than their peduncles. *Flowers* with rare exceptions, small and "round" (Fig. 1), clear golden yellow without other colouring or markings, pedicellate in the axils of small, finally recurved bracts. *Calyx* broadly campanulate, with five subequal teeth, the upper two usually a little shorter and narrower than the lower three, slightly exceeding the tube; tube slightly longer than and contracted into a short, basal receptacle; bracteoles minute, inserted usually near the top of the receptacle. *Standard* glabrous, nearly circular but slightly broader than long, scarcely auriculate, the calli extending on to the short claw, the apex shortly retuse or with a point in the indentation. *Wings* rounded-oblong to triangular, without a distinct auricle. *Keel* rounded, the beak slightly incurved. *Ovary* pubescent or hoary but sometimes apparently glabrous on the sides, with a more or less broad band of longer, rather coarse hairs along the upper suture, especially at its distal end. *Style* in a geniculate curve within the keel, bearded only on the inner side. *Pod* sparsely pubescent all over with short hairs outside, the sutural band now scarcely distinguishable, glabrous and shiny inside, oblong-clavate, tapered asymmetrically into a slender stipe. *Seeds* rarely seen mature on herbarium specimens though many apparently fully grown pods are present, only slightly longer than broad, yellow-ochre in colour with a marked but scarcely hooked radicular lobe, the surface almost smooth.

Terminal leaflets often 15-30 (-40) mm long. *Stipules* (0.5-) 1-3 (-5) mm long. *Rachis* of inflorescence 6- more than 16 cm, peduncle 2- more than 10 cm long. *Flowers* mostly 8-11 mm, rarely -14 mm long; pedicels 2-3 mm, bracts 1-2.5 mm long. *Calyx* c. 5 mm, bracteoles to c. 1 mm long. *Standard* (6.5-) 7-9 (-10.5) mm long and (7-) 8-11 (-14) mm broad, the claw c. 2 mm. *Wings* 7-9 (-10) mm long, 3-5 mm broad, the claw 1-3 mm. *Keel* 7-10 (-12) mm long, 4.5-6 mm broad, the claw 1-1.5 mm. *Pod* often 15-25 mm long, 5-8 mm diam., stipe 5-7 mm; seeds 2-3 mm.

Map 1. *Crotalaria benthamiana*, *C. dissitiflora* and *C. eremaea*.

DISTRIBUTION (see Map 1): Across tropical Australia, from the Ord District of Western Australia to the Northern Territory and Queensland (subsp. *rugosa*), and into subtropical Queensland and New South Wales (subsp. *dissitiflora*), but not reaching the coast in the east.

This species falls into two subspecies between which there is probably considerable intergradation, but allocation of specimens to their subspecies depends ultimately on two somewhat inexact characters, the length and density of the pubescence. Many of the specimens available are inevitably rather weatherbeaten, coming from areas of erratic and severe climatic conditions, and it is often difficult to judge the true nature of the plants represented. Thus application of subspecific names here has sometimes been arbitrary and, though indicated in the citations, the intergradant specimens are not distinguished on the map. Nevertheless a zone of intergradation can be detected chiefly in the Gregory South and Mitchell Districts of Queensland and is seen particularly in specimens from around Blackall.

Another aspect of this intergradation is that it has been more apparent in the western part of the specific range than in the eastern, and it is in this western area that *C. dissitiflora* abuts upon *C. eremaea*. It may be significant, in view of the possibly imperfect discreteness of these species, that the same two characters help to separate the species as distinguish the two subspecies of *C. dissitiflora*. For if the primary diagnostic character of the leaflets (see key on p. 324) is not decisive, then a specimen of indeterminate length and density of pubescence may appear either as an intergrade between the species, *C. dissitiflora* (subsp. *dissitiflora*) and *C. eremaea* (subsp. *eremaea*), or equally between the two subspecies (*dissitiflora* and *rugosa*) of *C. dissitiflora*. Other distinguishing characters, e.g. the greater stature and less congested habit of *C. eremaea*, and the soil preferred, are often not shown by herbarium specimens, and considerable field and experimental work may be necessary to solve this problem with any finality.

In this treatment, the species are separated as in Table 1, and in the key on p. 324, while the subspecies of *C. dissitiflora* are distinguished as follows:

1. Plants with rather scanty pubescence of short, appressed hairs c. 0.5 mm long. Leaflets always glabrous above and minutely dotted at least above and usually on both surfaces, the veins not dividing raised areas. Flowers small (mostly 8–11 mm long) and "round" (Fig. 1); keel with a short, often slightly incurved beak. Mostly subtropical, in the southeastern part of the range subsp. *dissitiflora* a.
- 1.* Plants with moderate to dense pubescence of rather long, loosely appressed, shining hairs c. 2 mm long on stems and leaves, though leaflets sometimes glabrous above, especially when old; young leaflets often but not always with a rather weakly bullate ("rugose") appearance from arrangement of pubescence at veins. Flowers as in subsp. *dissitiflora*, 6–10 mm long (very rarely –14 mm and approaching the "straight" form*). Tropical and occasionally subtropical, in the northwestern part of the range subsp. *rugosa* b.

1 ii a subsp. *dissitiflora*.—*C. dissitiflora* var. *dissitiflora*; Chippendale in Proc. Linn. Soc. New South Wales 96: 239 (1972).

[*C. dissitiflora* sensu auctt., non Benth. sensu str.: ? Mueller (1875) in part (see also subsp. *rugosa*); Moore & Betche (1893); Bailey (1900) excl. var. *rugosa*; Maiden & Betche (1916); Domin (1925) in part (see also subsp. *rugosa*); in references above. These authors there used the binomial for that part of the species occurring in the area they treated, viz. subsp. *dissitiflora*, though their descriptions sometimes covered other elements].

* These specimens are marked * in the citations of subsp. *rugosa* and approach the form seen in *C. benthamiana*.

Diagnostic characters as given in the key. *Keel* 8–10 (–12) mm long, 3–5 (–6) mm broad, the beak 6.5–8 (–9.5) mm long.

DISTRIBUTION: Inland parts of subtropical Queensland and northern New South Wales, on "black" and other heavy soils, and apparently in a few disjunct localities providing similar habitats in the Northern Territory.

SPECIMENS EXAMINED: QUEENSLAND: **Mitchell:** "Vinetree Downs", 16 miles [25 km] NE. of Blackall, *Swinburne* 2.1940 (BRI); "Malvern Hills", 22 miles [35 km] W. of Blackall, *Everist* 2021, 2.1940 (BRI); Blackall, *Ranking* (BRI); "Tambo" Station, E. of Tambo, *Smith & Everist* 876, 10.1940 (BRI). **Leichhardt:** Clermont, *Kearney* 12.1940 (BRI); 30 miles [48 km] N. of Springsure township, *Story & Yapp* 254, 9.1962 (BRI, MEL, NSW); "Karingal", Springsure, 12 miles [20 km] N. of Springsure-Rolleston road, *Gifford* 5.1964 (BRI); 10 miles [16 km] S. of Springsure on road to Rolleston, *Johnson & Blaxell* 953, 11.1972 (NSW); c. 15 miles [c. 24 km] S. of Moura on road to Theodore, *Henderson* 245, 3.1967 (BRI*, MEL, K); Dawson district, *Bancroft* 11.1915 (BRI); Theodore district, *James I*, 1.1956 (BRI); Mt Rose, 25 miles [40 km] NE. of Taroom, *Johnson* 1664, 5.1960 (BRI); "Richon", 38 miles [61 km] S. of Taroom, *Webster* 12.1961 (BRI). **Gregory South:** Windorah, 25° 26' S, 142° 39' E, *Cockburn BPS* 42, 7–8.1969 (BRI); 2 miles [3 km] from Betoota, *Beadle NSW* 113668, 5.1963 (NSW)†. **Warrego:** c. 1 mile [c. 1.6 km] S. of Charleville, *Rodd* 52, 8.1964 (NSW); Cunnamulla, *Copeman* 4.1963 (BRI), *White* 11801, 3.1941 (BRI); "Gilruth Plains" to Cunnamulla, *Martensz CSIRO W/L Herb.* 3938, 12.1967 (BRI); "Gilruth Plains", Cunnamulla, *McKee* 10313, 4.1963 (NSW, BRI); 6 miles [10 km] on Cunnamulla-Barrington road, *Ebersohn* E149, 4.1962 (BRI). **Maranoa:** between Amby and Eureka, *Hubbard & Winders* 6366, 1.1931 (BRI, K); Roma, *Stevens* 2.1958 (BRI); St George, *Winn* 5.1894 (BRI); Curriwillighi, *Dalton NSW* 45876 (NSW). **Darling Downs:** 3 miles [5 km] E. of Toobeah, *Pedley* 1190, 1.1963 (BRI)*.

NEW SOUTH WALES: **Western Plains:** 10 km W. of Barrington, *Mills & Cox* 211, 3.1975 (NSW); Moree-Boggabilla road, *Jackson NSW* 115136, 5.1972 (NSW); Collarenebri, *Waterhouse NSW* 45852, 9.1950 (NSW); Moree, *Mactier NSW* 113667, 2.1969 (NSW), *Cutting NSW* 113670, 2.1961 (NSW); County Narran, *Beadle* 3.1942 (SYD); Walgett, per *Glenfield Veterin. Res. Sta.* 43/350, 2.1943 (NSW); "Omco", Burren Junction, *Mitchell NSW* 11363, 2.1967 (NSW); "Boolcarrol", *Breakwell NSW* 113662, 4.1967 (NSW); 17 miles [27 km] N. of Narrabri, *Walker NSW* 116650, 3.1976 (NSW); Narrabri, *Crogan NSW* 48928, 2.1955 (NSW). **North Western Slopes:** Tamworth, *Barnier NSW* 45863, 3.1953 (NSW).

NORTHERN TERRITORY: **Central Australia:** the following three specimens do not come within the normal geographic range of *C. dissitiflora* subsp. *dissitiflora*; see comments on p. 322–3: 5 miles [8 km] W. of "Tarlton Downs" homestead, *Latz* 158, 2.1968 (NSW, MEL); 17 miles [27 km] W. of Arltunga, *Chippendale NT* 4080, 3.1958 (PERTH); near old "Hamilton Downs" homestead, *Johnson NT* 55, 10.1956 (AD).

1 ii b subsp. *rugosa* (Benth.) A. Lee, stat. nov.—*C. dissitiflora* var. *rugosa* Benth., Fl. Austral. 2: 184 (1864); F. M. Bailey, Queensland Fl. 2: 376 (1900); Chippendale in Proc. Linn. Soc. New South Wales 96: 239 (1972). **ISO-SYNTYPES:** (i) Sturts Creek [Western Australia–Ord], *F. Mueller* 2.1886 (MEL 1010367, "seen by Bentham"); also a print at NSW of CSIRO Neg. No. Kew 1120; (ii) Newcastle Water [Northern Territory–Barkly Tableland], [*F. Mueller*] (MEL 1010366, "seen by Bentham").

[*C. dissitiflora* sensu auct., non Benth. sensu str.: Ewart & Davies (1917) excl. "Glabrous form . . . Hill No. 54 . . ."; Domin (1925) in part (see also subsp. *dissitiflora*); Mueller (1862) in part (see also *C. benthamiana*) and (1875) ? in part (see also subsp. *dissitiflora*); in references above. These authors there used the binomial for that part of the species occurring in the area they treated, viz. subsp. *rugosa*].

See Holotype paragraph p 328.

? intergrade between subspecies.

Diagnostic characters as given in the key. Distinguished from subsp. *dissitiflora* chiefly by the pubescence of its leaves and stems, the hairs longer, dense at least on the underside of the leaflets and often on both sides, but less dense when old, and sometimes lacking on the upper leaf surface.

DISTRIBUTION (see Map 1): Tropical and occasionally subtropical; in Queensland, the Northern Territory and the Ord District of Western Australia, on heavy soils including some derived from limestones.

SPECIMENS EXAMINED: QUEENSLAND: **Burke:** Flinders Range, N. Queensland [sic; ?Flinders R, as mapped], *Browne NSW 45875*, 2.1914 (NSW); Flinders R, *White 8.1916* (BRI, NSW 45865); Barkly Tableland (? Q.), *Dunn 19*, 3.1934 (BRI); Camooweal, *Bailey NSW 45856* (NSW); "Sutherland", 60 miles [97 km] NW. of Maxwellton, *Nelson & Entwistle 34*, 4.1963 (BRI); "Sommerville", 50 miles [80 km] NE. of Maxwellton, *Entwistle 17*, 5.1963 (BRI); "Silver Hills", Richmond, *Mahoney 3*, 4.1957 (BRI); "Wongalee", c. 50 (sic) miles [c. 80 km] N. of Hughenden, 20° 30' S, 144° 00' E, *Hill 31*, 6.1968 (BRI); "Bundoran", near Nonda between Hughenden and Cloncurry, *Hubbard & Winders 7277*, 2.1931 (BRI)*; "Afton Downs", 20 miles [32 km] SW. of Hughenden, *Everist 3061*, 6.1947 (BRI). **Gregory North:** near McKinley, per *Mahoney 6.1953* (BRI); "Headingly", c. 30 miles [c. 48 km] N. of Urandangie, *Everist 3305*, 12.1947 (BRI); "Elderslie", Winton, *Kennedy 25*, 9.1935 (BRI); Winton-Boulia road, 52 miles [84 km] from Winton, *Gittins 674*, 4.1963 (BRI); Winton, *Lee NSW 45868*, 11.1949 (NSW); 30 miles [48 km] SE. of Winton, *Trapnell 169*, 7.1958 (BRI); "Roxborough Downs", *Bailey 12.1895* (BRI); between Glenormiston and Toko Ranges, *Boyle 1-2.1935* (BRI); Boulia-Bedourie road c. 20 miles [c. 32 km] S. of Boulia, per *Mahoney 6.1953* (BRI). **Mitchell:** Hughenden-Muttaborra road near "Kentle Downs", 21° 30' S, 144° 00' E, *Hill 6.1968* (BRI)*; "Tiree" Station, c. 68 miles [c. 110 km] S. of Torrens Creek, *Ford NSW 45853*, 9.1949 (NSW)*; Darr R, *de Burgh Birch* (BRI)†; Isisford road on cleared gidgea . . ., *Purcell 4.1962* (BRI)†; Reserve, Barcardine road, 10 miles [16 km] from Blackall, *Cull 2.1962* (BRI)†; "Mineeda" near Blackall, *Everist 1247*, 8.1935 (BRI)†; "Mineeda" Station near Blackall, *Everist & Smith 243*, 2.1937 (BRI)†.

NORTHERN TERRITORY: **Barkly Tableland:** 1 mile [1.6 km] E. of No. 7 Bore, near Elliott's Latz 437, 2.1969 (BRI); 12 miles [19 km] SE. of "Creswell Downs" homestead, *Chippendale NT 5581*, 3.1959 (NSW); 7.7 miles [12 km] S. of Monkaderry Waterhole, "Banka Banka", *Chippendale NT 7037*, 6.1960 (NSW); Bishops Bore, 17 miles [27 km] NW. of "Brunette Downs" homestead, *Chippendale NT 5011*, 10.1958 (PERTH); Bore No. 51, "Brunette Downs NT.", *Nicholls 798*, 5.1968 (BRI); 7 miles [11 km] N. of "Brunchilly", *Chippendale NT 7063*, 6.1960 (NSW); 15 miles [24 km] E. of "Brunchilly", *Chippendale NT 7092*, 6.1960 (NSW); 2.6 miles [4 km] SE. of Ranken [store], *Chippendale NT 7199*, 6.1960 (NSW); 23 miles [37 km] W. of Camooweal, *Chippendale NT 1485*, 8.1955 (BRI); 0.6 miles [1 km] E. of "Soudan" Station, *Perry 680*, 4.1948 (BRI); 3 miles [5 km] SSE. of "Soudan", *Nicholls 876*, 6.1968 (BRI). **Central Australia:** 30 miles [48 km] SW. of "Tobermorey" homestead, *Chippendale NT 393*, 9.1954 (NSW, BRI); Macdonnell Range, lat. 18° 30' [? McDouall Range], *Macd. Stuart* (MEL, NSW).

WESTERN AUSTRALIA: **Northern Province-Ord:** 40 miles [64 km] N. of "Sturt Creek" Station, *Beard 5627*, 6.1968 (PERTH).

1 iii *C. eremaea* F. Muell., Enum. Pl. coll. A. C. Gregory in Votes & Proc. New South Wales Legisl. Assemb., 2.v.1859 (Papers relative to Expedition in search of Dr Leichhardt); F. Mueller, *Fragm.* 3: 56 (1862).—*C. dissitiflora* var. *eremaea* (F. Muell.) Benth., *Fl. Austral.* 2: 184 (1864); F. Mueller, *Fragm.* 9: 157 (1875); Moore & Betche, *Handb. Fl. New South Wales*: 145 (1893); F. M. Bailey, *Queensland Fl.* 2: 376 (1900); Maiden & Betche, *Census New South Wales Pl.*: 106 (1916). HOLOTYPE: Coopers R [Queensland/South Australia], A. C. Gregory (MEL 1010369, "seen by Benthann"; the date of collection [6.1858] has been added by J. H. Willis).

* Flowers larger than usual and perhaps approaching *C. benthamiana*.

† ? intergrade between subspecies.

See also under the subspecies for synonymy.

[*C. dissitiflora* sensu auctt., non Benth.: Black, Fl. South Australia 2: 306 (1924), and ed. 2, 2:448 (1948); here *C. eremaea* and *C. strehlowii* are regarded as synonyms of *C. dissitiflora*; Eichler, Suppl. Black's Fl. S. Australia: 186 (1965)].

Soft-wooded shrubs, densely tomentose to completely glabrous, erect-growing to about 1 m high, the individuals apparently seed-derived but probably more often regenerated from underground parts (histologically, roots) which appear to be extensive and horizontal, rather than congested and ascending as in *C. dissitiflora*. *Leaves* unifoliolate, or if trifoliolate the lateral leaflets often much reduced or at most half the length of the terminal. *Leaflets* oblong, elliptical, ovate, trullate or obovate (mostly SAC 13–14, 36–37, 56–57), of thicker texture than in *C. dissitiflora* as shown by wrinkling in dried specimens, rarely showing the minute dotting of the surfaces characteristic of that species, glabrous in extreme forms of subsp. *strehlowii* to densely tomentose in those of subsp. *eremaea*, the hairs appressed to spreading, silvery or grey, but cream, fawn or occasionally golden brown in long-dried material; petioles in subsp. *eremaea* often persisting after the lamina has fallen. *Stipules* absent or extremely small (less than 1 mm long), very rarely larger (to 4 mm long) and spreading like those of *C. dissitiflora*. *Inflorescences* terminal on stems and branches, racemes usually of c. 15–30 flowers. *Flowers* variable in size and form, pedicellate and bracteate. *Calyx* open-campanulate with five subequal teeth, usually pubescent, densely or less so corresponding with the vegetative parts, the teeth triangular, the tube of about equal length and narrowed into a short receptacle with a pair of small bracteoles attached at or near the top of the pedicel. *Standard* nearly circular (SAC 6–7) but almost always slightly broader than long, the auricles scarcely developed, the claw rather broad. *Wings* more or less obtriangular (SAC 86–87), the claw upcurved. *Keel* varying from the short, “round” form through indeterminate specimens to the long, “straight” form (Figs. 1, 2), the former probably more common; beak slightly incurved in the “round” flowers to scarcely upturned in the “straight” form. *Ovary* and *pod* hoary to pubescent, in the young stages with the hairs along the upper suture longer and denser but not significantly so in the fruit; pod oblong-clavate, tapered asymmetrically into the stipe. *Seeds* chrome yellow with a faintly dimpled surface, the radicular lobe not prominently hooked.

Terminal leaflet (10–) 25–40 (–80) mm long, (2–) 5–15 (–26) mm broad, the laterals usually less than 1 cm long; stipules usually absent, or less than 1.5 mm, rarely more. *Flowers* rarely less than 10 mm long, often 12–15 mm, occasionally to 18 mm; pedicel often 3–5 mm; bract narrow, 1.5–4 mm. *Calyx teeth* (1.5–) 2–3 (–4) mm long, tube about equal, receptacle 1–1.5 mm; bracteoles (0.5–) 1.5–2 (–3) mm long. *Standard* (8–) 10–11 (–14) mm long, (9–) 10–12 (–14) mm broad, claw (2.5–) 3–4 mm long. *Wings* (8–) 9–10 (–11) mm long, claw mostly 2–3 mm. *Keel* (8–) 11–12 (rarely –18) mm long, (4–) 4.5–5.5 (–7) mm broad, claw 2–3 (–4) mm, beak 8–10 (rarely –14) mm long. *Pod* c. 2–3 cm long, 3–7 mm diam., stipe 3–4 mm. *Seeds* 4–5 mm long, c. 3 mm broad.

DISTRIBUTION (see Map 1): In sandy soils of Central Australia, extending into eremean parts of all the adjoining States, subsp. *eremaea* in the eastern and subsp. *strehlowii* in the central and western parts of the specific range.

A species of two races, the subspecies below:

1. Leaflets essentially oblong, 1 or 3, the laterals if present reduced and sometimes very small or, if approaching the terminal in form, very rarely more than half its length, not noticeably thick except sometimes by virtue of the tomentum, the leaf surface where visible through the hairs often with fine multidirectional wrinkling or pitting when dried, densely to sparsely pubescent but always with some hairs on the upper surface, the hairs appressed to spreading and usually denser below than above; stipules absent or less than 1 mm long, occasionally obvious, rarely to 4 mm, and often obscured by the pubescence; flowers variable in form, from small and "round" to larger and "straight" (Figs. 1, 2) subsp. *eremaea* a.
- 1.* Leaflets essentially ovate to trullate, commonly 1, or 3 with the laterals usually small, often circular to occasionally ovate-oblong (perhaps more fully developed and close to those of *C. dissitiflora* in some Western Australian plants); thickish in texture and drying with a wrinkled surface, quite glabrous above or rarely with a very sparse scattering of hairs on young leaves, the lower surface usually with a more or less sparse pubescence, or quite glabrous; hairs short, appressed, or occasionally (seen only in a few collections from Central Australia) sparse and stiffly spreading; stipules usually absent, rarely present and up to 1.5 mm long; flowers often large and "straight" but grading to the small "round" form particularly in the southwestern part of the range subsp. *strehlowii* b.

1 iii a subsp. *eremaea*.

Pentadynamis incana R. Br. in Bot. App. in Sturt, Narr. Exped. Centr. Australia 2: 76 (1848), non *Crotalaria incana* L., Sp. Pl.: 716 (1753). HOLOTYPE, n.v.: "on sand-hills with *Crotalaria sturtii*, D. Sturt". Sturt recorded *C. sturtii* (i.e. *C. cunninghamii*) as "on top of the ridges in pure sand, from S. lat. 28° to 26°", and his route crossed these latitudes in the vicinity of the Warburton R. The Type of *P. incana* would therefore have come from the Simpson Desert somewhere between the Warburton and Coopers Creek, in South Australia, c. 139° E long.

[*C. dissitiflora* sensu auctt., non Benth.: ? Tate, Fl. Extratrop. South Australia: 66, 223 (1890). Tate's description implies subsp. *eremaea*, but the localities cited and Tate's later (1896) usage suggest that his references applied to subsp. *strehlowii* q.v. and intergradant forms as well as to subsp. *eremaea*].

[*C. dissitiflora* var. *eremaea* sensu auctt., non Benth. sensu str.: Moore & Betche (1893); Bailey (1900); Maiden & Betche (1916); in references above in the synonymy of *C. eremaea*. These authors there used this equivalent of *C. eremaea* for that part of *C. eremaea* occurring in the area they treated, viz. subsp. *eremaea*, an interpretation supported by the descriptions given].

Characters as given in the key.

DISTRIBUTION: In the eastern and particularly southeastern parts of the specific range, i.e. in southwestern Queensland, northwestern New South Wales, northeastern South Australia and southeastern Northern Territory; nearly always in sandy soils and chiefly in the sandy expanses of the Simpson Desert and its extensions into semi-desert regions, showing various combinations of characters with typical subsp. *strehlowii* (see below) northwards from Spencers Gulf, and appearing occasionally with its own characteristic features within the area occupied by that subspecies; occasionally also on red and brown loams in northwestern New South Wales and Queensland.

SPECIMENS EXAMINED: QUEENSLAND: **Gregory North:** Glenormiston, No. 20 Bore, Colliver 8.1949 (BRI); between Glenormiston and the Toko Ranges, Boyle 1-2.1935 (BRI 126554); Boulia-Bedourie road, 20 miles [32 km] N. of Bedourie, Johnson 73/134, 9.1973 (NSW)*; Bedourie-Birdsville road, 32 miles [51 km] S. of Bedourie, Johnson 73/137, 9.1973 (NSW). **Gregory South:** Jundah to border, Cooper R, Sherman 8.1946 (BRI); "Retreat", on low sandhill, Everist 6321, 9.1960 (BRI); Bedourie-Birdsville road, 50 miles [80 km] N. of Birdsville, Johnson 73/139, 9.1973 (NSW)*; "Mooraberree", Everist 4098, 9.1949 (BRI); 36 miles [58 km] ENE. of Birdsville on Bedourie road, 25° 36' S, 139° 45' E, Briggs 4586, 5.1972 (NSW); 46 miles [74 km] E. of Birdsville on Betoota road, Rodd NSW 113675, 8.1964 (NSW); 20 miles [32 km] from

* *C. eremaea* (?—*C. dissitiflora*). See p. 323.

"Roseberth" Station towards Birdsville . . . *Johnson* 33, 6.1972 (NSW); Simpson Desert, widespread on sandhills, *Cockburn BPS* 40, 7-8.1969 (BRI); Poeppel Corner, 26° S, 138° E, *Boylard* 270, 9.1966 (BRI, K); "Ingella", c. 20 miles [c. 32 km] SE. of Windorah, *Everist* 4045, 6.1949 (BRI, K); Quilpie (but noted as coll. on red sandhill), *Smith* 11.1954 (BRI); "Thylungra", c. 75 miles [c. 121 km] NW. of Quilpie, *Everist* 5742, 10.1955 (BRI, K); 80 miles [129 km] W. [NW.] of Quilpie, *Stranger* 8, 3.1959 (BRI), *Everist* 6284, 9.1960 (BRI); "Copai", 55 miles [89 km] NW. of Quilpie, *Mackay* 8.1961 (BRI); "Nappamerrie" Station, Cooper Creek, *Jackson* 445, 8.1962 (AD); 26 mile tank-NE. "Nappamerrie", *Wilson* 9.1961 (BRI); between Innamincka S.A. and "Nappamerrie" Q., *Marks* 8.1967 (BRI); c. 10 miles [c. 16 km] NE. of "Nockatunga", *Carolin* 4150, 8.1969 (SYD); red sand dunes 80 miles [129 km] W. of Thargomindah, *Cockburn* 47, 9.1963 (BRI).

NEW SOUTH WALES: **Far Western Plains:** Queensland border, N. and a little W. of Broken Hill, *MacGillivray NSW* 45874, 4.1917 (NSW); Fort Grey, *Brough & Beadle* 8.1939 (SYD); near Fort Grey, *Brough & Beadle* 8.1939 (SYD); "Olive Downs" c. NNW. of Tibooburra, *Milthorpe* 880, 10.1972 (NSW); "Waka", c. between Fort Grey and Tibooburra, *Milthorpe* 882, 10.1972 (NSW); Bindara Tank, "Delalah Downs", *Cunningham SCS* 495, 8.1972 (NSW); 29° 22' S, 143° 03' E, "Delalah Downs", *Pickard* 2071, 3.1973 (NSW); "Gumvale" c. 25 km W. by S. of Tibooburra, *Johncock NSW* 101222, 101223, 4.1967 (NSW); Tibooburra, *Burgess & Beadle* 2.1950 (SYD); "Yandama" Station, *Collier NSW* 45857, 7.1910 (NSW); Milparinka, *Whalley NSW* 45849, 4.1957 (NSW), *Beadle* 10.1939 (SYD); "Winnathee" 60 miles [97 km] W. of Milparinka, *Johnson & Constable NSW* 33370, 6.1955 (NSW, K); Clifton Bore, E. of Milparinka, *Blaxell* 416, 4.1970 (NSW); Broken Hill + 138 miles [222 km] towards Tibooburra, *Gittins* 1910, 8.1969 (BRI, NSW); 20 miles [32 km] S. of Milparinka, *Beadle NSW* 113679, 5.1963 (NSW); Lake Yantara, *Beadle* 8.1941 (SYD); Silver City Highway, close to Salt Lake, *Martensz* 246, 12.1969 (NSW); "Callindry" Station, *Leigh W205*, 7.1968 (NSW); Cobham Lake, *Bauerlen NSW* 45866, 9.1887 (NSW), *Williams NSW* 45871, 10.1901 (NSW); "Cobham", *MacGillivray per Morris* 669, 8.1921 (NSW); sandhills, "Cobham", *Morris* 784, 10.1921 (NSW, BRI); Paldrumatta Bore, *Corbett NSW* 45869, 4.1900 (NSW); 3.2 km N. of Turleys Gate, S.A.-N.S.W. border, *Campbell & Pickard* 1256, 11.1970 (NSW); 180 km N. of Broken Hill, *Mulham W771*, 8.1974 (NSW); 35 km N. of Broughams Gate . . . 30° 35' S, 141° 00' E, *Lee* 226, 9.1971 (NSW); 0.8 km S. of "Tielta" homestead . . . 30° 50' S, 141° 14' E, *Lee* 216, 9.1971 (NSW); Sandhills, Barrier Range, *Collins* 9.1921 (SYD); Kinchega National Park, Menindee, *Ryan & Everley NSW* 113672, 6.1959 (NSW).

SOUTH AUSTRALIA: **Far North:** South of Charlotte Waters, *Kempe NSW* 45872, 9.1883 (NSW); Outer Birdsville Track, 35 miles [56 km] ex Birdsville, *Johnson* 73/140, 9.1973 (NSW); Walkers Well [prob. vicinity "Cordillo Downs"], *Basedow* 51, 10.1919 (NSW, K)*; Bradys Waterhole near Cordillo Downs, *Basedow* 61, 10.1919 (NSW); Outer Birdsville Track, 48 miles [77 km] S. of "Clifton Hills", *Johnson* 73/141, 9.1973 (NSW); Seven-mile Creek, S. of Goyders Lagoon, *Crocker* 6, 7.1939 (AD); Innamincka, *Herb. Tate* 391, 6.1884 (AD); Lake Eyre, c. 50 km NW. of Marree, *Hill* 213, 7.1955 (AD); Birdsville Track, 95 miles [153 km] N. of Marree, *Johnson* 73/144, 9. 1973 (NSW); "Killalpaninna" (Mission Station), Coopers Creek near Lake Eyre, *Howitt NSW* 45854, 12.1883 (NSW); Natarranni sandhills, Birdsville Track 85 miles [137 km] from Marree, *Johnson* 21, 6.1972 (NSW); near Cannawulicaninna Bore, Birdsville Track, c. 110 km NE. of Marree, *Crisp & Noble* 18, 4.1969 (AD); 108 km N. of Marree, *Milthorpe* 3044, 10.1974 (NSW); Birdsville Track, 7 miles [11 km] c. N. of Marree, *Johnson* 73/146, 9.1973 (NSW); Mt Lyndhurst, *Koch* 541, 10.1900 (NSW). **North Lake Torrens Basin:** Red Lake c. 130 km WSW. of Marree, *Aitken* 12.1964 (AD); "Mulgaria" homestead, 30° 14' S, 137° 38' E, *Aitken* 12.1964 (AD). **Lake Torrens Basin:** c. 26 miles [42 km] N. of Andamooka, *Macfarlane* 6.1964 (AD); Andamooka, sandhills, *Williams* 2812, 6.1966 (AD); Andamooka Opal Fields near landing strip, *Weber* 1378, 9.1968 (AD, K); c. 30 km due N. of Woocalla and c. 7 km E. of Mt Gunson, *Lorinczi & Tonkin Type D no. 1*, 9.1966 (AD).

NORTHERN TERRITORY: **Central Australia:** 7 miles [11 km] NW. of "Numagalong" Station homestead, *Nelson & Swinbourne NT* 11801, 8.1965 (NSW); Sandy Valley, Wycliffe Creek, *Allen* 598, 7.1922 (NSW); Alice Springs, *Tate* 1894 (AD); "Andado" to Georgina R, *Crocker-Simpson Desert Expedition* 1939 (AD).

* *C. eremaea* (?—*C. dissitiflora*). See p. 323.

The following specimens may be included in subsp. *eremaea* (see Map 1, subsp. *eremaea* atypical) though the diagnostic characters are less clearly shown, having less dense to sparse pubescence though still with some hairs on the upper surface of the normally oblong leaflets, or the tomentose leaflets tending to ovate shapes though often narrower than in subsp. *strehlowii* and about 5–10 mm wide.

QUEENSLAND: **Gregory North:** 13 miles [21 km] from Diamantina Shire boundary-Bedourie road on "Monkira" Station, *Williams* 153, 10.1968 (BRI). **Mitchell:** 48 miles [77 km] W. of Yaraka, *Everist* 7363, 8.1963 (BRI). **Gregory South:** sandhills about Georgina, Simpson Desert Expedition Camp 21, *Crocker* 5, 7.1939 (AD); 10 miles [16 km] from Windorah towards "Currawilla" Station, c. 25° 21' S, 142° 28' E, *Gittins* 1938, 8.1969 (BRI, NSW); 17 miles [27 km] W. of Windorah, *Everist* 1555, 6.1937 (BRI); 45 miles [73 km] S. of Windorah, *Everist* 4044, 6.1949 (BRI); 27.7 miles [c. 44 km] NW. of Tenham Bore, *Trapnell* E52, 6.1969 (BRI); sandhill at outlet of Eulbertie Waterhole, Coopers Creek, *Ogilvie* 8.1961 (BRI); 50 miles [81 km] W. of Quilpie, 5 miles [8 km] N. of the Quilpie to Eromanga road . . . "Kyra" homestead, *Emmerson* 8.1965 (BRI); "Nockatunga" Station, *Carolin* 4239, 8.1964 (SYD); SW. Queensland, *Little* 1910 (BRI).

SOUTH AUSTRALIA: **Far North:** 38 miles [c. 61 km] E. of Dalhousie Springs, *Lothian* 1432, 8.1963 (AD); also a series of other specimens from around this locality, *Lothian* 1487, 1622, 1657, 1702, 1796, 1807, all 8.1963 (AD; 1796 also at NSW). **Woomera:** "Purple Downs", *Weber* 1541, 9.1968 (AD). **Lake Torrens Basin:** Yeltacowie, *Murray* 33, 6.1927 (AD, bis); Arcoona, *Murray* 33 (sic), 8.1927 (AD).

NORTHERN TERRITORY: **Central Australia:** 17 miles [27 km] SE. of Bottom Bore, Hale R, *Chippendale* NT 4938, 9.1958 (AD, NSW, PERTH); Simpson Desert, 25° 15' S, 136° 43' E, *Symon* 4379, 11.1966 (AD).

As well as the specimens cited above, there were many in AD which I examined in 1971, determinable then as *C. eremaea*. Not having re-examined them and realizing the possibility of concept changes over the intervening period, they are not cited here, but their distribution is in agreement with that shown in Map 1.

The following specimens do not fall clearly into either of the subspecies (see Map 1, intergrades); nearly all occur in a zone between the main populations of the races, but they represent a number of combinations of the different subspecific characters, rather than a clinal intermediate series, though a near-clinal change is probably seen in South Australia. The commonest combination of subspecific characters is the leaflet shape of one subspecies with the pubescence of the other, e.g. the leaflet may be clearly oblong as in subsp. *eremaea*, but glabrous above and wrinkled as in subsp. *strehlowii*; or the leaflet shape may be rather indefinite or narrow-ovate, as in those plants of subsp. *strehlowii* with narrow leaflets, but this is combined with some, albeit sparse, pubescence on the upper surface of the leaflet. The nature of the pubescence also varies, and a few of these plants have a sparse but rather stiff and spreading pubescence which is not seen in either of the 'pure' subspecies and suggests a reduction in density of hairs of some of the densely tomentose forms of subsp. *eremaea* found in the Simpson Desert:—

NORTHERN TERRITORY: **Central Australia:** c. 15 miles [c. 24 km] S. of Wycliffe Well (c. 220 miles [c. 353 km] N. of Alice Springs), *Everist* 4279, 1.1950 (BRI); c. 55 km N. of Barrow Creek, v. *Dam* 52, 6.1969 (AD); Elkedra R, 15.2 miles [c. 24 km] E. of "Elkedra" homestead, *Chippendale* NT 3126, 10.1956 (NSW); 5 miles [8 km] SW. of "Tarlton Downs", *Winkworth* 594, 10.1954 (PERTH); creek 64 km N. of Arltunga, *Cleland* 9.1930 (AD); 4.2 miles [c. 6.5 km] E. of Allua Well, Ross R, *Chippendale* NT 4965, 9.1958 (NSW); 70 miles [113 km] SE. of "Ringwood", *Winkworth* 1273, 9.1955 (BRI, NSW, PERTH); 37 mile peg S. of Alice Springs, *Carolin* 5153, 8.1966 (SYD); Window Hill, Simpson Desert, *Chippendale* NT 1607, 9.1955 (AD, BRI, NSW).

1 iii b subsp. *strehlowii* (Pritzel) A. Lee stat. nov.—*C. strehlowii* Pritzel in Feddes Rept. 15: 356 (1918); Eichler, Suppl. Fl. South Australia: 186 (1965); Polhill in Kew Bull. 22: 247 (1968); Chippendale in Proc. Linn. Soc. New South Wales 96: 239 (1972). HOLOTYPE: n.v., believed destroyed in B: Hermannsburg on the Finke R [Northern Territory-Central Australia], *Strehlow* 5, 1906–8. The description leaves little doubt of identity, and I here designate the following topotypic specimen as NEOTYPE: 16 km W. of Hermannsburg on the Areyonga road, *Latz* 389, 12.1968 (NSW).

[*C. dissitiflora* Benth. var. *eremaea* sensu auct., non Benth. sensu str.: Tate in Part III (Section II, Botany): 152 (1896) in Spencer, ed., Rep. Horn Exped. Centr. Australia. Tate here used this synonym of *C. eremaea* for that part of *C. eremaea* occurring in the habitat and localities cited, i.e. for subsp. *strehlowii*].

[*C. dissitiflora* sensu auctt., non Benth. sensu str.: Ewart & Davies, Fl. N. Territory: 141 (1917) in part, viz. "Glabrous form . . ."; Gardner, Enum. Pl. Austral. Occ.: 62 (1930); Beard, ed., (Descr. Cat.) W. Austral. Pl. ed. 2: 65 (1970). These authors here used this binomial for that part of the *dissitiflora* complex occurring in the habitat and localities cited, i.e. for *C. eremaea* subsp. *strehlowii*. Gardner's usage is implicit in his listing also *C. benthamiana*; Beard later, in Wildfl. Northwest: 9, 28 (c. 1972) used *C. dissitiflora* for *C. benthamiana*, regarding the Fortescue taxon referred to as a part of *C. dissitiflora*].

Characters as given in the key, differing from subsp. *eremaea* in its leaf form and pubescence, the leaflets often single, trullate or ovate, and glabrous or nearly so; the correlation between these characters, however, is by no means perfect though it applies in the majority of specimens; its breakdown produces the "intermediate" specimens showing associations of characters different from the usual.

DISTRIBUTION (see Map 1): Northwest of South Australia, cremean Western Australia and Northern Territory (Central Australia), i.e. in the western and northwestern part of the specific range, and showing combination with the Type subspecies chiefly along the line of junction in South Australia, northwards from Spencers Gulf, and in Central Australia: often on sandhills but also on sandplains and sandy soils near watercourses.

SPECIMENS EXAMINED: SOUTH AUSTRALIA: **Far North:** 20 miles [32 km] S. of Ernabella, *Turvey NSW* 113664, 4.1964 (NSW); 40 miles [64 km] N. of Oodnadatta, *Lothian* 1400, 8.1963 (NSW); 50 km N. of Oodnadatta, *Milthorpe* 3058, 10.1974 (NSW); Alberga R, *Sheaffe NSW* 45785, 5.1936 (NSW); "Nilpinna", 28° 29' S, 135° 55' E, *Milthorpe* 3051, 10.1974 (NSW); 16 miles [26 km] SE. of Emu, *Forde* 449, 8.1956 (PERTH).

WESTERN AUSTRALIA: **Ereman Province-Carnegie:** 3 miles [5 km] NE. of Walter James Range, N. of Rawlinson Range, *George* 8344, 10.1966 (PERTH); 69 miles [111 km] SE. of Giles Weather Station on Gunbarrel Highway, *Johnson* 73/4, 6.1973 (NSW); ± 100 miles [c. 161 km] E. of Warburton Mission, *George* 5283, 7.1963 (PERTH); Lake Carnegie, *Fairall* 2097, 7.1966 (PERTH)*; 38 miles [61 km] SE. of Warburton, *George* 2938, 8.1961 (PERTH); 36 miles [58 km] SE. of Warburton, *George* 3898, 8.1962 (PERTH); E. of Laverton, *Geol. Survey* 1916 (PERTH).

NORTHERN TERRITORY: **Central Australia:** Tanami Track, 107 miles [172 km] SE. of Rabbit Flat, *Johnson* 73/61, 6.1973 (NSW); near Ti Tree Creek, N. of Alice Springs, *Hutton NSW* 98221, 5.1967 (NSW); Boomerang Waterhole, Lander R, 42 miles [63 km] NW. of "Willowra" homestead, *Chippendale NT* 4733, 7.1958 (AD, NSW); 40 miles [64 km] N. of Barrow Creek, Stuart H'way [21° 04' S, 134° 10' E], *Must* 168, 7.1968 (NSW); Hanson R-Anningie turnoff, *Parker* 112, 7.1969 (NSW); 9.3 miles [15 km] W. of "Amaroo", *Chippendale NT* 2528, 8.1956 (NSW); 3.7 miles [6 km] SE. of Yuendumu, *Chippendale NT* 4191, 4.1958 (PERTH); "Gibeannie" Station [now abandoned], 125 miles [200 km] N. of Alice Springs, *Rose* 349 SN 52/244 (S)

* A trifoliolate plant occurring a vast distance further west than any similar collection; the lateral leaflets are less than half as long as the terminal but scarcely modified in shape.

(BRI); 5 miles [8 km] NW. of "Aileron", *Nelson* 386, 8.1962 (NSW); Carinya Bore, "Central Mount Wedge" Station, *McKee* 8579, 2.1961 (NSW); 13 miles [21 km] E. of "Alcoota" Station, *Perry* 3393, 3.1953 (BRI); Narwietooma, *Gardner* 11578, 3.1953 (PERTH, MEL [as *C. arida* ined.]); Ehrenberg Range, 60 miles [97 km] E. of Sandy Blight Junction, *Butler* 65, 5.1967 (PERTH); 32 miles [51 km] NW. of "Hamilton Downs", *Winkworth* 361, 6.1954 (BRI); North-west Stock Route, c. 6 miles [c. 10 km] NW. of Desert Bore, *Everist* 4173, 1.1950 (BRI); 2 miles [3 km] W. of No. 2 Desert Bore, "Hamilton Downs", *Chippendale NT* 1685, 9.1955 (NSW); in steppes near Arltunga, *Gauba CBG* 001355, 9.1950 (NSW, bis); Alice Springs, *Animal Industry Branch* 381 (PERTH); Alice Springs district, *Gardner* (PERTH); Standley Chasm, 40 miles [64 km] W. of Alice Springs, *Dale NSW* 45784, 7.1939 (NSW); 9 miles [15 km] SE. of Alice Springs, *Nelson* 137, 12.1961 (NSW); 19 miles [30 km] S. of Alice Springs, *Swinbourne* 399, 8.1962 (PERTH); Deep Well road, 19 miles [30 km] S. of Alice Springs, *Nelson NT* 1980, 5.1966 (AD, NSW); 20 miles [32 km] S. of Alice Springs on Deep Well road, *Cunningham & Mulham* 2045, 3.1974 (NSW); 33 miles [53 km] SW. of Alice Springs, *Chippendale NT* 492, 11.1954 (AD); Palm Valley, *George* 5002, 7.1963 (PERTH); Illara Creek, 4.5 miles [7 km] NW. of "Tempe Downs", *Chippendale NT* 7400, 10.1960 (NSW); 29 miles [47 km] E. of "Tempe Downs" homestead, *Chippendale NT* 572, 11.1954 (BRI, NSW, PERTH); Finke R, Stuart Highway, *George* 5049, 7.1963 (PERTH); Finke R at Henbury bridge, *Briggs* 3500, 6.1970 (NSW); Lake Amadeus, 24° 57' S, 131° 12' E, *Latz* 5713, 9.1974 (NSW); between Shaw and Docker Rivers, *Carolin* 5389, 8.1966 (SYD); Irving Creek, N. of Petermann Ranges, *Nelson* 942, 3.1964 (AD, NSW); 3 miles [5 km] W. of Lasseters Cave, Petermann Ranges, *Maconochie* 692, 1.1969 (NSW); c. 17 miles [c. 27 km] NW. of "Andado" homestead, *Must* 88, 7.1968 (BRI); 75 miles [121 km] E. of Ayers Rock, *Jancey* 5.1963 (SYD); Ayers Rock, *Woolnough & Johnson NSW* 113665, 8.1968 (NSW); Simpson Desert, 46 miles [74 km] SW. of "Andado" homestead, *Chippendale NT* 6608, 9.1959 (NSW, AD); Simpson Desert, 25° 47' S, 125° 15' E, *Mitchell* 99, 8.1974 (NT, NSW).

In addition, the following may be included in subsp. *strehlowii* (see Map 1, subsp. *strehlowii* atypical) though the diagnostic characters are less definite; this is shown chiefly in the development of leaflets usually narrower than in typical subsp. *strehlowii* and tending, sometimes strongly, to the oblong shape seen in subsp. *eremaea*, these shapes being combined with the glabrous or near glabrous and usually wrinkled surface of subsp. *strehlowii*, and sometimes in trifoliolate leaves:

SOUTH AUSTRALIA: Far North: "Macumba" Station, *Ising* 11.1950 (AD). **Lake Torrens Basin:** 8 miles [13 km] N. of "Parakylia", *Filson* 3251, 9.1960 (AD). **Lake Torrens Plateau:** c. 3 km N. of South Gap, *Weber* 1215, 9.1968 (AD). ? loc.: Alice Springs to Marree, *Ward* 8.1947 (SYD).

NORTHERN TERRITORY: Central Australia: 42 miles [63 km] NE. of Barrow Creek, *Winkworth* 957, 3.1955 (BRI, PERTH); 61 miles [98 km] from Jervois Mine towards Plenty R, *Gittins* 1983, 9.1969 (BRI, NSW); Benstead Creek, 32 miles [51 km] ENE. of Alice Springs, *Forde* 86, 2.1956 (NSW); 2.5 miles [4 km] E. of Atherrita Bore, "Todd R" Station, *Nelson* 1343, 8.1964 (NSW); Finke R valley (south of Alice Springs), *Hutton NSW* 98220, 5.1967 (NSW).

For other specimens of this species which do not fall clearly into either of the subspecies but show atypical combinations of their characters, see under subsp. *eremaea*, p. 337.

As with subsp. *eremaea* I examined, and determined as *C. strehlowii*, a number of specimens at AD in 1971, in addition to those cited here for subsp. *strehlowii*; they are similarly excluded from Map 1 but again support the distributions shown there.

2. *C. MITCHELLII* AND *C. SMITHIANA*

C. mitchellii, based on Mitchell's collection from near a lagoon on the Balonne R, Queensland, had at first a limited circumscription equivalent to that of the Type

subspecies distinguished below. Later Bentham enlarged it by including specimens of a smaller-leaved, less pubescent variant, the southwestern race, among his citations in *Flora Australiensis*; this is formally distinguished as subsp. *laevis* below. Still later Ewart described a Central Australian plant as *C. mitchellii* var. *tomentosa*, but the taxon based on Ewart's Type, while superficially similar in its flowers to *C. mitchellii*, is distinct in its vegetative parts (see Table 2) and comprises a taxon of eremean occurrence quite disjunct from that of *C. mitchellii*, which is coastal in Queensland and penetrates inland scarcely beyond the western slopes divisions of New South Wales. It is described below as *C. smithiana*.

TABLE 2
DIFFERENCES BETWEEN *C. MITCHELLII* AND *C. SMITHIANA*

	<i>C. mitchellii</i>	<i>C. smithiana</i>
Leaves	Simple, shortly petiolate to almost sessile, glabrous and dotted above, glabrous or hoary to quite densely pubescent with $\pm$ long hairs below.	Unifoliate, the leaflet shortly petiolulate and articulate on a longer petiole, the joint sometimes obscure in the pubescence, tomentose on both sides at first, later more sparsely hairy to glabrous above.
Stems	Erect, branched, glabrous or with short upturned hairs.	$\pm$ prostrate after the first, forming a cushion-like growth, densely spreading-pubescent with the hairs later usually downturned.
Stipules ..	Soon caducous.	Persistent.

2 i *C. mitchellii* Benth. in Mitch. Trop. Australia: 122 (1848); Bentham, Fl. Austral. 2: 181 (1864); F. Mueller, Fragm. 9: 156 (1875), in part; F. M. Bailey, Queensland Fl. 2: 374 (1900); Moore & Betche, Handb. Fl. New South Wales: 145 (1893); Domin in Biblioth. Bot. 22 (89²): 733 (1925) excl. var. *tomentosa* and occurrence in South Australia. HOLOTYPE: bed of Baloon [Baloone] R [Queensland-Maranoa], Mitchell, 10 April 1846, photograph only seen: (K, print at NSW of CSIRO Neg. No. Kew 1112). Two specimens marked * in the citations below were matched with the Type by L. Pedley.

[In references under *C. mitchellii*, other than those above, the name has often been used with some degree of misapplication, either by misdetermination or by changes in circumscription, as follows: by F. Mueller (1862) when describing *C. novae-hollandiae* q.v.; by Tate (1890), Black (1924, 1948), Chippendale & Murray (1958), the Poison Plants Comm., CSIR (1934), Beard (1970), and probably Gardner (1930), when referring in part or in various ways to *C. smithiana* q.v.].

Soft-wooded erect or decumbent shrubs about 0.5–1 m in height with glabrous or pubescent stems, the hairs rather short and ascending, and simple leaves. Leaves often elliptical or narrow-elliptical, glabrous and obscurely dotted above or sometimes on both sides and variously pubescent below, shortly petiolate to almost sessile with scarcely more than the pulvinus, with subulate to setaceous stipules often fallen from herbarium specimens; hairs of the pubescence from very short in subsp. *laevis* to softly curved and c. 1 mm long in subsp. *mitchellii*. Inflorescences terminal or

leaf-opposed, racemes with numerous, crowded flowers. *Flowers* deep yellow, pedicellate and bracteate. *Calyx* with teeth subequal and slightly longer than the tube, the tube slightly protracted on the lower side, the receptacle scarcely discernible. *Standard* more or less circular but often slightly broader than long, with a pair of plate-like calli, or linear converging thickenings, at the base of the lamina. *Wings* rounded-oblong. *Keel* with a rounded, right-angled bend about the middle, the beak twisted at the apex. *Ovary* glabrous, almost sessile, the style geniculate but scarcely constricted at the bend, the pod oblong-clavate, tapered to the base, the stipe lengthening to equal the calyx tube. *Seeds* 8–12 in the pod, slightly flattened, somewhat glossy, greenish grey, with a hooked radicular lobe.

Inflorescence usually less than 20 cm long, often 8–15 cm. *Flowers* 7–10 (–11) mm long, the pedicel slender, 1–3 mm, in the axil of a small narrow bract (1.5–) 2–3 (–4.5) mm long. *Calyx teeth* (2–) 2.5 (–5) mm, the tube (1.5–) 2 (–3) mm long, the receptacle to 1 mm. *Standard* (5–) 6–7 (–9) mm long, (5–) 9 (–11) mm broad, the claw 1–2 mm. *Wings* 5–8 (–9) mm long, 2.5–4 mm broad, the claw 1.3–2 mm. *Keel* (6–) 7–10 (–13) mm long, 3–4 (–5) mm broad, the claw 1.5–2 mm, the beak (4.5–) 6 (–8) mm long. *Pod* mostly 2–3 cm long, (6–) 8 (–10) mm in diam. *Seeds* c. 4 mm × 3 mm.

DISTRIBUTION (see Map 2): Eastern Queensland, and the northern half of the Coast, Tablelands and Western Slopes in New South Wales.

C. mitchellii comprises two geographical races, one coastal in the northern and northeastern parts of the specific range, the other further south, often on the western slopes of the Great Dividing Range in southern Queensland and New South Wales, the races overlapping with considerable intergradation in southeastern Queensland and northern New South Wales; the relevant specimens are cited separately and shown on the map.

1. Stems pubescent, very rarely glabrescent; leaves large, 3–13 cm long, 2–5 cm broad, usually elliptical, acute or acuminate at the apex and tapered at the base, pubescent with loosely and irregularly arranged hairs on the underside; flowers in a very dense inflorescence, often with a thick peduncle and rachis..... subsp. *mitchellii* a.
- 1.* Stems glabrous or hoary, occasionally short-pubescent; leaves smaller or at least narrower, often 4–6 cm long and 1–1.5 cm broad, commonly narrow-obovate or narrow-elliptical with a more or less retuse or at least obtuse apex, glabrous or hoary below; flowers more widely spaced on a more slender peduncle subsp. *laevis* b.

2 i a subsp. *mitchellii*.

Diagnostic characters as given in the key. Height of plants 0.3–1 m, occasionally to 1.3 m. *Leaves* large and soft, elliptical or narrow-elliptical (SAC 3–4, or 2), occasionally narrow-ovate (SAC 36–38) or nearly obovate (SAC 46–47), (4–) 5–8 (–14) cm long, (1.5–) 3–3.5 (–more than 5) cm broad; stipules 2–5 mm long. *Inflorescences* (6–) 8–13 (–18) cm long, on a rather thick peduncle up to 8 mm in diameter, with often more than 40 crowded flowers.

DISTRIBUTION: From NE. Queensland (Cook District) to the North Coast of New South Wales, largely in coastal districts and on adjacent plateaux but extending inland to the Carnarvon Range and Maranoa District in Queensland; often on sandy and other light-textured soils, frequently near streams or behind beaches, sometimes in open forest or weedy in rough pasture or cultivation.

Map 2. *Crotalaria mitchellii* and *C. smithiana*.

SPECIMENS EXAMINED: QUEENSLAND: **Cook:** Cairns, *Betche NSW 45978*, 8.1901 (NSW); Mareeba, *Darnell-Smith NSW 45977*, 4.1929 (NSW); Walsh R, c. 10 miles [c. 16 km] E. of Dimbulah, *Everist 5144*, 5.1952 (BRI); Parada, along Walsh R, *Keefer 3.1958* (BRI); Yungaburra, *White 1.1918* (BRI). **North Kennedy:** Barron Beach (? loc.), *Flecker 1614*, 4.1936 (BRI); 6 miles [10 km] from Ravenshoe on Mt Garnet road, *Davis 25*, 3.1943 (NSW); Blunder Creek . . . area, c. 15 miles [c. 24 km] SW. [? SE.] of Mt Garnet, *Johnson 4.1964* (BRI); "Meadowbank", 45 miles [73 km] S. of Mt Garnet, *Hoskin 5*, 2.1963 (BRI); Cardwell, *Michael 550* (BRI); 60 miles [97 km] SSE. of Mt Garnet, *Clague 4.1971* (BRI); Magnetic I, *White 2.1918* (BRI); Townsville, *White 8.1916* (BRI), NSW 45979 with flowering shoot from Brisbane R); Animal Health Station, Oonoonba, *S/N 50/138*, 4.1950 (BRI); 6 miles [10 km] W. of Ayr, *Freshwater 8.1964* (BRI); Bowen District, *Young* (BRI); "Range View" c. 30 miles [c. 48 km] SE. of Ravenswood, *Everist 5183*, 6.1952 (BRI); Edgumbe Bay, *Michael 1203* (BRI). **South Kennedy:** Perpetua Point, E. of Sarina, *Francis 4.1927* (BRI). **Leichhardt:** Carnarvon Gorge, 70 miles [113 km] NNW. of Injune, *McDonald 208*, 4.1966 (BRI); Carnarvon Creek, Carnarvon Range, *Butler 2.1960* (BRI). **Port Curtis:** beach c. 5 km N. of Yeppoon, *Rodd 1149*, 8.1970 (BRI); Lamamou Beach near Yeppoon, 23° 07' S 150° 45' E, *Beeston 3.1971* (BRI); Rockhampton, *Court 5.1925*, *Bailey* (BRI); Marmor, near Rockhampton, *Francis 3.1920* (BRI); Gladstone, *Hedley 1888* (BRI); Rosedale, *Dovey* (BRI). **Wide Bay:** Wide Bay, *Leichhardt NSW 114019*, 5.1843 (NSW). **Maranoa:** "Rowallan", 2 miles [3 km] NE. of Mitchell, *Martin 2.1962* (BRI); "Lagoonville", Orallo, *Hando 1.1918* (BRI); Bokkara [Bokhara] Flats, *Leichhardt NSW 115724*, 2.1847 (or 1848) (NSW). **Darling Downs:** Chinchilla, *Darnell-Smith NSW 45980*, 4.1926 (NSW, "good match for the Type"*; Moonie R, 2 miles [3 km] S. of Weranga, *Pedley 963*, 2.1962 (BRI). **Moreton:** Moreton I, *White Easter 1924* (BRI); Redcliffe (Moreton Bay), *Bick 1.1917* (BRI); Nudgee, *White 1.1918* (BRI); South Brisbane Cemetery, *Bailey (as F.M.B.) 3.1875* (BRI); South Brisbane, *herb. Simmonds 97*, 5.1889 (BRI); Brisbane R, *Dietrich 836*, 1863-5 (NSW); Cedar Creek, *White 6.1913* (BRI); "Binnaburra", Macpherson Range, *Johnson NSW 45981*, 5.1951 (NSW). A specimen "Mouth of Althaus Creek, about 4 miles NW. of Yabulu [Queensland-North Kennedy] *Smith T.131*, 9.1942" (BRI), seems to have the dimensions of subsp. *mitchellii* but is almost completely glabrous; it is perhaps a local littoral ("on low dune") ecotype of which no other material is known.

NEW SOUTH WALES: **North Coast:** Tweed Heads, *Forsyth NSW 45986*, 10.1898 (NSW); Tooloom Scrub, 15 miles [24 km] SW. of Urbenville, *Constable 6648*, 12.1965 (NSW); Kyogle, *Vane NSW 113906*, 3.1961 (NSW); Jackadgery, W. of Grafton, *Brooks NSW 113905*, 2.1963 (NSW); Grafton [district], *Glenfield Veterin. Res. Sta. SN 43/2404*, 12.1943 (NSW); Willawarrin, Upper Macleay R, *Noonan NSW 113904*, 3.1964 (NSW); *Toorooka, Upper Macleay R, *Noonan NSW 45975*, 4.1958 (NSW, "a reasonably good match for the Type"); Port Macquarie, *Brown NSW 45988*, 12.1897 (NSW); Kendall district, *Bailey 86*, 8.1929 (NSW); Craven State Forest [c. 13 km S. of Gloucester], *de Beuzeville NSW 45989*, 7.1917 (NSW).

2 i b subsp. laevis A. Lee, subsp. nov. HOLOTYPE: 1 mile [1.6 km] W. of Baradine [New South Wales-North Western Slopes], *P. Hodge NSW 84861*, 6.1967 (NSW).

A subspecies *mitchellii* pubescentia minore vel absenti, foliis minoribus vel saltem angustioribus apicibus plerumque obtusis vel retusis, et saepe floribus paullo maioribus dissitioribusque, differt.

Diagnostic characters as given in the key. Differs from subsp. *mitchellii* also in intergrading characters: height of plants often c. 60 cm. *Leaves* (30-) 51-65 (-80) mm long, (8-) 15 (-28) mm broad, usually narrow-elliptical (SAC 1-2) or a little broader, or occasionally obovate (SAC 46-47) or ovate (SAC 37); hairs of the pubescence grading in length and curvature from very short and nearly straight

* See Holotype paragraph.

to the condition in subsp. *mitchellii*; stipules often present, 1.5–3 mm long. *Inflor-escence* often with less than 30 flowers in the same length, the peduncle and rachis more slender than in subsp. *mitchellii*.

DISTRIBUTION (see Map 2): From the Maranoa and Moreton Districts of Queensland to the Northern Tablelands, North Western and northern part of the Central Western Slopes, the eastern edge of the North Western Plains and the southern and inland parts of the North Coast of New South Wales; often on sandy or granitic (but not highly deficient) soils though occasionally on basalt, in open forest or woodland, sometimes near streams and occasionally weedy in pastures or cultivation, not known from the sea coast.

SPECIMENS EXAMINED: **QUEENSLAND:** **Wide Bay:** Gympie, *Kenny* (BRI); Archers Station, Durundur near Beerwah, *Leichhardt* NSW 45959 in part, 1.1844 (NSW). **Darling Downs:** Bungunyah, *Everist* 849, 12.1934 (BRI); Warwick, *Boorman* NSW 45982, 3.1911 (NSW); 7 miles [11 km] S. of Warwick, *Pedley* 1159, 12.1962 (BRI); 15 miles [24 km] S. of Warwick on Stanthorpe road, *Everist & Webb* 1276, 11.1946 (BRI); Inglewood, *Jones (as W.T.J.)* 1189, 3.1959 (BRI); Thulimbah, *Schindler* (BRI); Stanthorpe, *Davidson* 2.1891 (BRI), *Boorman* NSW 45983, 11.1904 (NSW), *Hockings* 2, 1.1963 (BRI). **Moreton:** Esk, *Stone* 10.1961 (BRI); Ipswich, *Hall* 154, 1909 (BRI); near Logan Village, *McCarthy* 1.1922 (BRI); 10 miles [16 km] NW. of Beaudesert, *Henderson* 11.1964 (BRI).

NEW SOUTH WALES: **North Coast:** Gloucester, *Betche* NSW 45990, 1.1881 (NSW); Glendon & Singleton, Hunter R, *Leichhardt* NSW 45959 in part, 12.1842–1.1843 (NSW); Belford, Branxton District, *Brown* NSW 45994, 1.1954 (NSW). **Northern Tablelands:** c. 5 miles [c. 8 km] past the Summit Border Gate on road to Maryvale [? error for Maryland], *Morrow* 39, 12.1961 (BRI); Track to Bald Rock Mtn, 15 miles [25 km] N. of Tenterfield, *Constable* 2078, 3.1962 (NSW); Boonoo Boonoo, *Boorman* NSW 45972, 2.1905 (NSW); Tenterfield to Sandy Flat, *Maiden* NSW 45967, 12.1898 (NSW); Deepwater, *Fraser & Vickery* 1.1936 (SYD); Head of Gwydir R, *Leichhardt* NSW 45959 in part, probably 3–6.1843 (NSW); Mt Lindsay, *Rupp* NSW 45960, 4.1914 (NSW). **North Western Slopes:** Wallangra, *Boorman* NSW 45969, 11.1912 (NSW); Warialda, *Rupp* NSW 45974, 1.1907 (NSW); Terry Hië Hie, *Inspector of Stock* NSW 45962, 12.1910 (NSW); Pilliga Scrub, *Rolls* 56, 2.1974 (NSW); Dead Mans Waterhole, Pilliga Scrub, *de Beuzeville* NSW 45966, 11.1915 (NSW); Coonabarabran to Narrabri road, *Althofer* NSW 114108, 2.1971 (NSW); Teridgerie Creek, W. of Baradine, *Constable* NSW 45961, 3.1951 (NSW); Coolie Camp, Baradine, *Chippendale & Constable* NSW 18539, 5.1951 (NSW); Baradine, *Fraser* NSW 45970, 1.1936 (NSW), *Fraser & Vickery* 1.1936 (SYD), *Rupp* 12.1932 (SYD); first Moonbi hill, *Martin* 11.1967 (SYD, Pollen Sample 1388); Warrumbungles [Mts] near Coonabarabran, *Hartley* 36, 1.1966 (NSW); Coonabarabran, *Nicholson* NSW 45963, 2.1933 (NSW), *Betche* NSW 45993, 1.1883 (NSW); Coonabarabran to Timor Rock, *Salasoo* 2202, 1.1962 (NSW); Liverpool Plains, NSW 45965 (NSW). **Central Western Slopes:** Merrygoen, *Brown* NSW 45973, 6.1889 (NSW); Coolbaggie Creek, 2 miles [3 km] S. of Eumungerie, *Constable* 27, 12.1961 (NSW); Birriwa, *Bowman* NSW 116651, 3.1976 (NSW); Merriwa, *Martin* NSW 45971, 3.1911 (NSW); Narrromine, *Scott* NSW 45992, 2.1915 (NSW), *Hovenden* NSW 45968, 2.1922 (NSW); Dubbo, *Simmons* NSW 113907, 2.1964 (NSW). **Western Plains:** annual weed collected in a regeneration plot, Narraway State Forest [c. 32 km W. of Coonamble], *Hoffman* NSW 45958, 5.1957 (NSW).

A few specimens intergrading between the subspecies (see Map 2) have been observed, as follows:

QUEENSLAND: **Burnett:** Gayndah, *Kenny* 1913 (BRI). **Darling Downs:** "Goondarin", Texas, *Myring* NSW 84682, 4.1967 (NSW); Wallangarra, *Betche* NSW 45985, 12.1891 (NSW), *Boorman* NSW 45984, 4.1914 (NSW). **Moreton:** 18 km S. of Esk, *Pedley* 04003, 12.1972 (BRI).

NEW SOUTH WALES: **North Coast:** Bonalbo, *Marks* NSW 45995, 1.1915 (BRI, NSW); Barrington, *Vickery* NSW 121552, 2.1937 (NSW).

2 ii C. smithiana A. Lee, nom. et stat. nov.—C. mitchellii Benth. var. tomentosa Ewart in Proc. Roy. Soc. Victoria 19: 37 (1906); Ewart & Davies, Fl. N. Territory: 141 (1917). **HOLOTYPE:** between the Finke R and Lady Charlottes Waters [Northern Territory-Central Australia], *Kempe* 23, 1885 (MEL 1010370).

[*C. mitchellii* sensu auctt., non Benth.: Tate, Handb. Fl. Extratrop. S. Australia: 66, 223 (1890); Black, Fl. S. Australia 2: 306 (1924) and ed. 2, 2: 448 (1948); Gardner, Enum. Pl. Austral. Occ.: 62 (1930); Poison Pl. Committee, CSIR, Pamph. 49: 26 (1934); Chippendale & Murray, N. Territory Admin., Anim. Industr. Branch, Extens. Article 2 (II): 34 (1958); Beard, ed., (Descr. Cat.) W. Austral. Wildfl., ed. 2: 65 (1970). These authors here used *C. mitchellii* for that part of the *mitchellii-smithiana* complex occurring in the area they treated, viz. *C. smithiana*, although conflicting features of description and other implications are found in some of the references].

A soft-wooded subshrub with a strong tap root and, at first, an erect stem, later with several to many branches from near the base, ascending, procumbent or prostrate, often 30–50 cm long (including the short terminal inflorescences), greyish to bluish green due to the dense pubescence, drying pale, the hairs of the pubescence at first stiffly spreading then often downturned on the stems or loosely appressed on the leaves. *Stems* terete, sometimes with a fine, obscure ribbing under the dense, pale hairs. *Leaves* unifoliate, stipulate and petiolate, the petiole articulate and usually angled on to a short petiolule, the joint somewhat obscured by the pubescence, the stipules very narrow, acute, at first erect then recurved to show the glabrous inner surface, the outer surface pubescent with long, spreading hairs. *Leaflets* mostly obovate to elliptical (SAC 3–4, 47–48), variable on the one plant, tapered at the base, at first densely tomentose on both sides, later the hairs less stiffly spreading and clearly sparse on the upper side which is sometimes almost or quite glabrous and then obscurely dotted, the apex obtuse or slightly emarginate, rarely broad-acute. *Inflorescences* ascending or erect, terminal on the branches and maturing basipetally, the somewhat crowded flowers pedicellate and bracteate, the bracts like the stipules but a little broader, the pedicels with stiffly spreading hairs. *Flowers* rather small, yellow, often with some reddish colour at the centre. *Calyx* more or less symmetrical, the teeth and tube nearly equal, the teeth triangular, the whole sparsely clothed with stiffly spreading hairs shorter than those of the stem and leaf. *Standard* nearly circular, sometimes with fine, dark stripes around the eye becoming obscure in most dried specimens, and with a pair of horizontal or rarely oblique calli extending downwards, but doubtfully into the claw. *Wings* more or less oblong with a very small auricle. *Keel* shortly beaked and with only a very slight twist at the apex, shortly fringed along the free margins of the petals near the base, with a pair of shallow pouches below the auricles. *Ovary* completely glabrous, or rarely with a very sparse scattering of minute hairs near the base of the suture, tapered into a very short stipe. *Style* downcurved, sometimes protruding through the base of the keel, then curving sharply erect, somewhat flattened, with a two-rowed beard, the adaxial row evident, the other sometimes obscure. *Stigma* terminal, brush-like. *Pod* inflated, glabrous, with only a few seeds (c. 4) in the few fruiting specimens seen, more or less clavate or sometimes truncate at the apex, the walls thin but firm, tapering at the base into a stipe about as long as the calyx tube. *Seeds* oblique-cordiform, light brown, slightly reddish, with a hooked radicular lobe.

Petiole 1.6–6 mm long, the petiolule 1–2 mm; stipules 1.5–2.5 mm long. *Leaflets* (1–) 1.5–3 (–5) cm long, (0.6–) 1.3–2.5 (–3) cm broad. *Inflorescences* (3–) 5–10 (–12) cm long on peduncles of 1–2 cm. *Flowers* (4–) 6–8 (–10) mm long; pedicels 1.5–2.5 mm; bracts 1.5–2.5 mm long. *Calyx teeth* mostly 2–3 mm, about equal to the tube. *Standard* 5–7 mm long, 6–7 mm broad the claw 1 mm. *Wings* 5–6 mm long, the claw 1.5 mm. *Keel* 6.5–8 mm long, c. 3 mm broad, the claw slender, 1–2 mm, the beak (4.5–) 6 (–8) mm long. *Ovary* 4–5 mm, tapered into the

stipe, c. 1 mm long. *Pod* usually 1.5–2 cm long, 5–9 mm in diam. *Seeds* 3.5–4.5 mm long, 2.5–4 mm broad.

DISTRIBUTION (see Map 2): On sands and sandy loams, often coarse-textured, of hills, creek beds and banks in Central Australia [Northern Territory], and in adjacent parts of Queensland and South Australia, extending into the extreme northwest of New South Wales.

SPECIMENS EXAMINED: QUEENSLAND: Gregory North: Box Flat near Georgina R, C[amp] 21 Simpson Desert Exped. 1939, *Crocker* 7.1939 (AD); 4 miles [6 km] from "Thorner" [Hut] on Dajarra road, *Pedley* 1113, 10.1962 (BRI); "Carandotta" Station, *Everist & Smith* 158, 2.1937 (BRI); 5 miles [8 km] SE. of Boulia, *Pedley* 2010, 5.1966 (BRI); Boulia-Bedourie road, 5 miles [8 km] ex Boulia, *Johnson* 73/127, 9.1973 (NSW); Boulia-Bedourie road c. 20 miles [c. 32 km] S. of Boulia, per *Mahoney*, 6.1953 (BRI); Currawilla, *Everist* 4081, 8.1949 (BRI). **Mitchell:** c. 15 miles [c. 24 km] SW. of Yalleroi, *Smith & Everist* 915, 10.1940 (BRI); 2.5 miles [4 km] SE. of Blackall, *Everist* 2586 (BRI). **Gregory South:** 60 miles [97 km] NE. of "Noccundra", *Carolin* 4275, 8.1964 (SYD); 35 miles [56 km] S. of Coopers Creek on road to Tibooburra (20 miles [32 km] to "Orientos" Station), *Johnson* 51, 6.1972 (NSW). **Warrego:** "Curragh", near Cunnamulla, *Hubbard & Winders* 6242, 1.1931 (BRI); airfield 4 miles [6 km] W. of Cunnamulla, *Ebersohn* E 207, 4.1962 (BRI); "Carbean" near Cunnamulla, *White* 11800, 3.1941 (BRI).

NEW SOUTH WALES: Far Western Plains: "Thurlow Downs" to "Berrawinia Downs" in the Paroo R District, *Boorman* NSW 116467, 10.1912 (NSW); Toona gate NW. of Tibooburra, *Milthorpe & Cunningham* 3421, 6.1975 (NSW).

SOUTH AUSTRALIA: Far North: c. 38 miles [c. 61 km] E. of Dalhousie Springs, *Lothian* 1534⁹ 8.1963 (AD); R[iver] Stevenson, *Horn Exped. NSW* 45786, 8.1894 (NSW); Lower part of R[iver] Stevenson, *Tate?* No. 865, 1894 (AD); "Macumba" Station, *Ising* 11.1950 (AD); near Macumba Hill, "probably *Horn Exped.*" 5.1894 (AD); 38 miles [62 km] N. of camp on Strzelecki Creek, *Milthorpe* 3034, 10.1974 (NSW); Coopers Creek, 4 km downstream from Innamincka, *Briggs* 4619, 5.1972 (NSW); c. 60 miles [c. 95 km] N. of Marree on Clayton R, *Hill* 479, 11.1955 (AD); Clayton R, c. 50 km NE. of Marree, *Hill* 319, 7.1955 (AD).

NORTHERN TERRITORY: Central Australia: Boomerang Waterhole, Lander R, 42 miles [68 km] NW. of Willowra, *Chippendale NT* 4761, 7.1958 (NSW, AD); Stirling Sandhills, 160 miles [258 km] N. of Alice Springs, *Dale* NSW 45781, 9.1939 (NSW); Wycliffe to Taylor Creek, *Ewart* 6.1924 (MEL); Barrow Creek, *Chippendale NT* 906, 3.1955 (BRI, NSW); c. 2 miles [c. 3 km] NE. of "Ammaroo" homestead, *Chippendale* 8.1957 (AD); 19 miles [31 km] NE. of "Ooratippra" homestead, *Swinbourne* 577, 11.1962 (NSW); 9 miles [15 km] W. of "Ooratippra" homestead, *Swinbourne* 431, 9.1962 (NSW, MEL); 2 miles [3 km] N. of "Ti Tree Well" homestead, *Winkworth* 869, 3.1955 (BRI); Rockhole between Yuendumu and Cockatoo Creek [c. 270 km NW. of Alice Springs] (AD); Cockatoo Creek, *Cleland* 8.1931 (AD 95833039); 8 miles [13 km] NE. of "MacDonald Downs" Station, *Perry* 3405, 3.1953 (BRI, NSW, AD); "MacDonald Downs", Fraser R, *Cleland* 8.1930 (AD 95833048); Aileron Lagoon, *Chippendale NT* 4521, 6.1958 (MEL, AD); Irukaru Creek [tributary of Bunday R c. 145 km NE. of Alice Springs], *Cleland* 9.1930 (AD 95833047, bis); Plenty R road, 1 mile [2 km] N. of "Mt Riddock" homestead, *Nelson NT* 11981, 5.1966 (NSW, AD); Plenty R to Harts Range Depot, *Burbidge & Gray* 4520, 10.1955 (AD); 6 miles [10 km] NE. of "Derwent", *Winkworth* 309, 6.1954 (BRI); "Napperby", c. 100 miles [c. 161 km] NW. of Alice Springs, *Everist* 4207, 1.1950 (BRI); Alice Springs, *Tate* NSW 45790, 1894 (NSW, AD); 0.7 miles [1 km] N. of Bastard Bore, "Todd R" Station, *Maconochie* 469, 9.1967 (AD); 9 miles [15 km] W. of "Ringwood" homestead, *Chippendale NT* 4925, 9.1958 (BRI, NSW); Deep Well road, 13 miles [21 km] S. of Alice Springs, *Nelson* 1844, 1.1969 (BRI); Simpson Desert, 24° 29' S 135° 44' E, *Mitchell* 71, 8.1974 (NT, NSW); SE. of "Henbury", *Basedow* 5–9.1920 (AD 96212262).

This species is named in honour of the late Lindsay Smith of the Queensland Herbarium, who had also recognized but not published it as distinct from *C. mitchellii*.

3. *C. NOVAE-HOLLANDIAE* (including *C. CRASSIPES*)

In his treatment of *C. novae-hollandiae* DC. and *C. crassipes* Hook. in *Flora Australiensis*, Bentham divided the former into "three rather marked forms" and published their names as: a. *parviflora*, b. *oblongifolia* and c. *lasiophylla*. These can reasonably be considered to have the rank of forma. He also gave *C. oblongifolia* Hook. as a synonym of *C. novae-hollandiae*. He cited a number of specimens without referring these individually to the three forms, nor did he distinguish a Type *forma* in any way. From his inscription on the sheets at K however, it is possible to allocate each of these specimens to the appropriate form, and these are accepted as the Types of Bentham's names. However, the characters of flower size and leaf pubescence which Bentham used to differentiate his three forms are not satisfactorily correlated in the larger collections now available. The species is now seen to fall clearly into two geographic races, treated as subspecies below, and all except one of Bentham's cited specimens fall into the Type subspecies. That specimen belongs to subsp. *lasiophylla* and is designated as its Lectotype below, since among the Syntypes of f. *lasiophylla*, Bentham included other elements which are referable to subsp. *novae-hollandiae*.

The rather sparse collections of subsp. *novae-hollandiae* from the Northern Territory and Western Australia show some variation suggesting that further collecting may reveal an east-west trend within that subspecies.

I have seen only a few specimens to which the name *C. crassipes* could be applied, two from Western Australia whence came the Type, the others from Queensland and the Northern Territory, and they differ from *C. novae-hollandiae* only in being glabrous. One collector remarks on the similarity between his specimens of glabrous and pubescent plants from the same locality, but notes that the specimens were collected from different soils. The stipule character described by Bentham for "*C. crassipes*" is seen in some but not all of the glabrous specimens and Bentham does not mention it for "*C. novae-hollandiae*" though it is similar there. There is variation in the shape of the leaflets in the glabrous specimens, as there is in those with pubescence, and within each of two Queensland localities (Lawn Hill, Tower Valley) glabrous and pubescent plants have been examined which are similar in leaf shape, although there is some difference in leaf shape between plants from the two localities. Thus, it seems that glabrous individuals or populations may occur in various parts of the range of subsp. *novae-hollandiae* but that they do not show sufficient coherence or distinctness to warrant formal taxonomic recognition.

3 *C. novae-hollandiae* DC., Prodr. 2: 127 (1825); Bentham, Fl. Austral. 2: 181 (1864); F. M. Bailey, Queensland Fl. 2: 97 (1900); Domin in Biblioth. Bot. 22 (89²): 733 (1925). HOLOTYPE: cited as "in Nova-Hollandia orient." A specimen in G-DC is accepted as the Type. Its label reads: "Nouvelle Hollande côte orient. Mus. de Paris 1822". Photograph only seen, a print at NSW. In spite of this citation, *C. novae-hollandiae* does not occur on the east coast of Australia. My interpretation of the identity of the Type has been confirmed by L. Pedley (pers. comm.) after examination of the specimen at G.

See under the subspecies for synonymy.

[F. Mueller in *Fragm.* 3: 50 (1862) gives a description of *C. novae-hollandiae* (probably subsp. *novae-hollandiae*) erroneously under "*C. mitchelli*", as pointed out by Bentham, loc. cit. and Domin, loc. cit.]

Usually small shrubs to about 1.5 m in height, soft-wooded but probably perennial, with a more or less dense pubescence on some or most parts, or occasionally quite glabrous. *Leaves* unifoliate, the petiole and usually shorter petiolule articulate and angled, stipulate with sometimes small and obscure or deciduous stipules; stipules always present on young stems but deciduous from the lower leaves, narrow, acuminate, erect then recurved, pubescent outside, glabrous within, often leaving a ridge, like that from the back of the petiole, decurrent on the stem (as described by Bentham for *C. crassipes*). *Leaflets* very variable in size and pubescence (see under the subspecies). *Inflorescences* terminal and leaf-opposed, the often numerous flowers somewhat variable in size and opening in basipetal or irregular order. *Flowers* yellow (a few, very old, with blackish discoloration—? the "Fl. ex sicco vid. purpurascens . . ." of the Candolle's description), often fairly large, on ascending then recurved pedicels and subtended by small, acuminate bracts similarly erect then recurved. *Calyx* broadly campanulate with the tube slightly protracted on the lower side, the teeth subequal, narrow deltoid and often slightly longer than the tube, the whole variously pubescent or occasionally quite glabrous. *Standard* more or less circular but often slightly broader than long with a pair of plate-like calli at its base not extending into the claw, the apex sometimes broadly pointed. *Wings* rounded oblong. *Keel* with a rounded right-angled bend at or slightly below the middle, slightly but constantly twisted at the apex, usually with a pair of shallow folds or pouches near the auricle. *Ovary* glabrous to partly or entirely pubescent, the style more or less geniculate but scarcely constricted at the bend. *Pod* clavate or somewhat truncate, tapered into a short stipe. *Seed* yellow ochre, with a marked radicular lobe.

Leaflet often 3–5 cm long, 1–3 cm broad; petiole 5–15 mm, petiolule 1.5–5 mm; stipules 2–3 mm long. *Flowers* mostly 10–15 mm long, occasionally only 6.5–8 mm, on pedicels of c. 4–8 mm with minute bracteoles near the middle; bracts 1–3 mm. *Calyx teeth* (2.5–) 3–4 (–5) mm long, the tube (2–) 3 (–4) mm. *Standard* (6–) 12 (–14) mm long, 8–14 mm broad, the claw mostly 1.5–2.5 mm long. *Wings* (6–) 9–12 mm long, the claw 1.5–3 mm. *Keel* (6.5–) 12 (–15) mm long, c. 5–7 mm broad, the claw 4–6 mm, the beak (3–) 4–6 (–7) mm long. *Pod* usually 2.5–3.5 cm long and c. 1 cm diam. *Seeds* c. 4 mm long, 3.5 mm broad.

DISTRIBUTION (see Map 3): Northern Australia from Queensland, mostly west of the Great Dividing Range, to Western Australia and Central Australia.

The species comprises two geographic races: subsp. *novae-hollandiae*, in north-western Queensland and the "Gulf country" of Queensland and the Northern Territory, and in comparable parts of Western Australia, and subsp. *lasiophylla*, limited to inland areas and known chiefly from Central Australia but occasionally overlapping the territory of subsp. *novae-hollandiae* near the Gulf of Carpentaria. There is apparently little intergradation between the subspecies, and the only specimen difficult to allocate is from this junction area: NORTHERN TERRITORY—**Darwin & Gulf District**: Junction of McArthurs & Glyde Rivers, Rice 2002, 9.1975 (NSW), shown as intergrade on Map 3.

Map 3. *Crotalaria novae-hollandiae* (incl. *C. crassipes*)

1. Leaflets elliptical to ovate or narrower, tapered into the petiolule at the base or occasionally approaching a cordate base, variously pubescent but often with the upper surface less densely clothed than the lower and sometimes quite glabrous above or on both sides; in the northern and northeastern parts of the range of the species subsp. *novae-hollandiae* a.
- 1.* Leaflets triangular or with a more or less cordate base, not tapered into the petiolule, densely pubescent on both surfaces with a grey tomentum becoming golden to deep reddish brown with long drying; in the inland, southern part of the range of the species subsp. *lasiophylla* b.

3 a subsp. *novae-hollandiae*.

C. crassipes Hook., Hooker's Icon. Pl. 9: t. 830 (1852); Benth., Fl. Austral. 2: 182 (1864); Fitzgerald in J. & Proc. Roy. Soc. Western Australia 3: 158 (1916-7); Ewart & Davies, Fl. N. Territory: 140 (1917); Gardner, Enum. Pl. Austral. Occ.: 62 (1930). HOLOTYPE, photograph only seen: North-east coast of Australia [Western Australia], *Bynoe, Voyage of the Beagle* (K, print at NSW of CSIRO Neg. No. Kew 1116).

C. novae-hollandiae f. *parviflora* Benth., loc. cit. ISOSYNTYPES: (i) Upper Victoria R [Northern Territory-Victoria R], *Mueller* 12.1855 (MEL 1010376, "seen by Bentham"; annotated "*C. concinna* F. Muell." [ined.] in a photograph of the sheet at K; (ii) Alligator Point [Queensland-Burke], scripsit *Mueller* (? MEL 1010371, "seen by Bentham").

C. oblongifolia Hook., Hooker's Icon. Pl. 9: sub t. 830 (1852).—*C. novae-hollandiae* f. *oblongifolia* Benth., Fl. Austral. 2: 182 (1864). HOLOTYPE, photograph only seen: N.W. Coast of Australia [Western Australia], *Bynoe, Voyage of the Beagle* (K-Herb. Hook., print at NSW of CSIRO Neg. No. 1115).

C. novae-hollandiae f. *lasiophylla* Benth., loc. cit., in part (see also subsp. *lasiophylla*), viz. as concordant with the following ISOSYNTYPES: (i) Albert R [Queensland-Burke], *Henne* (MEL 1010374, "seen by Bentham"); (ii) Gulf of Carpentaria [as last], *Henne* (MEL 1010375, "seen by Bentham"); (iii) Gulf of Carpentaria [as last], *Landsborough* (MEL 1010377).

[*C. mitchellii* (as "*Mitchelli*") sensu auct., non Benth.: F. Mueller in *Fragm.* 3: 50 (1862); *C. crassipes* and *C. oblongifolia* are here said to be allied to subsp. *novae-hollandiae* under this misapplied name].

[*C. novae-hollandiae* var. *novae-hollandiae*; Chippendale in *Proc. Linn. Soc. New South Wales* 96: 239 (1972), nom. invalid.,—from the areas of occurrence given for this and for "*var. lasiophylla*" Benth., also listed, the names seen to have been confused, as well as being unpublished in varietal rank].

Diagnostic characters as given in the key; pubescence very variable in this subspecies, that of the stems moderately dense, greyish, of short simple hairs at first stiffly ascending, later curved upwards, becoming light brownish with drying and age, but rarely approaching the rich reddish brown seen in subsp. *lasiophylla* below, occasionally (in the form described as "*C. crassipes*") absent; rarely longer, spreading and downturned in zone of contact with subsp. *lasiophylla*. Leaflet narrow-oblong (SAC 13), narrow-ovate (SAC 37-38), narrow-elliptical (SAC 1-2) or almost narrow-trullate (SAC 55), or broader: elliptical, ovate to trullate, always tapered at the base except in a few specimens grading towards subsp. *lasiophylla*, the apex acute, obtuse or retuse; upper surface glabrous to densely pubescent, the lower always with some hairs except in the few completely glabrous specimens, and usually but not always more densely pubescent than the upper surface; minute dotting of both surfaces visible in sparsely pubescent or glabrous leaves. Calyx with rather sparse, appressed hairs. Recorded height of plants (40-) 45-60 (-75) cm.

DISTRIBUTION (see Map 3): Tropical Australia, widespread in the monsoon belt. Often in sandy soils and in burnt or disturbed weedy situations, also in red-brown loamy soils, some volcanics; not infrequently near streams.

SPECIMENS EXAMINED: **QUEENSLAND:** **Cook:** Kennedy R, *Hann Exped.* 380 (BRI); Lynbrook, Etheridge [railway] Line, *Towers* 4.1960 (BRI); Gilbert R, *White* 2.1922 (BRI, NSW 45776). **Burke:** Settlement Creek, *Brass* 48, 6.1921 (BRI); Settlement Creek, 35 miles [56 km] N. of Wollgorang, *Perry* 1200, 6.1948 (BRI); s. loc., *Gittins* 810, 6.1963 (BRI); Burketown (Q.)-Borroloola (N.T.) road, 25 miles [40 km] W. of Gregory R, *Johnson* 73/101, 8.1973 (NSW); Leichhardt R at "Floraville" crossing, *Pedley* 2098, 6.1966 (BRI); 17 km W. of Alexandra R on Cloncurry-Burketown road 18° 35' S, 140° 07' E, *Jacobs* 1290, 4.1974 (NSW); 12 miles [19 km] W. of Alexandra R crossing on Cloncurry-Burketown road, *Carolyn* 8856, 4.1974 (SYD, NSW); 3 miles [5 km] S. of "Armravnald", 17° 36' S, 139° 45' E, *Carolyn* 8929, 4.1974 (SYD, NSW); Lawn Hill, *Jensen* 69, 69A*, 5.1940 (BRI); 10 miles [16 km] SSE. of "Morstone" Station, *Perry* 1055, 5.1948 (BRI); 6 miles [10 km] S. of "Boomarra" Station, *Perry* 3985, 8.1953 (BRI); "Chudleigh Park" Station, 110 miles [117 km] N. of Hughenden, *Hubbard & Winders* 7605, 2.1931 (BRI); c. 20° 00' S, 139° 50' E, on "Calton Hills" Station c. 40 miles [c. 64 km] N. of Mt Isa, *Brown* 8.1968 (BRI); 3 miles [5 km] N. of "Zingari" Station, *Speck* 4788, 8.1954 (BRI); Williams R, *Macgillivray* 2143, 8.1928 (BRI); 40 miles [64 km] ENE. of Richmond township, *Speck* 4477, 6.1954 (BRI)*; "Wongalee", c. 50 miles [c. 80 km] N. of Hughenden, *Hill* 29, 30, 6.1968 (BRI); Cloncurry, *Scarth-Johnson* 530, 8.1970 (BRI); W. of Mt Isa, *Beadle NSW* 115727, 9.1972 (NSW); Mt Isa, *Morris* 5.1952 (BRI); Mt Isa district, *Greig* 11.1956 (BRI); Hann Highway, 4 miles [6 km] N. of Hughenden, *Hill* 1, 6.1968 (BRI); Galah Creek, Hughenden, *McCarthy* 89, 6.1933

(BRI); "Hughenden" Station, *Inspector of Stock* 5.1954 (BRI)*; Tower Valley, Hughenden, *Sewell* 5.1954 (BRI 126637, 004829*); 20 miles [32 km] NW. of McKinlay, *per Forestry Commission of N.S.W.* 2334 (NSW); 6 miles [10 km] W. of Mary Kathleen, *Trapnell* 16, 7.1958 (BRI); between Duchess and Mt Isa, *Gittins* 726, 5.1963 (BRI; NSW as: 2 miles [3 km] W. of Duchess); 100 miles [161 km] S. of Cloncurry on the Longreach road, *Jancey* 5.1963 (SYD). **North Kennedy:** "Blue Range" Station via Ingham, *per Newton* 3.1953 (BRI); Ravenswood Junction, *Hubbard & Winders* 6954, 1.1931 (BRI). **Gregory North:** 44 miles [71 km] SSE. of Mt Isa on the Dajarra-Mt Isa road, *Williams* 177, 10.1968 (BRI); "Ardmore", *Everist* 3188, 11.1947 (BRI); Dajarra, *Everist & Smith* 139, 1.1937 (BRI); "Carandotta" Station, *Everist & Smith* 158, 2.1937 (BRI). A depauperate specimen: "Rockmount", Helidon [Moreton], *Klose* 7.1963 (BRI) seems likely to have been introduced into this locality.

NORTHERN TERRITORY: **Darwin & Gulf District:** 7 miles [12 km] E. of Rum Bottle Creek, 12° 04' S, 133° 48' E, *Macconchie* 1591, 6.1972 (NSW)*; Spring Vale, Port Darwin, *Giles* (BRI); above UDP Falls, *Gittins* 2701, 8.1973 (NSW)*; 41 miles [66 km] from Pine Creek → UDP Falls, *Gittins* 2569, 7.1973 (NSW)*; 6 miles [10 km] N. of Katherine, *McKee* 8526, 2.1961 (NSW); Stuart Highway c. 3 miles [c. 5 km] NW. of Katherine, *Adams* 881, 2.1964 (NSW); 30.2 miles [48.6 km] S. of Katherine town, *Chippendale NT* 3738, 9.1957 (NSW, MEL); Bryans Waterhole, 1 mile [1.6 km] off Carpentaria Highway, 18 miles [29 km] S. [E. by S. or W. by S.?] of Borroloola, *Johnson* 73/117, 8.1973 (NSW); 36 miles [58 km] E. of "O.T. Downs" homestead, *Chippendale NT* 5529, 3.1959 (NSW); Tablelands Highway 9 miles [15 km] S. of Carpentaria Highway, *Johnson* 73/122, 8.1973 (NSW). ? **Barkly Tableland:** 3 miles [5 km] W. of Calvert R crossing, 17° 13' S, 137° 22' E, *Carolyn* 9257, 5.1974 (SYD, NSW). **Victoria R District:** N.T. Kimberleys, 14 km W. of Victoria R crossing, *Symon* 10337, 6.1975 (NSW); 13.6 miles [22 km] S. of "Wave Hill" homestead, *Chippendale NT* 5761, 4.1959 (NSW).

WESTERN AUSTRALIA: **Northern Province-Hann:** Barnet R Gorge, 86 km S. of Gibb R Crossing, *Symon* 10264, 5.1975 (NSW)*; 14 km N. of Napier Range, 17° 11' S, 124° 51' E, *Symon* 10139, 5.1975 (NSW)*. **Ord:** Around Kununurra, Ord Dam & Wyndham road, 15° 45' S, 128° 44' E, *Scarth-Johnson* 548, 9.1970 (BRI); Ord R, *Fitzgerald NSW* 45777 & 45778, 10.1906 (NSW); in Ord R, 69 miles [111 km] S. of Kimberley Research Station, *Perry* 2961, 7.1952 (NSW, BRI); 17 miles [27 km] NE. of "Springvale" Station, *Lazarides* 5062, 4.1955 (BRI). **Fortescue:** Nickol Bay, scripsit *Maitland Brown* (MEL 1010373 "seen by Bentham", and MEL 1010372); W. Australian Hamersleys, Yampire Gorge, *Symon* 10059, 5.1975 (NSW).

* Glabrous specimen ("C. crassipes").

3 b subsp. lasiophylla (Benth.) A. Lee stat. nov.—*C. novae-hollandiae* f. *lasiophylla* Benth., Fl. Austral. 2: 182 (1864) as to the Lectotype (see also subsp. *novae-hollandiae*). LECTOTYPE, here designated: near Mt Humphries, 25° 38' [Northern Territory-Central Australia], Macd. Stuart (MEL 1010378, "seen by Bentham").

[*C. novae-hollandiae* var. *lasiophylla* Tate, Part III (Sect. II, Botany): 152 (1896), in Spencer, ed., Rep. Horn Exped. Centr. Australia, nom. invalid.; Chippendale in Proc. Linn. Soc. New South Wales 96: 239 (1972)—as for var. *novae-hollandiae* above, p. 350].

[*C. novae-hollandiae* sensu. auct., non DC. sensu str.: Black, Fl. S. Australia 2: 306 (1924) and ed. 2, 2: 448 (1948). Black here used the binomial for the only element of the species occurring in South Australia, viz. subsp. *lasiophylla*].

Diagnostic characters as given in the key; differs from subsp. *novae-hollandiae* also in its denser and more spreading tomentum, at first greyish, drying golden to deep reddish brown, the hairs c. 1 mm long, spreading and obliquely up- and down-turned in a dense mass on stems and leaves; stem hairs occasionally as in subsp. *novae-hollandiae* (in zone of contact). *Leaflets* with a characteristic truncate or slightly cordate base, more or less triangular (SAC 75–76), the apex obtuse or slightly retuse, very variable in size on the one plant. *Flowers* and their parts like those of subsp. *novae-hollandiae*, but the calyx more densely pubescent with rather longer hairs.

DISTRIBUTION (see Map 3): On sandy soils chiefly in eremean areas of the Northern Territory and in adjacent parts of Western Australia and Queensland; in the north of its range occasionally overlapping the territory of subsp. *novae-hollandiae*; perhaps to be expected in the far north of South Australia also.

SPECIMENS EXAMINED: QUEENSLAND: **Burke:** Gregory R Crossing, c. 20 miles [c. 32 km] SW. of Burketown, *Pedley* 2067, 6.1966 (BRI, K); Doomadgee Mission, *Whitehouse* (BRI), 17° 56' S, 138° 43' E, *Jacobs* 1461, 5.1974 (NSW). **Gregory South:** Simpson Desert and many areas inland, *Cockburn BPS* 53, 7–8.1969 (BRI).

NORTHERN TERRITORY: **Barkly Tableland:** Settlement Creek, *Brass* 343, 5.1923 (BRI); 18 miles [29 km] W. of Stuart Highway, Top Spring beef road, *Must* 454, 2.1969 (BRI); 10 miles [16 km] N. of Elliott, Stuart Highway, *Latz* 95, 1.1968 (BRI); 17 miles [27 km] N. of "Helen Springs" Station, *Perry* 1908, 8.1949 (BRI); 15 miles [24 km] S. of Elliott, *Byrnes* 1441, 3.1969 (NSW); Renner Springs 18° 20' S, 133° 47' E, *Williams* 81, 7.1975 (NSW); "Banka Banka" Station, *Allen* 661, 8.1922 (NSW); 50 miles [80 km] N. of Tennant Creek, *Perry CANB* 603, 4.1948 (NSW, BRI); Gibson Creek, 19° 12' S, 134° 08' E, *Must* 221, 7.1968 (NSW); 124 miles [199 km] W. of Camooweal in N.T., per N.S.W. Forestry Commission 2330, (NSW). **Victoria River District:** 40 miles [64 km] S. of Hookers Creek, *Chippendale NT* 2329, 7.1956 (NSW); 34.7 miles [55 km] NE. of Tanami, *Chippendale* 4.1959 (MEL). **Central Australia:** 15.4 miles [24.8 km] E. of The Granites, *Chippendale NT* 4275, 5.1958 (NSW, MEL); 69 miles [111 km] NW. of Willowra, *Chippendale NT* 4808, 7.1958 (NSW, MEL); "Numagalong" homestead, *Nelson & Swinbourne* 32, 8.1965 (NSW, MEL); 14 km S. of Devils Marbles, 7 km N. of Hurst Creek, *Symon* 10373, 6.1975 (NSW); "Singleton", 240 miles [386 km] N. of Alice Springs, *Everist* 1.1950 (BRI); 7 miles [11 km] S. of Wycliffe Well, *Rose* 11.1949 (BRI); "North Murray Downs", *Filson* 234, 5.1957 (MEL); Barrow Creek, *Baldwin Spencer* 1902 (MEL); "Central Mount Stuart", *Ewart* 6.1924 (MEL); 10 miles [16 km] S. of Wycliffe Well, *Everist* 4274, 1.1950 (BRI); 10 miles [16 km] E. of "Mt Riddock" homestead, *Winkworth* 606, 10.1954 (BRI); 40 miles [65 km] SE. of Alice Springs, Santa Teresa road, *Latz* 113, 1.1968 (NSW); 1 mile [1.6 km] N. of Renners Rock, *Winkworth* 34, 2.1954 (BRI, MEL); "Renners Rock" homestead, *Winkworth* 765, 11.1954 (BRI); Hugh R, 12 miles [19 km] NW. of "Maryvale" homestead, *Nelson NT* 11985, 4.1966 (NSW); 0.6 miles [1 km] N. of Deep Well Siding, *Nelson* 1270, 8.1964 (NSW); 2 miles [3 km] N. of Bandooma Siding, *Swinbourne* 419, 8.1962 (NSW); NW. Simpson Desert, *Cockburn BPS* 24, 7–8.1969 (BRI); 7.3 miles [11.7 km] SE. of Bottom Bore, Hale R, *Chippendale NT* 4931, 9.1958 (BRI, NSW).

WESTERN AUSTRALIA: **Ereman Province-Carnegie:** 123 km WNW. of Tanami towards Billiluna, 128° 30' S, 190° 40' E, *Gittins* 2324, 7.1971 (NSW); 78 miles [125 km] WNW. of Tanami 19° 50' S, 129° 50' E [? error for 128° 50' E], *Dunlop* 2333, 9.1972 (NSW).

4. C. MEDICAGINEA OR C. TRIFOLIASTRUM

Without adequate exotic material for comparison, it is impossible to evaluate critically the taxa denoted by these names, but at the Kew Herbarium, where there are larger collections from regions outside Australia *C. medicaginea* is recognized as a species distinct from *C. trifolium*; also, in the most recent publication involving both names, de Munk (1962, below) accepts them as two species and introduces a new diagnostic character, viz. auricles present on the standard in *C. trifolium*, absent in *C. medicaginea*.

In the past, however, other opinions have been held and published. The normal format of "synonymy" (which would be studded with "in part" qualifications), and "misapplied names", is abandoned here in favour of a chronological history of treatment and nomenclature of the plants involved.

The Australian plants referred to one or other of these names are represented in the Banks and Solander collection of 1770 (as *C. trifolium*): NSW 133329 and 113929 from Thirsty Sound, Queensland, and the taxon, variable in itself, can therefore be accepted as native; nevertheless the possibility cannot be ruled out that other, morphologically different, plants were introduced subsequently. The legitimate names since applied to the taxon or taxa were not published for some years after that collection:—

- 1786 *Crotalaria medicaginea* Lamk., *Encycl. Méth.* 2: 201.
- 1802 *Crotalaria trifolium* Willd., *Sp. Pl.* 3: 983.
- 1843 Bentham (in *London J. Bot.* 2: 578) noted that earlier writers had already confused the taxa. He recorded an Australian collection ["North Coast?"] by Bauer under *C. neglecta* Wight et Arn., not *C. medicaginea* or *C. trifolium*, and described a new species *C. luxurians* close to *C. neglecta* in flowers (but with these more numerous, and habit erect and more robust), from India.
- 1862 Mueller (in *Fragm.* 3: 56) referred specimens (Keppel Bay to Arnhem Land) to *C. medicaginea*, giving *C. trifolium* as a synonym. He repeated this treatment in *Fragm.* 9: 157 (1875) for other specimens from Broad Sound, Peak Downs and Springsure.
- 1864 Bentham (in *Fl. Austral.* 2: 183) referred Australian plants (from northwestern Australia to Queensland) to *C. trifolium* and specified distinctions from *C. medicaginea*. He made no mention here of *C. neglecta*.
- 1876 Baker (in *Hook. f., Fl. Brit. India* 2: 81) maintained *C. medicaginea* and *trifolium* as species and reduced *C. neglecta* and *C. luxurians* to varietal rank under *C. medicaginea*.
- 1893 Moore and Betche (in *Handb. Fl. New South Wales*: 145) referred New South Wales plants to *C. medicaginea* with *C. trifolium* as a synonym.
- 1916 Maiden and Betche (in *Census New South Wales Pl.*: 106) followed the last.
- 1925 Domin (in *Biblioth. Bot.* 22 (89²): 734) referred Queensland plants to *C. medicaginea*, setting up two varieties to accommodate variation, including in the synonymy of one of them, var. *australiensis* Domin, a reference to Bentham's alleged misuse of the name *C. trifolium* in *Flora Australiensis*. Domin considered this variety, the most common Australian form, close to *C. luxurians* Benth. in habit, and to *C. trifolium* in its often multiflorous inflorescences.

- 1962 de Munk (in Reinwardtia 6: 208) referred Malaysian plants, and a few specimens examined from Australia, to *C. medicaginea*, maintaining it as distinct from *C. trifoliatum*, which latter he considered not to occur in Malaysia.
- 1963 Backer (in Fl. Java 2: 584) referred Javanese plants to *C. medicaginea* f. [sic] *luxurians* Baker, and did not include *C. trifoliatum* as Javanese.

My interpretation of the foregoing is as follows: the name *C. trifoliatum* is at present applied outside Australia to an erect, relatively large-flowered subshrub of limited geographical range in India, while *C. medicaginea* (including *C. herniarioides* Wight et Arn., *C. neglecta* and *C. luxurians*) is accepted for plants of varying habit, with usually (but not always) smaller flowers, distributed over a wide area from India through Malaysia to China and Australia and with the "*luxurians*" element extending farthest to the south and southeast, perhaps continued into Australia as Domin's var. *australiensis*. Until a critical appraisal of the whole situation is undertaken, I propose to use the name which has been most commonly used in publications by Australian botanists and which is indicated by the most recent work, i.e. *C. medicaginea* Lamk. In recent years most of the collections from tropical Australia have been distributed as *C. trifoliatum* Willd., although the name *C. medicaginea* is in use at the Queensland Herbarium, where it is hoped that some investigation into variation in the Australian populations will be undertaken, since there is a large representation of this taxon in Queensland. Several variants have already been distinguished by L. S. Smith (1962), though under the name *C. trifoliatum*. In New South Wales there are relatively few of these plants, and critical distinction of the variants is not considered as important as it is in northern Australia, where stock-poisoning properties have been found in some. I am greatly indebted to L. Pedley for his painstaking determinations of numerous specimens of this group submitted to him while at Kew. They included many of the known variants from States other than New South Wales and all were considered to belong to *C. medicaginea*.

DISTRIBUTION: Widely distributed over northern Australia; in New South Wales occasional in drier parts of the North Coast, away from the sea, and one record from the North Western Slopes.

5. SPECIES NATURALIZED OR ESCAPED FROM CULTIVATION IN NEW SOUTH WALES

The species naturalized or adventive in New South Wales are grouped in the following list under the Sections enumerated in his recent classification of the African species by Polhill (1968), and a few brief notes are included where there have been recent changes in nomenclature.

Section *Grandiflorae* (Bak. f.) Polhill

C. agatiflora Schweinf.—cultivated and occasionally escaped.

Section *Incanae* (Benth.) Polhill

C. incana L. subsp. *incana* } both subspecies naturalized
subsp. *purpurascens* (Lamk.) Milne-Redh. } for many years.

C. goreënsis Guill. et Perr.—naturalized in Queensland, the Northern Territory and the Kimberleys of Western Australia, and appearing recently, probably as impurities in seed, in newly established "tropical" pastures in the North Coast of New South Wales.

Section *Macrostachyae* (Benth.) Polhill

C. lanceolata E. Mey.—naturalized.

C. pallida Ait. var. *obovata* (G. Don) Polhill—naturalized. This taxon has previously been identified at the National Herbarium of New South Wales by the synonyms *C. striata* DC. and, later, *C. mucronata* Desv.; earlier it was misidentified as the distinct species *C. saltiana* Andr. (see Senn in *Rhodora* 41: 356 (1939).

C. zanzibarica Benth.—naturalized. A few specimens were formerly identified as *C. usaramoensis* Bak. f., now regarded as a synonym.

Section *Calycinae* Wight et Arn.

C. juncea L.—an apparently temporary occurrence on the Clarence River at Upper Copmanhurst in 1910, but not collected since in New South Wales.

Section *Crotalaria*

C. grahamiana Wight et Arn.—naturalized. This species has previously been misdetermined in New South Wales as *C. quinquefolia* L.

C. semperflorens Vent.—cultivated and occasionally escaped.

C. spectabilis Roth—naturalized. For many years this species was confused with the native species *C. retusa* L., but our earliest specimen from a New South Wales locality is dated 1912 and described as a naturalized weed.

C. distans Benth.—apparently cultivated and occasionally escaped.

C. macaulayae Bak. f.—a single collection said to have been introduced as an impurity in seed.

C. virgulata Klotsch—cultivated and occasionally escaped.

ACKNOWLEDGEMENTS

I am indebted to the Directors of the National Herbarium of Victoria and the Queensland Herbarium, and to the Keeper of the State Herbarium of South Australia, for valuable loans. In 1974 I was able to examine material at the Kew Herbarium, including the Type of *C. dissitiflora*, while assisted by a grant from the Australian Biological Resources Study; thanks are also due to the Director and staff of the Kew Herbarium, and especially to Dr R. M. Polhill.

The abundance of material made available to me has permitted some confidence in the sorting adopted and in the nomenclature applied, and this confidence had been increased by the general agreement of Mr Les. Pedley of the Queensland Herbarium, who had intended to continue the unpublished work of his colleague, the late Mr Lindsay Smith. However, in view of my own progress at the time of Smith's death, Pedley generously assisted me with determinations, information and discussion in correspondence, while serving as Australian Botanical Liaison Officer at Kew. After the death of Smith in 1970, I found that he had been pursuing investigations similar to mine over a number of years, and had incorporated similar conclusions in a key constructed for use in the Queensland Herbarium. It is a pleasure to acknowledge

the support given by his work, even though my treatment differs somewhat from his, particularly in nomenclature. Smith's greater experience in the field yielded information about soil types, the importance of which I had not at that time fully recognized, nor had I looked closely into subdivisions of the species *C. linifolia* and *C. medicaginea* which figure in his key, and these last problems, less pertinent to the Flora of New South Wales, remain for his successor.

I am also indebted to Dr L. A. S. Johnson for a critical review of the content of this paper, and for many constructive suggestions for the improvement of its presentation.

The figures were prepared for publication by Christine Payne and the maps by Rhonda Barry.

LITERATURE CITED

- Jackson, E. A., 1962—Soil Studies in Central Australia. CSIRO Soil Publication No. 19.
- Litchfield, W. H., 1969—Soil Surfaces and Sedimentary History near the Macdonnell Ranges, N.T. CSIRO Soil Publication No. 25.
- Polhill, R. M., 1968—Miscellaneous Notes on African species of *Crotalaria* L. II. Kew Bull. 22: 169–348.
- Smith, L. S., 1962—Unpublished key to *Crotalaria*, in BRI.
- Stephens, C. G., 1961—The Soil Landscapes of Australia. CSIRO Soil Publication No. 18.
- Stephens, C. G., 1963—Soils, 2nd. ed.: 18, in Atlas of Australian Resources, 2nd. Series. Canberra.
- Systematics Association Committee, 1962—Descriptive Terminology, Chart 1a. Taxon 11: 246–247.