Original Article

Applying of theory of planned behavior to promote physical activity and exercise behavior among older adults

SEYEDEH AMENEH MOTALEBI¹, JAMILEH AMIRZADEH IRANAGH.^{1,2}, ABBAS ABDOLLAHI³, WAI KONG LIM⁴

¹Institute of Gerontology, Universiti Putra Malaysia, MALAYSIA

² Faculty of Health, Urmia Medical Sciences University, IRAN

³ Faculty of Human Ecology, Universiti Putra Malaysia, MALAYSIA

⁴ Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, MALAYSIA

Published online: December 30, 2014

(Accepted for publication December 15, 2014)

DOI:10.7752/jpes.2014.04087;

Abstract:

Regular physical activity (PA) is foremost important for well-being and independent living in older age. However, recently, more than 75% of older adults have not been involved in any PA/exercise program at the suggested levels. Understanding their perspective towards PA/exercise provide important information in development of interventions for improving their exercise behavior. A conceptual framework that has been successfully applied for exercise behavior is the theory of planned behavior (TPB). The TPB has well shown the relationship between attitudes and PA/exercise behavior directly or indirectly through intention. Attitude, subjective norms and perceived behavioral control are three independent concepts of this theory. In fact, positive attitude, better perceived social pressure, and greater behavioral control result in stronger intention to perform a given behavior. PA/exercise intention is explained more by the TPB constructs in older compared to younger adults. Advancing age is associated with reduction in PA/exercise intention; however, the translation of intention to behavior increases. Furthermore, subjective norm and perceived behavioral control are considered as the strongest determinants of PA/exercise intention among older adults. Hence, the interventions targeting these factors will be valuable in promoting exercise behavior in the elderly.

Key words: aging, exercise intention, behavior, psycho-social theory

Introduction

The aging of the world's population is a most important demographic phenomenon of the 21st century (Armitage & Conner, 2001; Ahmad et al., 2011). It is estimated that by the year 2050 one in every five persons will be old (Mujahid, 2006). Population ageing has become an important challenge for both developed and developing countries (Ahmad et al., 2011), particularly in relation to their health (Nakasato & Carnes, 2006; Singh & Hiatt, 2006; Lin et al., 2010). With advancing age, age-related illnesses, particularly chronic and degenerative conditions and their related disabilities become more obvious (Schulz et al., 2004). In this view, the major public health targets are reducing of frailty and maintaining functional independence in older adults (Landi et al., 2010).

Regular physical activity (PA) or exercise has many lifelong health benefits (Netz et al., 2005; Lautenschlager et al., 2008). It is linked to prevention of chronic diseases (Schutzer & Graves, 2004; Kruk, 2007; Nigg et al., 2009), falls (Heesch et al., 2008), and disability (Wallin et al., 2008) in older age. Furthermore, it is advocated for preventing and optimally managing cardiovascular risk factors and reduction of heart-related death (Akbaratabartoori et al., 2005; Sofi et al., 2008). Therefore, the more physically active old people are, the less probably to die early (Joseph et al., 2013). Regular PA plays a key role in maintaining and improving the quantity and quality of life (Elavsky et al., 2005; Guedes et al., 2012; Pernambuco et al., 2012). It is also well documented as an essential part of successful aging (Ni Mhaoláin et al., 2012). Generally, there is no other group who can benefit of exercise more than the old population (Breen et al., 2007).

Despite of the overwhelmingly PA-related health benefits, only a small number of old people meet the standard activity level of 30 to 60 minutes per day (Rahimi et al., 2011) and a high proportion of them are not physically active (Vallance et al., 2011). The prevalence of sedentary lifestyle increases with ageing process (Dumith et al., 2011), and it is one of the most important predictors of functional disability among the older adults (Landi et al., 2010). As, in 2010 more than three million deaths around the world were due to an inactive lifestyle (Bull & Bauman, 2011).

Theories are useful for explaining the determinant factors of PA/exercise involvement (Plotnikoff et al., 2010). Despite the number of available psychological theories, there is no agreement for the preeminent model for studying exercise behavior (Maddux & DuCharme, 1997). However, one conceptual framework that has been successfully applied for exercise behavior is theory of planned behavior [TPB; Ajzen (1991)]. The TPB has been applied to predict and understand PA/exercise behavior in different groups of people (Vallance et al., 2011). According to this theory, attitudes and PA behavior are correlated via intention (Grodesky et al., 2006; Vallance et al., 2011). A strong intention is required for being physically active (Stadler et al., 2009).

Given older adults health status gets to be a concern; therefore, a better understanding of the social cognitive correlates of PA/exercise behavior among them could ease the development of proper strategies on PA/exercise change in this group of population. This article, therefore, address the TPB application in understanding PA/exercise intention and behavior in older adults based on some previous studies.

Theory of Planned Behavior (TPB)

This social cognitive theory shows the relationship between attitudes and behaviors. It assumes that individual's attitude towards outcomes of a behavior is an important factor for engaging in that behavior (Kosma, 2012). The TPB (Ajzen, 1991) is an extension of the theory of reasoned action (Fishbein & Ajzen, 1975) by adding the concept of perceived behavioral control to give an explanation for behaviors that are not under a person's control (Ajzen, 1991; Benjamin et al., 2005).

The TPB seems to be a prominent theory for the prediction of health-related behaviors (Armitage & Conner, 2001; Conner & Sparks, 2005; Plotnikoff et al., 2008), such as smoking (Higgins & Conner, 2003), dieting (Nejad et al., 2005), and exercise (Hausenblas et al., 1997; Downs & Hausenblas, 2005; Rhodes et al., 2006). It combines social influences and personal factors as the behavior predictors (Rivis & Sheeran, 2003; Rhodes et al., 2006).

In the TPB, a fundamental and immediate antecedent factor of a behavior is intention that consists of both motivation and planning components (Vallance et al., 2011). Intention reflects one's motivation and willingness for behavior involving (Ajzen, 1991; Hagger & Chatzisarantis, 2005) that is the main and direct determinant of a behavior (Rhodes et al., 2006; Vallance et al., 2011; Kosma, 2012). Furthermore, intentions are explained by three independent concepts, including attitude toward the behavior, subjective norm (perceived social pressure regarding to the particular behavior), and perceived behavioral control (perceived capability for conducting the behavior) (White et al., 2012). Together, these constructs are representative of a person's total motivation for performing the behavior (Kwan & Bryan, 2010). The TPB actually provides a theoretical framework to describe these three elements of intention (Ajzen & Madden, 1986; Dean et al., 2007; Gretebeck et al., 2007). These constructs are effective on the individual's motivation to take part in the behavior (Kwan & Bryan, 2010). Briefly, with a more positive attitude and subjective norm, and better perceived behavioral control, intention to engage the behavior will be more powerful (Lin et al., 2012). Fig.1

Attitude

Attitude is a personal belief (behavioral beliefs) that weighted by the positive or negative evaluations of a behavior consequence (Ajzen, 1991; Conn et al., 2003; Glanz et al., 2008). Perceived outcomes of the behavior and assessment of these outcomes are crucial in the personal attitude (Lin et al., 2012). Attitude has both instrumental (e.g. harmful or beneficial) and affective (e.g. unpleasant or pleasant) components (Plotnikoff et al., 2010; Vallance et al., 2011). Major reported behavioral beliefs for PA among older adults have been weight control, reduced joint rigidity, social relations, and improved quality of life (Antikainen et al., 2010).

Subjective Norm

Subjective norm is defined as the perceived social force that affects the decision to perform the behavior or not (Nigg et al., 2009). It actually reflects a person's perceptions of social approval for performing the behavior (Cooke & French, 2008). It includes both injunctive (subjective) and descriptive components. Injunctive component is the others' view on a behavior performance. However, descriptive component represents the others' behavior that effects on the individual's performance (Vallance et al., 2011; Kosma, 2012). Family members, friends, physicians, and co-workers usually are significant people who their suggestions are important for performing a behavior in the elderly. For instance, if a woman believes that her spouse wants her to exercise, her subjective norm for the exercise will be increased (Ajzen, 1991; Conn et al., 2003; Antikainen et al., 2010).

Perceived Behavioral Control (PBC)

The PBC was included in the theory of reasoned action for description of behaviors that are not completely volitional (Ajzen, 1991). For these behaviors, intention is not a sufficient factor for action. Translating intentions into action are associated with many difficulties, such as distractions, forgetting or

conflicting bad habits that reduce the magnitude of the relationship between intention and behavior (Schwarzer, 2008; Luszczynska et al., 2011).

The PBC construct is similar to the concept of self-efficacy and reflects the degree of individual's control over internal and external obstacles for behavioral performance based on the perceived simplicity or difficulty (Cooke & French, 2008; Vallance et al., 2011; Linke et al., 2013) in earlier experiences. It is also determined by perceived power regarding to the perceived importance of each of factors that facilitate or hinder the action (Ajzen, 1991; Conn et al., 2003; Dean et al., 2007; Lin et al., 2012).

Control beliefs such as availability of resources and opportunities are the main determinants of people's intentions and actions. It may be supported by experience with the behavior, but acquaintances, friends, and others could influence on the effort of performing the behavior. In fact, more resources and opportunities and fewer barriers are determinants of the greater perceived control over the behavior (Ajzen, 1991). In older adults, usual reported control beliefs have been health problems, care for others, pain, cost, and insufficiency of energy, time, or motivation (Courneya, 1995; Eyler et al., 1998; Conn et al., 2003).

As depicted in Figure 1, perceived control is linked to both intentions and behavior. This proposes that the PBC has a motivational effect on intentions, for instance, individuals may desire to perform the behavior, but facing to some insurmountable barriers at the time reduces their intention or vice versa (Andrew Smith & Biddle, 1999).

Fig1: Components of the theory of planned behavior (Westaby, 2005)

The TPB and PA/Exercise in Older Adults

The TPB is a proper theory for application in the study of PA/exercise, as PA is a behavior that potentially has many obstacles and only to some extent is under personal control (Andrew Smith & Biddle, 1999). The TPB therefore has been widely used for predicting and explaining PA/exercise motivation and behavior in the general population (Hausenblas et al., 1997; Biddle & Nigg, 2000; Hagger et al., 2002a; Rhodes et al., 2006; Vallance et al., 2011). Specifically, a handful of studies has applied the TPB for exercise behavior in older adults (Courneya, 1995; Michels & Kugler, 1998; Conn et al., 2003; Benjamin et al., 2005; Lucidi et al., 2006; Dean et al., 2007; Gretebeck et al., 2007; Yardley et al., 2007; Kosma, 2012).

PA/exercise intention is reduced with advancing age (Rhodes et al., 1999; Yardley et al., 2007). Likewise, Yardley et al. (2007) observed a negative relationship between age and exercise intention in older adults. Because, older individuals believe that they are not capable of performing exercises as compared to younger individuals. However, older people are more inclined to translate their intentions to PA/exercise behavior (Hagger et al., 2002b). Downs and Hausenblas (2005) also found a larger intention-behavior contribution for older adults aged 65 to 80 compared to adults aged 26 to 64, and children aged 8 to 17.

Many studies have reported that the TPB constructs account for 39-44% of the variance of exercise behavior in the general population (Hausenblas et al., 1997; Armitage & Conner, 2001; Hagger et al., 2002a; Rhodes et al., 2006; Vallance et al., 2011). However, data from several sources have revealed that the TPB constructs are more applicable in explaining older adults' intention to exercise. For instance, Lucidi et al. (2006); Lin et al. (2012) and Gretebeck et al. (2007) reported that 45%, 55% and 72% of the variance in PA intention in the elderly was explained by the TPB variables, respectively.

Previous studies have emphasized that demographic characteristics may result in different associations of the TPB constructs to the explanation of intention and behavior (Nigg et al., 2009). Generally, effect of each of the TPB constructs will change based on the behavior and the group of population under study (Grodesky et al., 2006; Dean et al., 2007). In younger adults, intention is the strongest predictor of PA, mostly by attitude and PBC with moderate to large effect sizes, and to a less extent by subjective norm, with a small to negligible effect (Hausenblas et al., 1997; Downs & Hausenblas, 2005; Rhodes et al., 2006). In this regards, Hagger et al. (2002b), Hausenblas et al. (1997), and Downs and Hausenblas (2005) in their meta-analysis review reported that subjective norm is considered a weaker predictor of PA/exercise intentions in general population than attitude and PBC. However, subjective norm and PBC have been found as the strongest determinants of exercise behavior intention in older adults (Wankel & Mummery, 1993; Dean et al., 2007). Dean et al. (2007) also

observed that attitude was not a significant effective factor of strength-training intention among older adults. Although, the majority of aged people have positive attitudes towards exercise, and know its beneficial effect for their physic and mind but yet they still do not participate (Costello et al., 2011). The more influence of subjective norm in older age could be associated with the different motivations factors in physical activity attendance in older versus younger adults. For instance, social interaction is the main reason for joining an exercise program in older adults compared to their younger counterparts (Gill & Overdorf, 1994; Allender et al., 2006; Capalb et al., 2012). Physical activity gives older adults an opportunity for connecting with others, so, they may more interest to join in activities in which their important people are also involved.

These findings suggest that awareness of benefits of physically active lifestyle may not necessarily increase older adult's intention to attend an exercise program. So, health promotion programs, which attempt to change older adults' attitudes towards PA/exercise, might not be enough to facilitate participation in the activity. As a result, interventions would be better target to other TPB variables, such as subjective norm and PBC, for effective change. Hence, old people need a highly supportive settings for maintaining exercise behavior (Kosma, 2012). Additionally, it is important to use appropriate effort and type of exercise considering different elders' age groups and levels of physical and mental health conditions. This might help older adults to obtain an increased sense of PBC.

There are some contradictory ideas on the extension of the TPB theory by applying volitional variables such as planning. In this regards, Norman and Conner (2005) illustrated that planning can play the mediator role by facilitating the translation of intention to exercise behavior among undergraduate students. White et al. (2010) also documented that the TPB-centered intervention on exercise behavior is mediated by planning strategies among old people with diabetes and cardiovascular disease. However, Conn et al. (2003) reported that a planned intention cannot always predict the actual performance and might be modified by an unplanned program. For example, a planned intention to swim after work may be changed due to an unexpected meeting. Additionally, Linke et al. (2013) also stated that planning was not affective on habitual behaviors. In this case, even the strongest intentions are not powerful enough to change the everyday habits.

Conclusion

This paper has given an account of using the TPB theory to enhance PA/exercise behavior in the older adults. In general, the TPB provides a better understanding of the determinant factors of exercise. Previous studies showed that the TPB constructs define more variance of exercise intention in older compared to younger' adults. Subjective norm and PBC have been as the strongest determinants of exercise intention and behavior in the older population. Hence, improvement of attitudes toward a physical active lifestyle might not be sufficient to increase the exercise program involvement. Other TPB variables, such as subjective norm and PBC are required to make PA/exercise behavior change among old population.

Conflicts of Interest

There are no any conflicts of interest to declare.

References

- Ahmad WIW, Ismail Z, & Rahman AHA. (2011). Aging and economic participation of older population in rural malaysia. *International Journal of Humanities and Social Science*. 1(9): 128-133.
- Ajzen I. (1991). The theory of planned behavior. *Organ Behav Hum Dec*. 50(2): 179-211.
- Ajzen I, & Madden T. (1986). Prediction of goal-directed behavior: Attitudes, intentions, and perceived behavioral control. *Journal of experimental social psychology*. 22(5): 453-474.
- Akbartabartoori M, Hankey C, & Lean M. (2005). Healthy lifestyle behaviours are associated with lower probability of having cardiovascular disease risk factors. *Iranian Journal of Public Health*. 34(Sup):75-76.
- Allender S, Cowburn G, & Foster C. (2006). Understanding participation in sport and physical activity among children and adults: A review of qualitative studies. *Health education research*. 21(6): 826-835.
- Andrew Smith R, & Biddle SJ. (1999). Attitudes and exercise adherence: Test of the theories of reasoned action and planned behaviour. *Journal of Sports Sciences*. 17(4): 269-281.
- Antikainen I, Ellis R, Kosma M, Allen PD, Cherry KE, Monroe PA, & Wood RH. (2010). Examining change in theory-based physical activity beliefs of culturally diverse older adults. *J Appl Gerontol*. 29(4): 507-517.
- Armitage C, & Conner M. (2001). Efficacy of the theory of planned behaviour: A meta-analytic review. *Br J Sociol* 40(4): 471-499.
- Benjamin K, Edwards N, & Bharti V. (2005). Attitudinal, perceptual, and normative beliefs influencing the exercise decisions of community-dwelling physically frail seniors. *J Aging Phys Act*. 13(3): 276.
- Biddle S, & Nigg C. (2000). Theories of exercise behavior. *Int J Sport Psychol*.
- Breen L, Stewart CE, & Onambélé GL. (2007). Functional benefits of combined resistance training with nutritional interventions in older adults: A review. *Geriatrics Gerontology International*. 7(4): 326-340.

- Bull FC, & Bauman AE. (2011). Physical inactivity: The “cinderella” risk factor for noncommunicable disease prevention. *J Health Commun.* 16(sup2): 13-26.
- Capalb DJ, O'Halloran P, & Liamputtong P. (2012). Why older people engage in physical activity: An exploratory study of participants in a community-based walking program. *Australian journal of primary health.*
- Conn VS, Tripp R, Toni, & Maas ML. (2003). Older women and exercise: Theory of planned behavior beliefs. *Public Health Nurs.* 20(2): 153-163.
- Conner M, & Sparks P. (2005). Theory of planned behaviour and health behaviour. *Predicting health behaviour.* 2: 170-222.
- Cooke R, & French DP. (2008). How well do the theory of reasoned action and theory of planned behaviour predict intentions and attendance at screening programmes? A meta-analysis. *Psychol Health.* 23(7): 745-765.
- Costello E, Kafchinski M, Vrazel J, & Sullivan P. (2011). Motivators, barriers, and beliefs regarding physical activity in an older adult population. *Journal of geriatric physical therapy.* 34(3): 138-147.
- Courneya KS. (1995). Understanding readiness for regular physical activity in older individuals: An application of the theory of planned behavior. *Health Psychol.* 14(1): 80.
- Dean R, Farrell J, Kelley M, Taylor M, & Rhodes R. (2007). Testing the efficacy of the theory of planned behavior to explain strength training in older adults. *J Aging Phys Act.* 15(1): 1.
- Downs DS, & Hausenblas HA. (2005). Elicitation studies and the theory of planned behavior: A systematic review of exercise beliefs. *Psychol Sport Exerc.* 6(1): 1-31.
- Dumith SC, Hallal PC, Reis RS, & Kohl HW. (2011). Worldwide prevalence of physical inactivity and its association with human development index in 76 countries. *Prev Med.* 53(1): 24-28.
- Elavsky S, McAuley E, Motl RW, Marquez DX, Hu L, Jerome GJ, & Diener E. (2005). Physical activity enhances long-term quality of life in older adults: Efficacy, esteem, and affective influences. *Annals Behavior Medicine.* 30(2): 138-145.
- Eyler AA, Baker E, Cromer L, King AC, Brownson RC, & Donatelle RJ. (1998). Physical activity and minority women: A qualitative study. *Health Educ Behav.* 25(5): 640-652.
- Fishbein M, & Ajzen I. (1975). *Belief, attitude, intention and behavior: An introduction to theory and research.*
- Gill K, & Overdorf V. (1994). Incentives for exercise in younger and older women. *Journal of Sport Behavior.* 17(2): 87-97.
- Glanz K, Rimer BK, & Viswanath K. (2008). *Health behavior and health education: Theory, research, and practice:* John Wiley & Sons.
- Gretebeck KA, Black DR, Blue CL, Glickman LT, Huston SA, & Gretebeck RJ. (2007). Physical activity and function in older adults: Theory of planned behavior. *Am J Health Behav.* 31(2): 203-214.
- Grodesky JM, Kosma M, & Solmon MA. (2006). Understanding older adults' physical activity behavior: A multi-theoretical approach. *Quest.* 58(3): 310-329.
- Guedes DP, Hatmann AC, Martini FAN, Borges MB, & Bernardelli R. (2012). Quality of life and physical activity in a sample of brazilian older adults. *J. Aging Health.* 24(2): 212-226.
- Hagger MS, & Chatzisarantis NL. (2005). First and higher order models of attitudes, normative influence, and perceived behavioural control in the theory of planned behaviour. *Br J Health Psychol.* 44(4): 513-535.
- Hagger MS, Chatzisarantis NL, & Biddle SJ. (2002a). The influence of autonomous and controlling motives on physical activity intentions within the theory of planned behaviour. *Br J Health Psychol.* 7(3): 283-297.
- Hagger MS, Chatzisarantis NL, & Biddle SJ. (2002b). A meta-analytic review of the theories of reasoned action and planned behavior in physical activity: Predictive validity and the contribution of additional variables. *J Sport Exercise Psy.*
- Hausenblas HA, Carron AV, & Mack DE. (1997). Application of the theories of reasoned action and planned behavior to exercise behavior: A meta-analysis. *J Sport Exercise Psy.*
- Heesch KC, Byles JE, & Brown WJ. (2008). Prospective association between physical activity and falls in community-dwelling older women. *Journal of Epidemiology and Community Health.* 62(5): 421-426.
- Higgins A, & Conner M. (2003). Understanding adolescent smoking: The role of the theory of planned behaviour and implementation intentions. *Psychology, Health & Medicine.* 8(2): 173-186.
- Joseph RP, Durant NH, Benitez TJ, & Pekmezi DW. (2013). Internet-based physical activity interventions. *American Journal of Lifestyle Medicine:* 1559827613498059.
- Kosma M. (2012). An expanded framework to determine physical activity and falls risks among diverse older adults. *Res Aging.*
- Kruk J. (2007). Physical activity in the prevention of the most frequent chronic diseases: An analysis of the recent evidence. *Asian Pacific Journal of Cancer Prevention.* 8(3): 325.
- Kwan BM, & Bryan AD. (2010). Affective response to exercise as a component of exercise motivation: Attitudes, norms, self-efficacy, and temporal stability of intentions. *Psychology of Sport and Exercise.* 11(1): 71-79.

- Landi F, Abbatecola AM, Provinciali M, Corsonello A, Bustacchini S, Manigrasso L, Cherubini A, Bernabei R, & Lattanzio F. (2010). Moving against frailty: Does physical activity matter? *Biogerontology*. **11**(5): 537-545.
- Lautenschlager NT, Cox KL, Flicker L, Foster JK, van Bockxmeer FM, Xiao J, Greenop KR, & Almeida OP. (2008). Effect of physical activity on cognitive function in older adults at risk for alzheimer disease. *Journal of American Medical Association*. 300(9): 1027-1037.
- Lin M-H, Chou M-Y, Liang C-K, Peng L-N, & Chen L-K. (2010). Population aging and its impacts: Strategies of the health-care system in taipei. *Ageing Research Reviews*. 9: S23-S27.
- Lin SF, Lee JW, Modeste N, & Johnson EG. (2012). Attitudes and beliefs predicting taiwanese older adults' intentions to attend strength and balance training programs. *J Appl Gerontol*. **31**(2): 260-281.
- Linke SE, Robinson CJ, & Pekmezi D. (2013). Applying psychological theories to promote healthy lifestyles. *American Journal of Lifestyle Medicine*.
- Lucidi F, Grano C, Barbaranelli C, & Violani C. (2006). Social-cognitive determinants of physical activity attendance in older adults. *J Aging Phys Act*. 14(3): 344.
- Luszczynska A, Schwarzer R, Lippke S, & Mazurkiewicz M. (2011). Self-efficacy as a moderator of the planning-behaviour relationship in interventions designed to promote physical activity. *Psychol Health*. 26(2): 151-166.
- Maddux JE, & DuCharme KA. (1997). *Handbook of health behavior research* New York: Plenum Press.
- Michels TC, & Kugler JP. (1998). Predicting exercise in older americans: Using the theory of planned behavior. *Mil Med*. 163(8): 524.
- Mujahid G. (2006). Population age ing in east and south-east asia, 1950-2050: Implications for elderly care. *Asia-Pacific Population Journal*. 21(2): 25-44.
- Nakasato YR, & Carnes BA. (2006). Health promotion in older adults: Promoting successful aging in primary care settings. *Geriatrics*. 61(4): 27-31.
- Nejad LM, Wertheim EH, & Greenwood K. (2005). Comparison of the health belief model and the theory of planned behavior in the prediction of dieting and fasting behavior. *Relaunching soon*. **1**(1): pp. 63-74.
- Netz Y, Wu M, Becker BJ, & Tenenbaum G. (2005). Physical activity and psychological well-being in advanced age: A meta-analysis of intervention studies. *Psychological Aging*. 20(2): 272-284.
- Ní Mhaoláin AM, Gallagher D, O Connell H, Chin A, Bruce I, Hamilton F, Teehee E, Coen R, Coakley D, & Cunningham C. (2012). Subjective well-being amongst community-dwelling elders: What determines satisfaction with life? Findings from the dublin healthy aging study. *Int Psycho Geriatr*. 24(02): 316-323.
- Nigg CR, Lippke S, & Maddock JE. (2009). Factorial invariance of the theory of planned behavior applied to physical activity across gender, age, and ethnic groups. *Psychol Sport Exerc*. 10(2): 219-225.
- Norman P, & Conner M. (2005). The theory of planned behavior and exercise: Evidence for the mediating and moderating roles of planning on intention-behavior relationships. *J Sport Exercise Psy*. 27(4).
- Pernambuco C, Rodrigues B, Bezerra J, Carrielo A, Fernandes A, Vale R, & Dantas E. (2012). Quality of life, elderly and physical activity. *Health*. 4(2): 88-93.
- Plotnikoff RC, Lippke S, Courneya K, Birkett N, & Sigal R. (2010). Physical activity and diabetes: An application of the theory of planned behaviour to explain physical activity for type 1 and type 2 diabetes in an adult population sample. *Psychol Health*. 25(1): 7-23.
- Plotnikoff RC, Lippke S, Courneya KS, Birkett N, & Sigal RJ. (2008). Physical activity and social cognitive theory: A test in a population sample of adults with type 1 or type 2 diabetes. *Applied Psychology*. 57(4): 628-643.
- Rahimi A, Anoosheh M, Ahmadi F, & Foroughan M. (2011). Exercise and physical activity among healthy elderly iranians. *SE Asian J Trop Med*. 42(2): 444.
- Rhodes RE, Blanchard CM, & Matheson DH. (2006). A multicomponent model of the theory of planned behaviour. *Br. J. Health Psychol*. **11**(1): 119-137.
- Rhodes RE, Martin AD, Taunton JE, Rhodes EC, Donnelly M, & Elliot J. (1999). Factors associated with exercise adherence among older adults. *Sports Medecine*. 28(6): 397-411.
- Rivis A, & Sheeran P. (2003). Descriptive norms as an additional predictor in the theory of planned behaviour: A meta-analysis. *Curr Psychol*. 22(3): 218-233.
- Schulz E, Leidl R, & König H-H. (2004). The impact of ageing on hospital care and long-term care—the example of germany. *Health Policy*. 67(1): 57-74.
- Schutzer KA, & Graves BS. (2004). Barriers and motivations to exercise in older adults. *Preventive Medicine*. 39(5): 1056-1061.
- Schwarzer R. (2008). Modeling health behavior change: How to predict and modify the adoption and maintenance of health behaviors. *Applied Psychology*. 57(1): 1-29.

- Singh GK, & Hiatt RA. (2006). Trends and disparities in socioeconomic and behavioural characteristics, life expectancy, and cause-specific mortality of native-born and foreign-born populations in the united states, 1979–2003. *International Journal of Epidemiology*. 35(4): 903-919.
- Sofi F, Capalbo A, Cesari F, Abbate R, & Gensini GF. (2008). Physical activity during leisure time and primary prevention of coronary heart disease: An updated meta-analysis of cohort studies. *European Journal of Cardiovascular Prevention & Rehabilitation*. 15(3): 247-257.
- Stadler G, Oettingen G, & Gollwitzer PM. (2009). Physical activity in women: Effects of a self-regulation intervention. *Am J Prev Med*. 36(1): 29-34.
- Vallance JK, Murray TC, Johnson ST, & Elavsky S. (2011). Understanding physical activity intentions and behavior in postmenopausal women: An application of the theory of planned behavior. *International Journal of Behavioral Medicine*. 18(2): 139-149.
- Wallin M, Talvitie U, Cattani M, & Karppi S-L. (2008). Construction of group exercise sessions in geriatric inpatient rehabilitation. *Health Commun*. 23(3): 245-252.
- Wankel LM, & Mummery WK. (1993). Using national survey data incorporating the theory of planned behavior: Implications for social marketing strategies in physical activity. *J Appl Sport Psychol*. 5(2): 158-177.
- Westaby JD. (2005). Behavioral reasoning theory: Identifying new linkages underlying intentions and behavior. *Organ Behav Hum Dec*. 98(2): 97-120.
- White KM, Terry DJ, Troup C, Rempel LA, & Norman P. (2010). Predicting the consumption of foods low in saturated fats among people diagnosed with type 2 diabetes and cardiovascular disease. The role of planning in the theory of planned behaviour. *Appetite*. 55(2): 348-354.
- White KM, Terry DJ, Troup C, Rempel LA, Norman P, Mummery K, Riley M, Posner N, & Kenardy J. (2012). An extended theory of planned behavior intervention for older adults with type 2 diabetes and cardiovascular disease. *Journal of aging and physical activity*. 20(3): 281-299.
- Yardley L, Donovan-Hall M, Francis K, & Todd C (Cartographer). (2007). Attitudes and beliefs that predict older people's intention to undertake strength and balance training.