Original Article

Water balance changes associated with performance level in the Half Ironman

KOVÁŘOVÁ LENKA¹, HADŽEĞA TOMÁŠ²

^{1,2} Faculty of Physical Education and Sport, Charles University, Prague, CZECH REPUBLIC

Published online: September 28, 2015

(Accepted for publication september 09, 2015)

DOI:10.7752/jpes.2015.03070;

Abstract:

The objective of our study was to analyse the total body water (TBW), intracellular water (ICW) and extracellular water (ECW) dynamics for two groups of athletes (elite and hobby) during one race (the Half Ironman). Forty-two male triathletes volunteered to participate in the study. Their body composition parameters were determined with the Tanita MC-980 multi-frequency segmental body composition analyser. The results are presented as means and standard deviations. The pre- and post-race results for both groups were analysed with paired t-tests. The pre-race results for both groups were compared using non-paired t-tests. The associations between body composition variables and race times were investigated with Pearson's correlation analyses. The statistical significance was set at $P \leq 0.05$.

Professional athletes achieved higher hydration rates before performance in terms of both the TBW percentage ($P = 0.005$) and the ICW percentage ($P = 0.006$) parameters. However, there were no significant differences between the groups of athletes in water-control ability during endurance exercise. In both groups of athletes, significant changes in water balance and total weight loss were observed during the race towards Physical Activity.

Key words: triathlon, body water, hydration, performance.

Introduction

A triathlon is characterised by participation in three sports (swimming, cycling, and running) one after the other (Neumann, Pfützner, & Hottnerott, 2004). Several variations have occurred during the 30 year history of triathlons. The best-known variation, which is an Olympic Games event, is the so-called short (Olympic) triathlon with a 1.5-km swim, a 40-km bike ride and a 10-km run. The long triathlon does not have fixed trail distances. The most common race distances are a 3.6-km swim, a 180-km bike ride and a 42.2-km run (more commonly known as the Ironman); a trail of half these lengths is referred to as the Half Ironman. Because of the duration of the Half Ironman race (4–6 h, depending on the performance level), participants rely on aerobic energy release (Friel, 2009).

Total body water (TBW) is a significant component of body weight. The amount of water in the body depends on age, sex and body weight. The average TBW amount is 63 % in an adult male and 53 % in an adult female (Rokyta & Štátný, 2002). Depending on the water's localisation, TBW can be classified as intracellular (ICW) and extracellular (ECW). ICW accounts for ca 40 % of body weight. ECW accounts for approximately 20 % of total body weight and is divided into intravascular fluid (plasma) and interstitial fluid (lymph). Although TBW information is needed to estimate body composition and determine nutritional status, its sub-division into ICW and ECW is more important for research purposes than TBW alone (Moore, 1963; Segal, 1991).

Dehydration is a common finding in endurance athletes; it is defined as the loss of more than 2 % of body weight during an endurance performance (Sawka et al., 2007; Sawka & Montain, 2000; Sawka & Noakes, 2007). Many studies argue that dehydration leads to a poorer endurance performance (Stearns et al., 2009; von Duvillard, Braun, Markofski, Beneke, & Leithäuser, 2004). One study (Kovacs, 2011) mentions that scientists who have investigated the effect of liquid intake limitations on the volumes and functions of the human body note that hyperthermia (elevated body temperature caused by dehydration) directly affects brain function by changing the blood flow and metabolism in the brain. This reduces the performance of the central cognitive or neuromuscular unit, which may impair muscular function, change the perception of effort, or both. Dehydration may result in clinical symptoms similar to those of a concussion: fatigue, drowsiness, headaches, inability to concentrate and a loss of balance.

However, the conclusion that a 2 % loss of body weight adversely affects performance is derived only from laboratory results based solely on studies with relatively few participants (Noakes, 2007). In contrast to these studies, there is evidence that the fastest contestants in athletic competitions usually lose the largest amount of body weight during competition. Previous studies (Pugh, Corbett, & Johnson, 1967; Wyndham & Strydom,

1969) have reported that the athletes who had lost the greatest amounts of body weight usually had the highest body temperatures after the race and had won the competitions. Other studies (Byrne, Lee, Chew, Lim, & Tan, 2006; Chevront & Haymes, 2001; Zouhal et al., 2009) reveal that a loss of body weight greater than 3–4 % is common among faster-finishing athletes, and there was a weak inverse relationship between body weight loss and sports performance. Thus, the fastest-finishing athletes tend to lose the greatest amounts of body weight. Fudge and Pitsiladis (2009) note that the world record holder for the 42 km in 2008 lost 10 % of his body weight in achieving this record. If it were true that the loss of 10 % of body weight worsened sports performance by more than 50–100 %, it would have been impossible for that athlete to achieve this record performance (Craig & Cummings, 1966; Sawka, Francesconi, Young, & Pandolf, 1984).

Sharwood, Collins, Goedecke, Wilson, & Noakes (2002) also fail to find any evidence that body weight loss or dehydration were related to poor performances in a survey of 297 triathletes participating in the Ironman race. The athletes who lost the greatest amount of body weight completed the race in the shortest time. This result was confirmed by a survey conducted in 1967. Pugh et al. (1967) report that the winner (2 h, 38 min) of the national marathon in England lost 5.23 kg, which was 6.9 % of his body weight. Furthermore, the first four athletes who completed the race had an average body weight loss of 5.8 %.

Recent studies of ultra-marathon runners (23 ultra-marathon runners participating in a 24-h run) reveal that body weight loss is positively correlated with competitive performance (Kao, Shyu, & Yang, 2008). However, no change in body weight was registered for swimmers during ultra-long distance swims (Knechtle, Knechtle, Kaul, & Kohler, 2009a). It is possible that the body weight loss in ultra-performance athletes occurs not only as a result of dehydration but as a result of skeletal muscle and fat mass loss (Knechtle et al., 2009a; Knechtle et al., 2010; Knechtle et al., 2011; Knechtle, Wirth, Knechtle, Rosemann, & Senn, 2011; Zouhal et al., 2009). Another study (Tam, Nolte, & Noakes, 2011) devoted to half-marathon and ultra-marathon runners reveals a body weight change in participants in both types of races that exceeded the changes in their TBW results. This finding means that TBW loss alone does not account for the body weight loss during the race.

Knechtle, Wirth, Knechtle, and Rosemann (2009b) investigated whether ultra-performance runners participating in a 100-km race suffered from body weight loss and whether this loss involved fat mass, muscle mass or TBW. The ultra-runners were evaluated before and after racing. The results were a 1.6-kg ($P < 0.01$) loss in body weight, a 0.4-kg ($P < 0.01$) loss in fat mass and a 0.7-kg loss in muscle mass ($P < 0.01$). The TBW increased by 0.8 l ($P < 0.05$). Jurimae and Jurimae (2001) report a total weight decrease from 84.7 $\pm$ 9.1 to 83.6 $\pm$ 9.6 kg and a TBW decrease from 46.5,1 to 46.0 $\pm$ 3.01 l among their study participants after a 30-min maximum workout on a rowing machine. In contrast, after an Alpine marathon in Spain, 24 amateur athletes (aged 39 $\pm$ 10 years) showed a significant increase ($P < 0.05$) in ICW (27.4 vs. 28.1 l) and TBW (43.6 vs. 45.1 l) and a decrease in body weight (71.8 vs. 68.2 kg) (Clemente, Ramos, & Gonzalez, 2011).

An interesting issue is the correlation between body composition dynamics during the race and the performance achieved. Knechtle, Knechtle, Wirth, Rüst, and Rosemann (2012) report changes in body weight and TBW in 219 amateur male runners during a 100-km ultra-marathon. The athletes lost 1.9 kg of their body weight ($s = 1.4$), which equals 2.5 % ($s = 1.8$) of body weight ($P < 0.001$). The athletes also lost 0.2 kg ($s = 1.3$) of their presumed fat mass ($P < 0.05$) and 0.9 l ($s = 1.6$) of their presumed TBW ($P < 0.001$). The study concluded that faster runners lost more body weight; the runners also lost more body weight if they drank less liquid. Faster runners also drank more fluids compared with slower runners. Yet another study by Sharwood, Collins, Goedecke, Wilson, and Noakes (2004) found a significant correlation between the rate of body weight loss and completion time in the Ironman race. The athletes who completed the race with the highest body weight losses obtained the fastest times in the final results. Zouhal et al. (2011) note the significant linear relationship between the rate of body weight loss and the completion time for marathon runners. The athletes with the highest body weight losses obtained the best times. In their study of the association between changes in body weight and the total 42-km finish time of marathon runners Zouhal et al. note massive differences in body water among the runners. The body water changes ranged from an 8 % loss to a 5 % gain. The researchers also state that 55 % of the runners who lost more than 2 % of their body weight during the race ran significantly faster compared with runners who lost less body weight. This finding contradicts the principal component of the ACSM Position Stand knowledge (Sawka et al., 2007) but agrees with other studies claiming that the fastest athletes in marathon races and other endurance competitions tend to lose the greatest amounts of body weight (Chevront, Carter, & Sawka, 2003; Kao et al., 2008; Zouhal et al., 2009).

The objective of our study was to analyse the TBW, ICW and ECW dynamics in two groups of athletes (elite and hobby) during one race (the Half Ironman). We compared these parameters prior to and after the race to identify potential changes that may be associated with physiological predispositions for endurance performances.

Method

Participants

Forty-two male triathletes volunteered to participate in the study.

Procedures

According to their performance level and finish time, the athletes were divided into an elite (G-1) and a hobby (G-2) group. The runners' anthropometric characteristics are provided in Table 1. The athletes were evaluated before (M-1) and 10 min after (M-2) the "Czechman Half Ironman" competition (1.8 km swimming, 90 km cycling and 21 km running).

The runners signed an informed consent waiver and were submitted to the experimental protocol according to the study design. The procedures and techniques proposed in this study were approved by the ethics committee FPES CU.

Table 1. Basic anthropometric characteristics of the athletes

	G-1	G-2	<i>P</i>
N	16	26	
Age (years)	32.75 ± 6.87	36.15 ± 7.39	0.145
Race time (h:mm:ss)	4:42:38 ± 24:00	5:26:59 ± 13:50	0.000
Body weight (kg)	76.71 ± 6.83	80.18 ± 7.51	0.140
Body height (cm)	180.06 ± 5.94	179.77 ± 4.82	0.862

Instruments

The body composition parameters were collected using a Tanita MC-980 multi-frequency segmental body composition analyser.

Statistical analysis

The results are presented as means and standard deviations. The pre- and post-race results of both groups were compared using paired t-tests. The pre-race results of both groups were compared using non-paired t-tests.

Associations between body composition variables and race times were investigated using Pearson's correlation analyses. Significance was set at $P \leq 0.05$.

Results

The results for each variable at M-1 and the p-values for the statistical comparisons for G-1 and G-2 are presented in table 2.

For the pre-race measurement (M-1), no differences were found in body mass (BM) ($P = 0.14$), TBW ($P = 0.966$), ECW ($P = 0.313$) or ICW ($P = 0.734$). However, when taking into account the other parameters (total weight) and assessing the parameters as a percentage of total weight, significant differences emerged for TBW % ($P = 0.005$) and ICW % ($P = 0.006$). ECW % ($P = 0.072$) did not reach significance.

Table 2. Water balance before the race and the p-values for the statistical comparisons of G-1 and G-2
M1 before the race

	G-1	G-2	<i>P</i>
BM (kg)	76.71 ± 6.83	80.18 ± 7.51	0.140
TBW (L)	48.81 ± 3.67	48.76 ± 3.83	0.966
ECW (L)	17.02 ± 0.83	17.29 ± 0.85	0.313
ICW (L)	31.79 ± 2.88	31.47 ± 3.05	0.734
TBW (% BM)	63.73 ± 2.41	60.97 ± 3.23	0.005
ECW (% BM)	22.27 ± 1.01	21.66 ± 1.06	0.072
ICW (% BM)	41.48 ± 1.84	39.31 ± 2.58	0.006

The results for each variable (G-1, G-2), the differences between M-1 and M-2, and the p-values for statistical comparisons are presented in Table 3.

Table 3. Water balance before and after the race and the p-values for the statistical comparisons independent of G-1 and G-2

G-1				G-2		
	M1	M2	<i>P</i>	M1	M1	<i>P</i>
BM (kg)	76.71 ± 6.83	75.89 ± 6.45	0.032	80.18 ± 7.51	79.53 ± 7.48	0.003
TBW (L)	48.81 ± 3.67	47.83 ± 3.85	0.000	48.76 ± 3.83	48.01 ± 4.00	0.002
ECW (L)	17.02 ± 0.83	16.84 ± 0.83	0.000	17.29 ± 0.85	17.13 ± 0.87	0.000
ICW (L)	31.79 ± 2.88	30.99 ± 3.20	0.000	31.47 ± 3.05	30.88 ± 3.20	0.005

In both groups of triathletes, water balance and total weight changed significantly. Total weight decreased in G-1 ($P = 0.032$) and G-2 ($P = 0.003$). The same results were registered for TBW ($P = 0.000$ and $P = 0.002$, respectively), ECW ($P = 0.000$ and $P = 0.000$, respectively) and ICW ($P = 0.000$ and $P = 0.005$, respectively).

We further analysed the water balance changes caused by the race in connection with performance. The results for each variable and the p-values for G-1 and G-2 are presented in Table 4.

Table 4. Water balance changes during the race and the p-values for the statistical comparisons of G-1 and G-2

	M1 - M2 G-1	G-2	<i>P</i>
Δ BM _(M1 - M2) (kg)	0.82 ± 1.38	0.65 ± 1.01	0.658
Δ TBW _(M1 - M2) (L)	0.98 ± 0.73	0.75 ± 1.11	0.465
Δ ECW _(M1 - M2) (L)	0.18 ± 0.16	0.16 ± 1.16	0.695
Δ ICW _(M1 - M2) (L)	0.80 ± 0.62	0.59 ± 0.97	0.442

No significant differences in water balance changes were found for any of the analysed groups. Additionally, the association between water balance changes and race times was insignificant (table 5).

Table 5. Pearson's correlation between water balance changes and race times

	Pearson's correlation	<i>P</i>
Δ BM _(M1 - M2) (kg)	-0.129	0.415
Δ TBW _(M1 - M2) (L)	0.034	0.830
Δ ECW _(M1 - M2) (L)	-0.039	0.805
Δ ICW _(M1 - M2) (L)	0.046	0.771

Discussion

The objective of our study was to analyse water balance changes during the Half Ironman race. To enhance the quality of the study, the analysis was performed for two separate groups (elite [G-1] and hobby [G-2]). In both groups, the water balance parameters decreased as a result of the competitive performance. This finding agrees with the results of some studies (Jurimae & Jurimae, 2001; Knechtle et al., 2012; Tam et al., 2011); however, it contradicts the results of other studies (Clemente et al., 2011; Knechtle et al., 2009b) that report increased TBW among runners after the race. A triathlon is a combination of swimming, cycling and running. The majority of available studies have examined endurance runners. There are very few studies devoted expressly to cycling (Knechtle et al., 2009c) or swimming (Knechtle et al., 2009a). The results available for long-distance swimmers do not report any changes in body weight (Knechtle et al., 2009a).

The results of our study may lead to the assumption that the dynamics of all parameters were independent of the triathletes' performances, which would imply that athletes' ability to control water metabolism during athletic endeavours is not a parameter that can be considered a talent.

The correlation between water balance changes and performance (race time) was not confirmed. This finding agrees with the results of some studies (Byrne et al., 2006; Cheuvront & Haymes, 2001) but contradicts the results of other studies (Knechtle et al., 2012; Zouhal et al., 2011) that present a linear relationship between the rate of body weight loss and race completion time.

Most studies do not present the environmental conditions during the test. Because the majority of cases represent field testing, this information is often essential for data interpretation. In our case, the conditions were approximately 18 degrees Celsius, windy, and partly rainy. Such conditions can dramatically affect the athletes' performance, refreshment system and drinking regimen.

The finding most relevant to the selection of talents is the significant difference in the TBW % ($P = 0.005$) and the ICW % ($P = 0.006$) between groups (G-1 and G-2) prior to the race. The better triathletes were better hydrated before the performance. It is evident from the results that the difference in total body hydration is mostly a consequence of the amount of ICW. ICW accounts for 40 % of body weight, or 66 % of TBW, which amounts to approximately 30–35 % of the TBW in the soft tissues (mainly in muscles). The remainder, corresponding to 8–10 % of TBW, is found within the connective tissues, cartilage and bones (Trojan, 1999). Intracellular fluid contains almost four times more protein compared with plasma and contains a relatively large amount of potassium and phosphate ions (but far less sodium and chlorine ions compared with plasma). Calcium ions, whose levels are kept low, are also present in the cells in small concentrations (Rokyta & Šťastný, 2002).

Very similar water metabolism levels during the race were observed for both groups; therefore, the hydration status before the performance plays the greatest observable role. If all athletes experience a similar loss of fluids while racing, the athletes with better hydration statuses prior to the race finish with values closer to

normal; thus, they are not totally dehydrated. Thus, pre-race hydration may play a key role, especially during the final phase of the race.

Our study did not address whether statistically higher hydration values before endurance racing result from more the efficient water or ICW metabolism necessary for maximum endurance performances or the extent to which this result was influenced by pre-race tune ups and potential super-compensation.

Conclusions

The main conclusions of our study are that (i) elite athletes achieve a higher hydration status in terms of both TBW and ICW before their performances. In contrast, (ii) the differences identified in the water balance parameters during the race were comparable for both groups. Therefore, there were no differences in water control ability during endurance exercise performances between the surveyed groups of athletes. We also found that (iii) significant water balance and total weight loss occurred in both groups of athletes during the workout.

References

- Byrne, C., Lee, J. K. W., Chew, S. A. N., Lim, C. L., & Tan, E. Y. M. (2006). Continuous Thermoregulatory Responses to Mass-Participation Distance Running in Heat. *Medicine & Science in Sports & Exercise*, 38(5), 803–810.
- Clemente, S. V., Ramos, C. D., & Gonzalez, J. M. (2011). Modifications to body composition after running an alpine marathon. *International SportMed Journal*, 12(3), 133–140.
- Craig, E. N., & Cummings, E. G. (1966). Dehydration and muscular work. *Journal of Applied Physiology*, 21(2), 670–674.
- Friel, J. (2009). *The Triathlete's Training Bible*. Boulder, CO: VeloPress.
- Fudge, B. W., & Pitsiladis, Y. P. (2009). Efficacy of prevailing fluid intake recommendations for elite marathon running. *Medicine & Science in Sports & Exercise*, 41, 190.
- Cheuvront, S., Carter, R., & Sawka, M. N. (2003). Fluid balance and endurance exercise performance. *Curr Sports Med Rep*, 2(4), 202–208.
- Cheuvront, S. N., & Haymes, E. M. (2001). Ad libitum fluid intakes and thermoregulatory responses of female distance runners in three environments. *Journal of Sports Sciences*, 19(11), 845–854.
- Jurimae, J., & Jurimae, T. (2001). Body composition analysis by bioelectrical impedance: effect of single training session in male rowers. *Papers on Anthropology*, 10, 67–72.
- Kao, W. F., Shyu, C. L., & Yang, X. W. (2008). Athletic performance and serial weight changes during 12- and 24-hour ultra-marathons. *Clinical Journal of Sport Medicine*, 18, 155–158.
- Knechtle, B., Knechtle, P., Kaul, R., & Kohler, G. (2009a). No Change of Body Mass, Fat Mass, and Skeletal Muscle Mass in Ultraendurance Swimmers After 12 Hours of Swimming. *Research Quarterly for Exercise & Sport*, 80(1), 62–70.
- Knechtle, B., Knechtle, P., Wirth, A., Rüst, A. C., & Rosemann, T. (2012). A faster running speed is associated with a higher body weight loss in 100-km ultra-marathoners. *Journal of Sports Sciences*, 30(11), 1131–1140.
- Knechtle, B., Senn, O., Imoberdorf, R., Joleska, I., Wirth, A., Knechtle, P., & Rosemann, T. (2010). No Fluid Overload in Male Ultra-Runners During a 100 km Ultra-Run. *Research in Sports Medicine*, 19(1), 14–27. doi: 10.1080/15438627.2010.510039
- Knechtle, B., Wirth, A., Knechtle, P., & Rosemann, T. (2009b). Increase of Total Body Water With Decrease of Body Mass While Running 100 km Nonstop--Formation of Edema? *Research Quarterly for Exercise & Sport*, 80(3), 593–603.
- Knechtle, B., Wirth, A., Knechtle, P., & Rosemann, T. (2009c). An Ultra-cycling Race Leads to no Decrease in Skeletal Muscle Mass. *International Journal of Sports Medicine*, 30(3), 163–167.
- Knechtle, B., Wirth, A., Knechtle, P., Rosemann, T., & Senn, O. (2011). Do ultra-runners in a 24-h run really dehydrate? *Irish Journal of Medical Science*, 180(1), 129–134. doi: 10.1007/s11845-010-0500-8
- Kovacs, M. (2011). Hydration, dehydration, and performance: A literature review. *Coaching & Sport Science Review (Spanish Version)*, 19(53), 21–23.
- Moore, F. D., Olesen, K. H., McMurrey, J. D., Parker, H. V., Ball, M. R., Boyden, C. M. (1963). *The body cell mass and its supporting environment in body composition in health and disease*. Philadelphia, PA: WB Saunders.
- Neumann, G., Pfützner, A., & Hottnerott, K. (2004). *Das grosse Buch vom Triathlon*. Aachen: Meyer and Meyer Vellag.
- Noakes, T. D. (2007). Drinking guidelines for exercise: What evidence is there that athletes should drink "as much as tolerable", "to replace the weight lost during exercise" or "ad libitum"? *Journal of Sports Sciences*, 25(7), 781–796.
- Pugh, L. G., Corbett, J. L., & Johnson, R. H. (1967). Rectal temperatures, weight losses, and sweat rates in marathon running. *Journal of Applied Physiology*, 23(3), 347–352.
- Rokyta, R., & Šťastný, F. (2002). *Struktura a funkce lidského těla*. Praha: TIGIS.

-
- Sawka, M. N., Burke, L. M., Eichner, E. R., Maughan, R. J., Montain, S. J., & Stachenfeld, N. S. (2007). American College of Sports Medicine position stand. Exercise and fluid replacement. *Medicine and Science in Sports and Exercise*, 39(2), 377–390.
- Sawka, M. N., Francesconi, R. P., Young, A. J., & Pandolf, K. B. (1984). Influence of hydration level and body fluids on exercise performance in the heat. *JAMA*, 252(9), 1165–1169. doi: 10.1001/jama.1984.03350090041020
- Sawka, M. N., & Montain, S. J. (2000). Fluid and electrolyte supplementation for exercise heat stress. *The American Journal of Clinical Nutrition*, 72(2 Suppl), 564S–572S.
- Sawka, M. N., & Noakes, T. D. (2007). Does dehydration impair exercise performance? *Medicine and Science in Sports and Exercise*, 39(8), 1209–1217.
- Segal, K., Burastero, S., Chun, A., Coronel, P., Pierson, R. N. Jr., & Wang, J. (1991). Estimation of extracellular and total body water by multiplefrequency bioelectrical- impedance measurement. *The American Journal of Clinical Nutrition*, 54(1), 26–29.
- Sharwood, K., Collins, M., Goedecke, J., Wilson, G., & Noakes, T. (2002). Weight changes, sodium levels, and performance in the South African Ironman Triathlon. *Clinical Journal of Sport Medicine*, 12(6), 391–399.
- Sharwood, K., Collins, M., Goedecke, J. H., Wilson, G., & Noakes, T. D. (2004). Weight changes, medical complications, and performance during an Ironman triathlon. *British Journal of Sports Medicine*, 38(6), 718–724.
- Stearns, R. L., Casa, D. J., Lopez, R. M., McDermott, B. P., Ganio, M. S., Decher, N. R., . . . Maresh, C. M. (2009). Influence of hydration status on pacing during trail running in the heat. *Journal of Strength & Conditioning Research (Lippincott Williams & Wilkins)*, 23(9), 2533–2541.
- Tam, N., Nolte, H. W., & Noakes, T. D. (2011). Changes in Total Body Water Content During Running Races of 21.1 km and 56 km in Athletes Drinking Ad libitum. *Clinical Journal of Sport Medicine*, 21(3), 218–255.
- von Duvillard, S. P., Braun, W. A., Markofski, M., Beneke, R., & Leithäuser, R. (2004). Fluids and hydration in prolonged endurance performance. [Article]. *Nutrition*, 20(7/8), 651–656. doi: 10.1016/j.nut.2004.04.011
- Wyndham, C. H., & Strydom, N. B. (1969). Danger of an inadequate water intake during marathon running. *South African Medical Journal*, 43(29), 893–896.
- Zouhal, H., Groussard, C., Minter, G., Vincent, S., Cretual, A., Gratas-Delamarche, A., . . . Noakes, T. D. (2011). Inverse relationship between percentage body weight change and finishing time in 643 forty-two-kilometre marathon runners. *British Journal of Sports Medicine*, 45(14), 1101–1105.
- Zouhal, H., Groussard, C., Vincent, S., Jacob, C., Abderrahman, A. B., Delamarche, P., & Gratas-Delamarche, A. (2009). Athletic performance and weight changes during the "Marathon of Sands" in athletes well-trained in endurance. *International Journal of Sports Medicine*, 30(7), 516–521. doi: 10.1055/s-0029-1202350.

Acknowledgements: This work was supported by GAČR under Grant 13-07776P and Charles University in Prague under Grant number 39 Social-Sciences Aspects of Human Movement Studies.