Endurance performance and stretch-shortening cycle in two male amateur soccer teams – a cross-sectional study

MARIANO RINGGENBERG¹; ROGER SAGER²; SLAVKO ROGAN³

^{1,2,3}Bern University of Applied Sciences, Department of Health, Discipline of Physiotherapy, SWITZERLAND

³Academy for integrative Physiotherapy und Training Education, Grenzach-Wyhlen, GERMANY

Published online: March 31, 2017

(Accepted for publication January 28, 2017)

DOI:10.7752/jpes.2017.01029

Abstract:

This study examined the investigation protocol and compares the endurance performance and the fast stretch-shortening cycle (F-SSC) in two male amateur soccer teams. Altogether, twenty amateur soccer players from two teams (third and fifth amateur division) participated in this cross-sectional study. Outcomes were yo-yo intermittent recovery test level 1 (YYIRT1) for endurance performance and a plyometric jump test (PJT) for the F-SSC. The recruitment rate of 45.5% is low while the attrition rate of 0% and the adherence rate of 100% are good. For endurance performance and F-SSC, no significant differences between the teams were determined. The findings from this study indicate that it is feasible with modification such as a familiarization to the PJT. It shows that the YYIRT1 is a practical tool to monitoring actual training levels for endurance performance in amateur soccer players. This study shows no significant difference between a third division and fifth division amateur soccer team in endurance performance and F-SSC.

Key Words: soccer, endurance testing, jump testing.

Introduction

Many research studies were conducted with professional soccer players or professional soccer players were compared with amateur soccer players. Kaplan et al. (2009) showed that in the shuttle run test, professional soccer players have significantly higher values than amateurs. Ostojic et al. (2004) found significant differences in favour of the professional soccer players regarding VO_{2max} and maximal heart rate (HR_{max}) during 20 m shuttle run tests, as well as during the squat jump (SJ).

Arnason et al. (2004), Cometti et al. (2001), Ostojic (2004), Dellal et al. (2011), Le Gall et al. (2010) and Wisloff et al. (2004) evaluated differences in MVC, slow stretch-shortening cycle (S-SSC) and explosive strength respectively. In contrast, there is a lack of adequate information for reactive strength, which described the fast stretch-shortening cycle (F-SSC). Güllich & Schmidtbleicher (1999) refer to reactive strength as a component of the rate of RFD. Reactive strength in the form of an F-SSC is also important in soccer (Reilly & Ekblom, 2005).

To our knowledge, appropriate studies are rare in amateur soccer players. Science has also to spread its knowledge to amateur sport. There is a lack of investigation protocols that describe which endurance strength components are to be measured and how these data must be collected. For this reason, and due to lack of data, it seems justified in this context to perform at first a study with amateur male soccer players to generate basis data for planning future studies. The aim of this study was 1) to test the feasibility of the investigation protocol and 2) to evaluate the endurance performance and the F-SSC between two amateur male soccer teams.

Material & methods

Participants

Twenty Swiss male amateur soccer players from a third division team ($n=10$; mean age: 24.04 ± 2.92 years) and fifth division team ($n=10$; mean age: 24.16 ± 3.81 years) voluntarily participated in this cross-sectional study. All the soccer players were recruited based on a) they had a minimum of eight years of training and weekend match day experience, and b) they trained in the evening, at least two days per week for an average of 90 minutes each time in addition to the weekend match, and c) were members of the same soccer team. Moreover, any players who experienced injuries, or missed 50% of training sessions were excluded from this study.

Table 1 Characteristics of the amateur soccer players with means and standard deviation of age and anthropometry

	First team (n = 10)	Second team (n = 10)
age (years)	24.04 ± 2.92	24.16 ± 3.81
height (cm)	178.05 ± 4.31	180.25 ± 4.92
weight (kg)	82.85 ± 8.84	80.45 ± 9.66
BMI (kg x m ⁻²)	26.10 ± 2.20	24.69 ± 1.96
abdominal circumference (cm)	88.15 ± 6.58	86.60 ± 5.99
thigh circumference (cm)	55.50 ± 3.41	53.30 ± 3.20

This study is consistent with the institutional ethical requirements for human experimentation in accordance with the latest version of the Declaration of Helsinki. All participants gave written informed consent.

Procedures

Two survey sessions were organized during the in-season soccer-training period on an outdoor training field. The first session was designed to collect the following anthropometric data (Table 1): Body height (tape rule, Stanley, Idstein, Germany), body weight (digital bathroom scale, Soehnle, Leifheit AG, Nassau, Germany).

The warm-up program for both test days was standardized and consisted of three minutes of running the 20 m distance back and forth at a set pace (i.e. 9.0 km/h) with the help of “beep” sounds. The surveys took place in the evening between 7-10 p.m. The same two investigators (MR, RS) performed all the measurements.

Outcome: measurements

To test the endurance performance, the yo-yo intermittent recovery test level 1 (YYIRTLL1) was applied (Krustrup et al., 2003). The test has an intraclass correlation coefficient (ICC) of 0.86 (Thomas, Dawson, & Goodman, 2006). The YYIRTLL1 consisted of 20 m shuttle runs performed at increasing velocities with ten seconds of active recovery between runs until exhaustion.

A Myotest® device (Sion, Switzerland) was used to test the F-SSC with the PJT. The Myotest® device is a small electronic triaxial accelerometer which is fixed on a Velcro belt around the waist. The Myotest® device has very good reliability for F-SSC with an ICC of 0.98 (Casartelli, Muller, & Maffiuletti, 2010). The participants started in an upright standing position with their hands on their hips; after a beep start signal they were then instructed to flex their knees around 90° as quickly as possible and then jump as high as possible in the ensuing concentric phase. When their feet touched the ground on landing they jumped up as high as possible another five times with the minimum possible ground contact time while keeping the knees and ankles extended. After five bounces a double beep signal occurs.

Statistical analyses

The variables are described as mean ± SD, as well as median and interquartile range (IQR). Due to the low number of participants, nonparametric analysis was used. The Mann-Whitney U test was used to analyze group differences regarding endurance performance and F-SSC. A p value < 0.05 was considered as statistically significant. The data were analyzed using the statistical package program IBM SPSS Version 19 (SPSS, Inc., Chicago, USA) and Microsoft Excel 2007 (Microsoft Corporation, Redmond, USA).

Results

After the verbal information session, all 20 amateur soccer players who were present were recruited. These comprised ten players (number of starters: 80%) from the third amateur division team (they ranked 7th) and ten players (number of starters: 60%) from the fifth amateur division team (they ranked 4th). All soccer players carried out the YYIRTLL1 and PJT measurements. The attrition rate was 0% and the adherence rate 100%. There were no reports of adverse effect or injury during the investigation.

Table 2 shows the median (IQR) of the variables of endurance performance while Table 3 represents the variable of the F-SSC index. No significant difference between the two teams was found for endurance performance ($P = 0.65$) and for F-SSC index ($P = 0.67$).

Table 2 Descriptive statistics of the YYIRTLL1 in meter

Team	n	distance median (IQR)
Third Division	10	1000 (440 – 1880)
Fifth Division	10	1080 (440 – 1520)

Mean and ± SD and median and interquartile range (IQR) values for YYIRTLL1. P-values compared between the two teams after YYIRTLL1 performance using Mann-Whitney U test. ± SD = standard deviation; YYIRTLL1 = yo-yo intermittent recovery test level 1

Table 3 Descriptive statistics of the F-SSC index

Team	n	Index Median (IQR)
Third Division	10	1.57 (0.71 – 2.42)
Fifth Division	10	1.50 (1.04 – 2.52)

Mean and $\pm$ SD and median and interquartile range (IQR) values for F-SSC index. *P*-values compared between the two teams after VJT performance using Mann-Whitney U test. $\pm$ SD = standard deviation; F-SSC = fast stretch-shortening cycle

Discussion

The present study aimed to test the feasibility of the investigation protocol and to evaluate the endurance performance and the F-SSC between two amateur male soccer teams.

Study Protocol

The compliance in this current study was excellent. The 0% attrition rate for the amateur soccer players is low and the 100% adherence rate is high. This finding agrees with the previous pilot studies from Rogan et al. (2013; 2012; 2013). Performance measurements are rarely performed nowadays in the field of amateur soccer players, so it can be a motivating factor for any individual soccer player to carry out performance measurements. Not only the coach but also the players want to know their performance.

Performance outcomes

Several studies in professional soccer (Arnason et al., 2004; Cometti et al., 2001; Dellal et al., 2011; Kalapotharakos et al., 2006; Kaplan et al., 2009; Ostojic, 2004) indicate that a soccer team with high physical performance consequently has a significantly higher chance of winning and thus ranks better in the league and plays at a higher level. With amateur soccer players that play in the third and the fifth divisions, the current study did not confirm the results of the studies of professional soccer players. The following rationale can be cited.

The YYIRTL1 is a test that measures soccer players' endurance performance. This test with the nature “run-turn-run-stop-recovery-run- etc.” is similar to soccer match-play movements. The 10 sec. rest interval after 40 m shuttle run to provide the ability to recover between high-intensity shuttle run (Bangsbo, Mohr, Poulsen, Perez-Gomez, & Krstrup, 2006). Bangsbo et al. (2006) described that after YYIRTL1 in the pre-season preparation of professional soccer players, endurance performance improved by 31% with only a minor change in VO_{2max} . The ability to recover after highly intensive activities depends on good development of cardiovascular systems. The lower the heart rate after highly intensive activity (ability to recover) the higher the capacity of a soccer player (Bangsbo et al., 2006). In this current study heart rate after YYIRTL1 was not measured. However, it's possible to hypothesize that the soccer players from the third division team have more developed cardiovascular systems compared to the those in the fifth division team.

Young et al. (2005) determined that starter player in the Australian Football League (AFL) achieved significantly greater distances in the Yo-Yo intermittent recovery as a substitute. Both teams included 22 soccer players. On average 12 players are starters and 10 substitutes. In the two-amateur soccer teams a total of 70% starters completed the investigation. Among these were both goalkeepers.

A factor was that the players had not previously been made familiar with the PJT. Chu (1998) emphasizes that to achieve the maximum result when learning a new exercise, coordination and technique in jump tests play an important role. Factors such as the timing, position of the legs or angle of the knee and ankle must be practiced. The invitation to jump as high as possible after the shortest possible ground contact time is at first opposed and therefore it is difficult to perform two things at the same time. Participants who focus more on the jump height make an increased knee and hip flexion during the jumps, which results in a longer ground contact time. This corresponds more to the counter movement jump (Rogan, Radlinger, et al., 2013). For this reason, it is important that the participants must be able to perform the PJT correctly, before measurements of the F-SSC are taken. In future, the investigation procedure can begin only if the player is familiar with the PJT.

Conclusions

The scientist not only has the task to spread knowledge in professional, Olympic or university sport, but also to draw attention to amateur sport. This study shows that the study protocol is feasible. It indicates that the YYIRTL1 is a practical and sensitive tool to evaluate actual training levels for endurance performance in amateur soccer players. This study shows no significant difference between a third division and fifth division amateur soccer team in endurance performance and F-SSC. Future studies must take into account that the process of recruitment should consist of several information sessions in order to guarantee that all players will participate. In addition, the players must be able to familiarize themselves with the measurement procedures YYIRTL1 and PJT. Only when the players know the process of the YYIRTL1 and PJT should measurements be carried out at the beginning and the end of the season.

Conflicts of interest

No conflicts of interest exist

References

- Arnason, A., Sigurdsson, S., Gudmundsson, A., Holme, I., Engebretsen, L., & Bahr, R. (2004). Physical fitness, injuries, and team performance in soccer. *Medicine & Science in Sports & Exercise*(36), 2.
- Bangsbo, J., Mohr, M., Poulsen, A., Perez-Gomez, J., & Krustrup, P. (2006). Training and testing in elite athlete. *J Exerc Sci Fit*, 4(1), 1-14.
- Casartelli, N., Muller, R., & Maffiuletti, N. A. (2010). Validity and reliability of the Myotest accelerometric system for the assessment of vertical jump height. *J Strength Cond Res*, 24(11), 3186-3193. doi:10.1519/JSC.0b013e3181d8595c
- Chu, A. D. (1998). *Jumping into plyometrics* (2 ed.): Human Kinetics Publishers.
- Cometti, G., Maffiuletti, N., Pousson, M., Chatard, J., & Maffulli, N. (2001). Isokinetic strength and anaerobic power of elite, subelite and amateur French soccer players. *International Journal of Sports Medicine*, 22(1), 45-51.
- Dellal, A., Hill-Haas, S., Lago-Penas, C., & Chamari, K. (2011). Small-sided games in soccer: amateur vs. professional players' physiological responses, physical, and technical activities. *J Strength Cond Res*, 25(9), 2371-2381. doi:10.1519/JSC.0b013e3181fb4296
- Güllich, A., & Schmidtbleicher, D. (1999). [Structure of motor strength and the training methods]. *Deutsche Zeitschrift für Sportmedizin*, 50(7-8), 223-234.
- Kalapotharakos, V. I., Strimpakos, N., Vithoulka, I., Karvounidis, C., Diamantopoulos, K., & Kapreli, E. (2006). Physiological characteristics of elite professional soccer teams of different ranking. *J Sports Med Phys Fitness*, 46(4), 515-519.
- Kaplan, T., Erkmen, N., & Taskin, H. (2009). The evaluation of the running speed and agility performance in professional and amateur soccer players. *The Journal of Strength & Conditioning Research*, 23(3), 774-778.
- Krustrup, P., Mohr, M., Amstrup, T., Rysgaard, T., Johansen, J., Steensberg, A., . . . Bangsbo, J. (2003). The Yo-Yo intermittent recovery test: Physiological response, reliability, and validity. *Medicine & Science in Sports & Exercise*, 35(4), 697-705.
- le Gall, F., Carling, C., Williams, M., & Reilly, T. (2010). Anthropometric and fitness characteristics of international, professional and amateur male graduate soccer players from an elite youth academy. *J Sci Med Sport*, 13(1), 90-95. doi:10.1016/j.jsams.2008.07.004
- Ostojic, S. (2004). Elite and nonelite soccer players: Preseasonal physical and physiological characteristics. *Research in Sports Medicine*, 12(2), 143-150.
- Reilly, T., & Ekblom, B. (2005). The use of recovery methods post-exercise. *J Sports Sci*, 23(6), 619-627. doi:10.1080/02640410400021302
- Rogan, S., Blasimann, A., Nyffenegger, D., Zimmerli, N., & Radlinger, L. (2013). [The relevance of core muscles in ice hockey players: a feasibility study]. *Sportverletz Sportschaden*, 27(4), 212-218. doi:10.1055/s-0033-1356113
- Rogan, S., Blasimann, A., Steiger, M., Torre, A., & Radlinger, L. (2012). [Acute effects of fast dynamic stretching on rate of force development in ice hockey players: a pilot study]. *Sportverletz Sportschaden*, 26(4), 207-211. doi:10.1055/s-0032-1325416
- Rogan, S., Radlinger, L., Portner-Burkhalter, C., Sommer, A., & Schmidtbleicher, D. (2013). Feasibility study evaluating four weeks stochastic resonance whole-body vibration training with healthy female student. *International Journal of Kinesiology & Sport Science*, 1(2), 1-10.
- Thomas, A., Dawson, B., & Goodman, C. (2006). The yo-yo test: reliability and association with a 20-m shuttle run and VO₂(max). *Int J Sports Physiol Perform*, 1(2), 137-149.
- Wisloff, U., Castagna, C., Helgerud, J., Jones, R., & Hoff, J. (2004). Strong correlation of maximal squat strength with sprint performance and vertical jump height in elite soccer players. *Br J Sports Med*, 38(3), 285-288.