Original Article

Effect of a game-based high intensity interval training program on body composition in primary education: comparison of the Sport Education model and traditional methodology

YESSICA SEGOVIA¹, DAVID GUTIÉRREZ²

^{1,2}Faculty of Education, Ciudad Real, Castilla-La Mancha University, SPAIN.

Published online: March 31, 2020

(Accepted for publication: March 03, 2020)

DOI:10.7752/jpes.2020.02113

Abstract:

The main aim of this study was to evaluate the effects on schoolchildren's body composition of a game-based high intensity interval training (GB-HIIT) programme delivered during physical education. The secondary aim was to compare the effect of the implemented programme using two different methodologies: TheSport Education model and traditional methodology (TM). The study had a pre-post quasi-experimental design, and used a sample comprising 154 students aged between nine and 13 years (10.66±.82 years) enrolled in 5th and 6th grades of primary education. The programme lasted 15 sessions. Measurements of waist circumference and body fat percentage through bioelectrical impedance were taken. The results showed significant differences between the experimental group and control ($p > .001$) in waist circumference and body fat percentage. In conclusion, GB-HIIT appears to be effective in countering effects on schoolchildren's body composition.

KeyWords: -Pedagogical Model, Sport Education Model, Physical Education, Functional Training Exercises, Waist circumference, Body fat.

Introduction

The rise in overweight and obesity in children and adolescents is one of the primary epidemics of the 21st century. López-Sánchez et al. (2019) showed that the prevalence in children and adolescents in southern Europe ranged from 32.2% to 39.8%, depending on the scale used. Furthermore, the highest values have been associated with children who do no physical activity (PA) (Ajejas Bazán, Jiménez-Trujillo, Wärnberg, Domínguez Fernández, López de Andrés, & Pérez Farinos, 2018) or whose physical fitness (PF) is lower (Gulías-González et al, 2014). Despite uncertainty about the best strategy (setting and working method) to counter low levels of PA and PF in school-aged children, there is a consensus on considering that, as regards context, school-based interventions are the most effective and reach the largest percentage of the population (Kriemler, Meyer, Martin, Van Slujs, Andersen, & Martin, 2011). The contribution of physical education (PE) to these key health indicators could be greater if classes were programmed with this goal in mind (Fairclough & Stratton, 2005).

Regarding the working method, the literature has shown that vigorous PA is more effective than lower-intensity PA in modifying body composition in both adults (Boutcher, 2018) and children and adolescents (Owens, Galloway, & Gutin, 2015). Of the high-intensity programmes available, High-Intensity Interval Training (HIIT) appears to be the system currently yielding the best results (Boutcher, 2018). In school-age children, studies have shown its effectiveness in modifying body composition in individuals with overweight or obesity (Lambrick, Westrupp, Kaufmann, Stoner, & Faulkner, 2016) and those with low PA levels (Melrose Logan, Harris, Duncan, Plank, Merien, & Schofield, 2016), in particular when compared with moderate-intensity PA (Racil, Coquart, Elmontassar, Haddad, Goebel, Chaouachi, Amri, & Chamari, 2016). However, in their systematic review, Reyes-Amigo, Gómez, Gallardo & Palmeira (2017) reported that there was still insufficient evidence on the effect of HIIT on body composition in six- to 12-year-olds, although it has been demonstrated that HIIT can be included in PE regardless of gender, sexual maturity and cardiorespiratory fitness (Reyes-Amigo, 2019).

Calahorra-Cañada, Torres-Luque, López-Fernández and Carnero (2017) found that school students only engage in between 5.7 and 8.7 minutes of moderate to vigorous PA in each PE. Such a small workload, if based on high-volume exercise such as continuous training, would mean minimal impact on achieving PF and health aims. HIIT represents a clear alternative in order to achieve these objectives since benefits for body composition in children aged between seven and 12 years have been reported with less than nine minutes of vigorous PA per session (Rosenkranz, Rosenkranz, Hastmann, & Harms, 2012). Moreover, this system would allow the teacher to have time to meet other educational objectives. In this sense, the emergence of interest in HIIT is a consequence of both its efficacy in modifying the cardiorespiratory fitness and cardiometabolic health of children and

adolescents (Bond, Weston, Williams, & Barker, 2017) and its efficiency (good results with a minimum investment of time).

Despite the growing interest in HIIT in recent years, as demonstrated by the publication of various systematic reviews of its application in school-aged children (Bond et al., 2017; Reyes-Amigo et al., 2017), few studies have been conducted in school settings to assess the impact of HIIT on body composition. This lack of studies is even more evident if we limit the search to the context of PE, where only five studies were found (Delgado-Floody, Espinoza-Silva, García-Pinillos, & Latorre-Román, 2018; Huerta et al., 2017; Muntaner & Palou, 2017; Reyes-Amigo, 2019; Weston, Azevedo, Bock, Weston, George, & Batterham, 2016).

Of these cited studies, only those by Delgado-Floody et al. (2018) and Reyes-Amigo (2019) were conducted in primary school settings. The study by Delgado-Floody et al. (2018) revealed changes in body composition in pupils with overweight or obesity. The boys with obesity exhibited significant reductions in waist circumference (WC) and body fat percentage (BF%), while girls only showed significant changes in BF%. In the children with overweight, significant improvements were only found for BF% and only in girls. In the control group (obesity and overweight), BF% increased significantly. In contrast, the study by Reyes-Amigo (2019) did not differentiate between the children's bodyweight status. The findings showed significant differences in PF between the HIIT group and the group engaging in a programme of moderate-intensity games, with this variable decreasing only in the moderate-intensity group. Both programmes yielded significant results in the reduction of BF%, with no inter-group differences.

HIIT, when used in PE at schools, could provide a response to the demand for efficacy and efficiency in work on PF. However, as a training system, it may not be sufficiently motivating and meaningful to make this work on PF effective in long-term educational goals, especially as regards adherence to PA. In this sense, teaching based on pedagogical models could be an answer to these core PE questions, since the rationale of the implementation of these models (what is known as models-based practice) is the achievement of a series of learning outcomes, for which multiple versions or "models" of PE would be needed (Kirk, 2013). Among the leading pedagogical models is the Sport Education model. The motivation it generates in students has been widely studied and its potential has been demonstrated in different genders, levels and sports, and in students with different motivational profiles (Chu & Zhang, 2018) and in different cultural contexts (Gutiérrez, García-López, Hastie, & Calderón, 2013). With regard to motivation, studies highlight that Sport Education, compared to a traditional approach, regulates motivation more effectively (Fernández-Río, Méndez-Giménez, & Méndez-Alonso, 2017), significantly improves intrinsic motivation (Cuevas, García-López, & Serra-Olivares, 2015) and can facilitate positive change in students with low levels of motivation (Perlman & Caputi, 2017).

The primary objective of Sport Education is to offer a more authentic sport experience in the context of school PE, with its main aims being the training of competent, literacy and enthusiastic sport practitioners (Siedentop, 1994). To do this, it maintains the core characteristics of sport (a season, affiliation with a team, competition, culminating event, record keeping and festivity), while adapting the sport content to the demands of education and introducing roles beyond that of player/learner (e.g., coach, referee, reporter, etc.) (Siedentop, Hastie, & van der Mars, 2019).

The plasticity characteristic of Sport Education has meant that, despite being a model with its basis in sports, teaching units (seasons, in Sport Education terminology) have been successfully implemented using content not directly related to sport games (García-López & Gutiérrez, 2016). Among these, fitness content stands out for its contribution to objectives related to health and FP. Various studies have shown the potential of Sport Education to improve fitness levels, fitness knowledge and PA levels in school students (Ward, Hastie, Wadsworth, Foote, Brock, & Hollett, 2017). However, to the best of our knowledge, no study has analysed the possible effect of this model on school-aged children's body composition.

The main aim of this study was to evaluate the effects on schoolchildren's body composition of a game-based HIIT (GB-HIIT) programme delivered during PE. The secondary aim was to compare the effect of the programme implemented using two different methodologies: Sport Education and traditional methodology (TM).

Material & methods

Participants

This study was conducted in three state primary schools in the province of Ciudad Real (Spain), with the participation of eight natural groups (five 5th grade classes and three 6th grade classes), comprising a total of 154 pupils (81 boys and 73 girls) with ages ranging from nine to 13 years (10.66 ± 0.82 years). Four teachers (two males and two female) also took part in the study.

The students were assigned to either the experimental group ($n=103$) or the control group ($n=51$). Table I shows the percentages of students according to BF% (McCarthy, Cole, Fry, Jebb, & Prentice, 2006). The experimental group was divided into two further groups depending on whether their programme followed Sport Education ($n=59$) or TM ($n=44$). The children in Sport Education group had no previous experience of this methodology.

Table 1. Descriptive characteristics of the initial sample. Body status according to BF% of the student body.

	Low	Normal	High or very high
Experimental group	17.8%	59.4%	22.8%
Control group	19.6%	62.7%	17.6%

The participants were selected by means of incidental sampling, using natural groups based on their viability and access. Participation in the study was voluntary with both parents and children giving their consent.

Method

The study design was quasi-experimental with pre- and post-test measures, using an experimental group (Sport Education and TM) and a control group. The participants were not probabilistically allocated to groups due to the prior assignation of the students in natural groups at the schools. Nonetheless, despite the natural groups being selected by convenience sampling, the students were randomly assigned to the TM, Sport Education and control group.

The study was divided into four phases. First, the GB-HIIT was designed and validated. Second, pre-test assessment was conducted in all the groups. Third, the GB-HIIT programme was implemented in the experimental group (Sport Education and TM), while the control group followed its study plan (15 sessions of a season of ringo). And fourth, once the programme had been completed, the post-test assessment was carried out.

Season design and lesson content

A 15-session, game-oriented PF programme was designed using the HIIT system and functional exercises. The intervention was implemented in the schools' PE classes. Although we used two different methodologies (Sport Education and TM), the workload and motor activities were the same in both programmes. Each week, the different class groups in the experimental group had either three weekly 45-minute sessions or two weekly 60-minute sessions.

The content was adapted to Sport Education following the guidelines described by García-López & Gutiérrez (2016) and Siedentop, Hastie & van der Mars (2019). The season was divided into four stages (organisation, pre-season, regular stage, and final stage). Each class group was organised into four socially compatible groups, which, in terms of PF, were heterogeneous at within-group level but balanced at between-group level. Apart from participating as an athlete, each student had a dual team and organisational role, taking on the responsibilities for both the team role they were given (coach, fitness coordinator, nutritionist, journalist, manager) and the organizational role (main referee, event referee, scorekeeper or competition coordinator). The TM intervention used none of the methodological strategies associated with Sport Education, such as permanent teams, roles or festivities, while the competition was replaced by activities in which techniques were co-evaluated.

With regard to content, the sessions consisted of two main motor activities: Moderate to Vigorous Intensity Game (MoVinG) and Recreational and Intense Circuit (Randin). These activities were designed specifically to include three characteristics: playful, high intensity and functional training exercises.

The MoVinG were games lasting five to seven minutes, incorporating the technique of functional exercises. The Randin were implemented following the HIIT system. The length of the Randin increased progressively from five (sessions 1-5) to eight minutes (sessions 6-15). Intervals lasted 40 seconds at an intensity of 85-90% and 20 seconds of recovery (no movement). Randin were used for both training and as the competitive activity.

In order to determine and validate the intensity of the sessions, in one typical training session and another competition session, we monitored 20 students, 10 from each methodology group. The participants were selected taking into account a balance of gender and different levels of physical fitness. Mean heart rate data were obtained for the complete session and the high-intensity periods. The data confirmed the children were working at an intensity of 85-90% during the MoVinG and the Randin. Data were collected using the Polar A300 heart rate monitor.

The design and validation of the programme is described in Segovia and Gutiérrez (2018).

Data collection

Body composition was measured during the PE classes, following the guidelines set out in the high priority ALPHA-Fitness battery, validated for children and adolescents (Ruiz et al., 2011). We measured WC and, instead of BMI, we took bioelectrical impedance measurements of BF%, which is considered to be a more reliable method (Nuttall, 2015) and is validated in children (Lee et al., 2017).

WC. A tape measure was used to measure the WC at the midpoint between the last rib and the iliac crest using a tape measure.

BF%. Bioelectrical impedance measurements of BF% were taken using the Tanita BC-418Body Composition Analyser. Although the children were measured barefoot and wearing light clothes, 1kg was subtracted from the measurements. Height was measured to the nearest 0.1cm, using the Seca 222 measuring rod, with the children barefoot and standing straight.

The examiners were given theoretical and practical assistance in all the tests by experts in healthcare research.

Data analysis

The statistical analysis was conducted using the IBM-SPSS 21.0 statistical software package. The normality distribution of the variables was calculated using the Kolmogorov-Smirnov test (more than 50 cases) and the Shapiro-Wilk test (fewer than 50 cases). The results showed a non-normal distribution and hence non-parametric tests were used. Descriptive pre-post statistics (intra-group analysis) and the variable “program effect” (post-test – pre-test) (inter-group effect) were calculated. An inter-group comparison was conducted using baseline values to determine initial differences and the differences in the programme’s effectiveness based on a comparison of the differences between pre-and post-test mean scores. The U Mann-Whitney test was used to analyse differences according to the programme content (two groups: GB-HIIT and control) and the Kruskal Wallis test was used to analyse differences according to methodology (three groups: TM, Sport Education and control). In the U Mann-Whitney test, Bonferroni’s correction was applied to the alpha value in order to control for Type 1 error ($\alpha = .05 / 3 = .017$). The Wilcoxon test was used to determine intra-group differences. The r statistic (effect size) for non-parametric tests was estimated and interpreted, using a scale of small (.10), medium (.30) and large (.50). For the Wilcoxon test, the effect size was calculated using $r = Z/\sqrt{N}$, where N was the number of measurements.

Results*Results according to programme content (GB-HIIT group versus control)*

No significant differences were found in the initial characteristics of the sample (GB-HIIT versus control) for any of the variables.

Intra-group analysis

In the overall sample (Table II), significant differences were only revealed in the control group, which showed significant increases in WC and BF%, with a small and large effect size, respectively. The analysis by gender (Table II) yielded significant changes in both the GB-HIIT group and the control group. In the GB-HIIT group, the WC in boys decreased significantly, with a small effect size. However, in the control group, significant differences were found in both boys and girls. The girls showed a significant increase in both WC and BF%, with a medium and a large effect, respectively. A significant increase only in BF%, with a medium effect size, was found in the case of the boys.

Table II. *Descriptive statistics (pre-test and post-test) and intra-group analysis of WC and %BF according to programme content. Wilcoxon test*

	<i>Pre</i>	<i>Post</i>			
	M(SD)	M(SD)	p	Z	r
WC					
		GB-HIIT			
Total	70.35(10.87)	69.94(10.62)	.147	-1.451	.10
Boys	70.67(11.21)	69.87(10.88)	.030	-2.170	.21
Girls	70.00(10.58)	70.01(10.44)	.972	-.035	.00
		Control			
Total	74.58(11.95)	75.79(12.06)	.001	-3.257	.32
Boys	77.13(12.05)	77.83(11.91)	.115	-1.576	.21
Girls	71.71(11.42)	73.50(12.06)	.003	-2.945	.43
BF%					
		GB-HIIT			
Total	22.55(8.81)	22.37(8.90)	.749	-.320	.02
Boys	20.09(8.48)	19.84(8.43)	.146	-1.453	.14
Girls	25.26(8.43)	25.16(8.65)	.269	-1.106	.11
		Control			
Total	21.30(9.03)	22.59(8.94)	<.001	-4.542	.45
Boys	20.28(7.84)	21.14(7.63)	.025	-2.247	.31
Girls	22.45(10.25)	24.23(10.12)	<.001	-4.287	.62

M=mean; SD= standard deviation

According to bodyweight status by BF%, significant differences were only found in the increase in BF% in normal status ($p > .001$) and low status ($p = .012$) and the increase in WC ($p = .006$) in normal status.

Inter-group analysis

The Mann-Whitney U test (Table III), used to test for differences between the GB-HIIT group and the control group, revealed significant differences in both variables for both the complete sample and by gender.

Table III. *Descriptive statistics (program effect) and inter-group analysis of WC and %BF according to programme content. Mann-Whitney U test*

	GB-HIIT	Control				
	M(SD)	M(SD)	<i>U</i>	<i>p</i>	<i>Z</i>	<i>R</i>
WC						
Total	-.35(2.43)	1.22(2.37)	1672.000	<.001	-3.537	.29
Boys	-.69(2.24)	.70(2.30)	468.000	.015	-2.432	.27
Girls	.01(2.59)	1.79(2.35)	356.500	.006	-2.725	.32
BF%						
Total	.02(1.88)	1.29(1.74)	1435.00	<.001	-4.594	.36
Boys	-.19(2.07)	.86(2.08)	445.500	.006	-2.749	.31
Girls	.26(1.63)	1.78(1.10)	280.500	<.001	-3.772	.42

M=mean; SD= standard deviation

According to bodyweight status by BF%, significant differences were found in children with normal status ($p=.001$) in both variables, and in BF% in low status ($p=.031$), coinciding with the data reported in previous section.

Results according to methodology (sport education model versus traditional methodology versus control group)
The Kruskal Wallis test revealed no significant differences in the initial characteristics of the sample.

Intra-group analysis

The analysis of the data dividing the experimental GB-HIIT group according to methodology (TM and Sport Education) revealed no significant changes either in the overall sample or by gender (in contrast to the results of the whole group analysis, see Results according to programme content (GB-HIIT group versus control) without-group results) (Table IV). Changes were found, however, in the control group, as reported in results according to programme content (GB-HIIT group versus control) without-group results.

Table IV. *Descriptive statistics (pre-test and post-test) and intra-group analysis of WC and %BF according to methodology. Wilcoxon test*

	<i>Pre</i>	<i>Post</i>			
	M(SD)	M(SD)	<i>p</i>	<i>Z</i>	<i>r</i>
WC					
TM					
Total	70.77(10.73)	70.42(11.21)	.404	-.835	.09
Boys	71.52(10.99)	70.78(11.30)	.208	-1.259	.19
Girls	70.02(10.68)	70.09(11.38)	.935	-.081	.01
Sport Education					
Total	70.03(11.05)	69.59(10.26)	.238	-1.179	.11
Boys	70.08(11.50)	69.30(10.76)	.071	-1.808	.23
Girls	69.98(10.70)	69.94(9.82)	.936	-.081	.01
Control					
Total	74.58(11.95)	75.79(12.06)	.001	-3.257	.32
Boys	77.13(12.05)	77.83(11.91)	.115	-1.576	.21
Girls	71.71(11.42)	73.50(12.06)	.003	-2.945	.43
%GC					
TM					
Total	23.51(9.13)	23.43(9.04)	.675	-.419	.04
Boys	21.44(9.16)	21.15(8.95)	.390	-.860	.13
Girls	25.57(8.83)	25.71(8.74)	.808	-.244	.04
Sport Education					
Total	21.81(8.55)	21.58(8.79)	.925	-.095	.01
Boys	19.14(7.98)	18.95(8.08)	.213	-1.244	.16
Girls	25.00(8.24)	24.71(8.72)	.236	-1.185	.16
Control					
Total	21.30(9.03)	22.59(8.94)	<.001	-4.542	.45
Boys	20.28(7.84)	21.14(7.63)	.025	-2.247	.31
Girls	22.45(10.25)	24.23(10.12)	<.001	-4.287	.62

M=mean; SD= standard deviation

Inter-group analysis

Table V and VI illustrates the inter-group differences by methodology and gender. The analysis revealed significant differences between the experimental group (TM and Sport Education) and the control, with a medium effect size in both WC and BF%.

By gender, we found significant differences in BF% in girls between the two experimental group (TM and Sport Education) and control, with large and medium effect size, respectively. With regard to WC, differences were only found between the TM group and control, with a medium effect size. In the boys, significant differences were found for both variables but only between Sport Education group and control, with a small effect size.

Table V. *Descriptive statistics (program effect) and inter-group analysis of WC and %BF according methodology. Kruskal-Wallis*

	TM	Sport Education	Control	Kruskal-Wallis	
	M(SD)	M(SD)	M(SD)	p	χ^2
WC					
Total	-.23(2.27)	-.44(2.56)	1.22(2.37)	.002	12.604
Boys	-.55(1.97)	-.78(2.43)	.70(2.30)	.044	6.245
Girls	.07(2.52)	-.04(2.70)	1.79(2.35)	.024	7.476
BF%					
Total	-.08(1.77)	.10(1.98)	1.29(1.74)	<.001	19.023
Boys	-.29(1.75)	-.12(2.30)	.86(2.08)	.022	7.612
Girls	.14(1.80)	.37(1.52)	1.78(1.10)	.001	14.402

M=mean; SD= standard deviation

Table VI. *Inter-group analysis of WC and %BF according methodology. Mann-Whitney U*

	Groups	U	p	Z	R
WC					
Total	MT-Sport Education	1176.000	.663	-.436	.08
	TM-Control	698.500	.004	-2.888	.32
	Sport Education-Control	973.500	.001	-3.191	.30
Boys	TM-Sport Education	287.000	.532	-.626	.16
	TM-Control	193.000	.095	-1.668	.30
	Sport Education-Control	275.000	.016	-2.398	.31
Girls	Sport Education-Control	289.000	.872	-.161	.02
	TM-Control	151.500	.013	-2.482	.36
	Sport Education-Control	205.000	.024	-2.252	.31
BF%					
Total	TM-Sport Education	1225.000	.842	-.199	.05
	TM-Control	577.500	<.001	-4.066	.42
	Sport Education-Control	858.000	<.001	-3.666	.37
Boys	TM-Sport Education	318.500	.684	-.407	.08
	TM-Control	182.000	.021	-2.315	.33
	Sport Education-Control	263.500	.016	-2.418	.33
Girls	TM-Sport Education	264.500	.656	-.446	.09
	TM-Control	110.500	.001	-3.378	.50
	Sport Education-Control	150.000	.002	-3.149	.46

Discussion

The main aim of this study was to evaluate the effects of GB-HIIT programme delivered during PE on the body composition of 5th and 6th grade schoolchildren. The secondary aim was to compare the effect of the methodology used to implement the programme (Sport Education and TM). The results showed GB-HIIT is effective in modifying the body composition of primary school children. This effectiveness was supported by comparing with the data from the control group, which increased significantly in the study variables. In the intra-group analysis, the participants in the experimental group exhibited a general decrease in all the values, although this decrease was only significant in the case of boys' WC. Moreover, the methodology used to implement the GB-HIIT programme had no impact on the findings.

The effectiveness of the GB-HIIT programme lies primarily in stabilization, as shown by the non-increase in the scores on the variables associated with the children's body composition, which did, however, increase in the control group. These findings are consistent with those of the study by Delgado-Floody et al.

(2018), where, following a 48-session programme conducted with children with overweight and obesity, the experimental group significantly improved their body composition compared with the control, whose body composition worsened. In contrast to our study, however, Delgado-Floody et al. (2018) reported both enhancements in the experimental group and deteriorations in the control group. This difference is arguably due to both the length of the programme and the characteristics of the sample, specifically the children's bodyweight status.

Furthermore, our results differ from those of the study by Reyes-Amigo (2019), despite similar in the diversity of status-weight and also in the duration of the program. In this case, the high-intensity programme reduced the body fat percentage but yielded less positive results in WC compared with the moderate-intensity programme.

Among the three studies we found that were conducted in secondary schools, only that by Weston et al. (2016) with 14 year-olds, reported significant differences in WC. These authors' findings are similar to ours, that is, they found differences between the intervention groups and control, mainly due to increased WC values in the control group.

As regards the methodology used, to the best of our knowledge, no previous study has analysed the effect of Sport Education methodology on body composition. Although the study yields no differences between methodologies in the study variables, the previous literature has demonstrated the educational potential of Sport Education compared to traditional methodology in variables that might improve PF, such as motivation (Chu & Zhang, 2018), as well as others such as moral and ethical development (Harvey, Kirk, & O'Donovan, 2014). Hence, given the similar results obtained with both methodologies using PF content and the GB-HIIT, we recommend the implementation of Sport Education, as it can enhance and promote a larger number of educational objectives in PE classes.

The present study has two main limitations. The first is the sample size, which prevents the generalization of the results obtained. However, despite its limited size, it is larger than that used in most HIIT studies in children and adolescents. The second limitation is that, despite having a control group to determine the effectiveness of the programme, its effects were not analysed taking possible confounders into account, such as biological maturity, socioeconomic status or diet.

Conclusions

The lack of studies in the field of PE with characteristics similar to those of the present work makes ours a novel study with the potential to open up a new research line addressing the efficacy of GB-HIIT in PE and its effect depending on the different pedagogical approaches implemented. In addition, it would also be interesting to study variables that might have a greater impact on the methodology used, such as variables of motivation, body image or adherence to PA. In sum, the results obtained in this study appear to show that increased body fat in primary school children, due to diet and/or body development, could be countered by the use of GB-HIIT sessions in PE classes. The impact on long-term educational goals is as yet unknown, but the methodology implemented appears to have no impact on our study variables.

Funding

Yessica Segovia is supported by a grant from the University of Castilla-La Mancha (2018-CPUCLM-7487).

Conflicts of interest -The authors declare no conflict of interest.

References

- Ajejas Bazán, M. J., Jiménez-Trujillo, M. I., Wärnberg, J., Domínguez Fernández, S., López-de-Andrés, A., y Pérez-Farinos, N. (2018). Prevalence of childhood overweight/obesity in Spain 1993-2011 and associated risk factors in 2011. *Nutrición Hospitalaria*, 35, 84-89.
- Bond, B., Weston, K., Williams, C., & Barker, A. (2017). Perspectives on high-intensity interval exercise for health promotion in children and adolescents. *Journal of Sports Medicine*, 8, 243-265, 2017.
- Boutcher, S.H. (2018). Ejercicio Intermitente de Alta Intensidad y Pérdida de Grasa. *Revista de Educación Física*, 36(4), 1-13.
- Calahorra-Cañada, F., Torres-Luque, G., López-Fernández, I., & Carnero, E. (2017). Is physical education an effective way to increase physical activity in children with lower cardiorespiratory fitness? *Scandinavian Journal of Medicine & Science in Sports*, 27(11), 1417-42.
- Chu, T.L., & Zhang, T. (2018). Motivational processes in Sport Education programs among high school students: A systematic review. *European Physical Education Review*, 24(3), 372-394.
- Cuevas, R., García-López, L.M., & Contreras, O. (2015). Influencia del modelo de Educación Deportiva en las necesidades psicológicas básicas. *Cuadernos de Psicología del Deporte*, 15(2), 155-162.
- Delgado-Floody, P., Espinoza-Silva, M., García-Pinillos, F., & Latorre-Román, P. (2018). Effects of 28 weeks of high-intensity interval training during physical education classes on cardiometabolic risk factors in Chilean schoolchildren: a pilot trial. *European Journal of Pediatrics*, 177, 1019-1027.

- Fairclough, S., & Stratton, G. (2005). Physical education makes you fit and healthy'. Physical education's contribution to Young people's physical activity levels. *Health Education Research*, 20(1), 14-23.
- Fernández-Río, J., Méndez-Giménez, A., & Méndez-Alonso, D. (2017). Efectos de dos formatos instructivos, Educación Deportiva e Instrucción Directa, en la respuesta psicológica de estudiantes de secundaria. *Sport TK: Revista Euroamericana de Ciencias del Deporte*, 6(2), 9-20.
- García López, L.M., & Gutiérrez, D. (2016). *Aprendiendo a enseñar deporte. Modelos de Enseñanza Comprensiva y Educación Deportiva*. Barcelona: INDE.
- Gulías-González, R., Martínez-Vizcaíno, V., Cañete García-Prieto, J., Díez-Fernández, A., Olivas-Bravo, A., & Sánchez-López, M. (2014). Excess of weight, but not underweight, is associated with poor physical fitness in children and adolescents from Castilla-La Mancha, Spain. *European Journal of Pediatric*, 173, 727-735.
- Gutiérrez, D., García-López, L. M., Hastie, P., & Calderón, A. (2013). Spanish Students' Perception of Their Participation in Season of Sport Education. *The Global Journal of Health and Physical Education Pedagogy*, 2(2), 111-129.
- Harvey, S., Kirk, D., & O'Donovan, T.M. (2014). Sport Education as a pedagogical application for ethical development in physical educations and youth sport. *Sport, Education and Society*, 19(1), 41-62.
- Huerta, A., Galdames, S., Cataldo, M., Barahona, G., Rozas, T., & Cáceres, P. (2017). Efecto de un entrenamiento intervalado de alta intensidad en la capacidad aeróbica de adolescentes. *Revista Médica de Chile*, 145(8), 972-979.
- Kirk, D. (2013). Educational value and models-based practice in physical education. *Educational Philosophy and Theory*, 45(9), 973-986.
- Kriemler, S., Meyer, U., Martin, E., van Sluijs, E.M., Andersen, L.B., & Martin, B.W. (2011). Effect of school-based interventions on physical activity and fitness in children and adolescent: a review of review and systematic update. *British Journal of Sports Medicine*, 45(11), 923-930.
- Lambrick, D., Westrupp, N., Kufmann, S., Stoner, L., & Faulkner, J. (2016). The effectiveness of a high-intensity games intervention on improving indices of health in young children. *Journal of Sports Sciences*, 34(3), 190-198.
- Lee, L.W., Liao, Y.S., Lu, H.K., Hsiao, P.L., Chen, Y.Y., Chi, C.C., y Hsieh, K.G. (2017). Validation of two portable bioelectrical impedance analyses for the assessment of body composition in school age children. *Plos One*, 12(2), e0171568.
- López-Sánchez, G. F., Sgroi, M., D'Ottavio, S., Díaz-Suárez, A., González-Villora, S., Veronese, N., & Smith, L. (2019). Body Composition in Children and Adolescents Residing in Southern Europe: Prevalence of Overweight and Obesity According to Different International References. *Frontiers in Physiology*. <https://doi.org/10.3389/fphys.2019.00130>
- McCarthy, H. D., Cole, T. J., Fry, T., Jebb, S. A., & Prentice, A. M. (2006). Body fat reference curves for children. *International journal of obesity*, 30(4), 598-602.
- Melrose Logan, G. R., Harris, N., Duncan, S., Plank, L. D., Merien, F., & Schofield, G. (2016). Low-Active Male Adolescents: A Dose Response to High-Intensity Interval Training. *Medicine & Science in sports & Exercise*, 481-490.
- Muntaner-Mas, A., & Palou, P. (2017). Effects of high intensity interval training (HIIT) intervention amongst school adolescents. *Journal of Physical Education & Health*, 6(10), 19-25.
- Nuttall, F.Q. (2015). Body Mass Index: Obesity, BMI, and Healty: A Critical Review. *Nutrition today*, 50(3), 117-128.
- Owens, S. Galloway, R., & Gutin, B. (2017). The Case for Vigorous Physical Activity in Youth. *American Journal of Lifestyle Medicine*, 11(2), 96-115.
- Perlman, D., & Caputi, P. (2017). Examining the influence of Sport Education on the precursors of amotivation. *European Physical Education Review*, 23(2), 212-222.
- Racil, G., Coquart, J.B., Elmontassar, W., Haddad, M., Goebel, R., Chaouachi, A., Amri, M., & Chamari, K. (2016). Greater effects of high-compared with moderate-intensity interval training on cardio-metabolic variables, blood leptin concentration and ratings of perceived exertion in obese adolescent females. *Biology of Sport*, 33(2), 145-152.
- Reyes-Amigo, T., Gómez, M., Gallardo, M., y Palmeira, A. (2017). Effectiveness of high-intensity interval training on cardiorespiratory fitness and body composition in preadolescents: a systematic review. *European Journal of Human Movement*, 39, 32-47.
- Reyes-Amigo, T. (2019). Ejercicio físico de alta intensidad en niños: Efecto de juegos intermitentes de alta intensidad sobre la aptitud cardiorrespiratoria y la composición corporal en escolares [Doctoral Thesis]. Universidade Lusófona de Humanidades e Tecnologias, Faculdade de Educação Física e Desporto, Lisboa. Available from: <http://dspace.ulusofona.pt/handle/10437/9551>
- Rosenkranz, S. K., Rosenkranz, R. R., Hastmann, T. J., & Harms, C. A. (2012). High-intensity training improves airway responsiveness in inactive nonasthmatic children: evidence from a randomized controlled trial. *Journal of Applied Physiology*, 112(7), 1174-1183.

- Ruiz, J.R., España Romero, V., Castro Piñero, J., Artego, E.G., Ortega, F.B., Cuenca García, M., ... Castillo, M.J. (2011). Batería ALPHA-Fitness: test de campo para la evaluación de la condición física relacionada con la salud en niños y adolescentes. *Nutrición Hospitalaria*, 26, 1210-1214.
- Segovia Y., & Gutiérrez, D. (2018). Percepción del profesorado y el alumnado sobre la aplicación HIIT en Educación Física. Modelo de Educación Deportiva Versus Metodología Tradicional. *Ágora para la Educación Física y el Deporte*, 20(2-3), 133-161.
- Siedentop, D. (1994). *Sport Education: Quality PE through positive sport experiences*. Champaign, IL: Human Kinetics.
- Siedentop, D., Hastie, P. A., & van der Mars, H. (2019). *Complete guide to Sport Education*. (3rd ed). Champaign, IL: Human Kinetics.
- Ward, J. K., Hastie, P., Wadsworth, D. D., Foote, S., Brock, S. J., & Hollett, N. (2017). A Sport Education Fitness Season's Impact on Students' Fitness Levels, Knowledge, and In-Class Physical Activity. *Research Quarterly for Exercise and Sport*, 88(3), 346-351.
- Weston, K. L., Azevedo, L. B., Bock, S., Weston, M., George, K. P., y Batterham, A. M. (2016). Effects of Nivel, School-Based High-Intensity Interval Training (HIT) on Cardiometabolic Health in adolescents: Project FFAB (Fun Fast activity Blasts)- An Exploratory Controlled Before-And-After Trial. *Plos One*, 11(8), e0159116.