Original Article

Physiological foundations of load modeling in the annual training cycle of highly qualified canoe slalom athletes

DARYA OKUN¹, MARYNA KOROLOVA², SVETLANA STADNIK³, VLADYSLAV ROZHKOVA⁴,
LARYSA TARAN⁵, MAKSYM MISHYN⁶, IRINA PETRENKO⁷, OLENA TARASEVICH⁸
^{1, 2, 3, 4, 5, 6, 7, 8}Kharkiv State Academy of Physical Culture, Kharkiv, UKRAINE

Published online: September 30, 2020

(Accepted for publication: September 22, 2020)

DOI:10.7752/jpes.2020.05365

Abstract:

The paper presents the results of a study of the volume and intensity of the load in the annual training cycle of the national team of Ukraine in canoe slalom. The interest in building a one-year training cycle in various sports always remains in the center of attention of specialists, since it is the one-year planning of the training process that is the culmination of all the trainer's creative work, embodied in a specific document, on the basis of which all other training cycles, individual classes and motor tasks are detailed. Purpose of the study is to determine the optimal volume and intensity of training loads in building the training process of the national team of Ukraine in canoe slalom. In the course of the study, it was revealed that the total rowing volume in various intensity zones was on average 5115.26 ± 283.49 km (100%). As for the ratio of volumes in different zones of muscle work intensity, the rowing volume in the zone of moderate power averaged 2383.36 ± 142.74 km. With such rowing intensity, when the concentration of lactate does not exceed 2 mmol / l, the energy supply of muscle activity, that is, physical work, is carried out exclusively due to the aerobic mechanism. Under such conditions, the heart rate (HR) of athletes is within 70–80% of the maximum, and the intensity of the training load corresponds to 80–90% of the level of the anaerobic threshold (ANT). It should be noted that the share of rowing in this power zone is quite large and averages 46.6% of the total rowing volume. It is established that the total mileage of rowing on the water is on average 5115 km. Of these: rowing in the moderate power zone - 2383 km (46.6%); rowing in the submaximal power zone - 2419 km (47.1%); rowing in the zone of high power - 229 km (4.5%); rowing in the zone of maximum power - 92 km (1.8%). The created model of the volume and intensity of training loads can be considered basic in the construction of the training process of highly qualified rowers.

Key words: training process, muscle work, rowing, training loads.

Introduction

The basis for building a one-year training cycle should be based on certain patterns that reflect the essential characteristics of the process of sports training. However, experts on this subject do not have an unambiguous opinion and, as a result, there are several models of the structure of the annual training cycle, moreover, fundamentally different from each other (Braghin, F., Cheli, F., Melzi, S. & Sabbioni, E. 2008, Harwood, C.G., & Knight, C.J. 2015, Messias, L. H. D., et al. 2014, Holt, N.L., Tamminen, K.A., Black, D.E., Mandigo, J.L., & Fox, K.R. 2009). The interest in building a one-year training cycle in various sports always remains in the center of attention of specialists, since it is the one-year planning of the training process that is the culmination of all the trainer's creative work, embodied in a specific document, on the basis of which all other training cycles, individual classes and motor tasks are detailed (Messias, L. H. D., et al. 2018, Hunter, A. 2009, Hunter, A., Cochrane, J. & Sachlikidis, A. 2007, Okun, D. 2016, Sigmund, M., Rozsypal, R., Kudlacek, M., Kratochvil, J., & Sigmundová, D. 2016, Zeimers, G., Anagnostopoulos, C., Zintz, T., & Willem, A. 2019).

Interest in building a one-year training cycle in various sports always remains in the center of attention of specialists. It is not at all accidental, since it is the annual planning of the training process that is the culmination of all the work of the trainer, embodied in a specific document, on the basis of which all other training cycles, individual lessons and motor tasks are detailed (Okun, D. 2016, Okun D., Mulyk K. 2017, Matveev, L.P. 1999). Certainly, the basis for the construction of the annual cycle should be based on certain patterns that reflect the essential characteristics of the process of sports training. However, specialists do not have an unambiguous opinion on this matter, and, as a consequence, there are several models of the structure of the annual training cycle, and, moreover, they are fundamentally different from each other.

Purpose of the study is to determine the optimal volume and intensity of training loads in building the training process of the national team of Ukraine in canoe slalom.

Material & methods

Studies were conducted at training facilities in the cities of Kharkov, Kiev and Nikolaev with members of the national rowing slalom team. A total of 58 rowers took part in the experiment from 2016 to 2019, including 24 male kayakers, 16 female kayakers, 18 male canoeists, 8 female canoeists.

In the course of the study, we used the following materials and methods:

1. Analysis of scientific and methodological literature on the problem of modeling the training loads of athletes.
2. Analysis of the effectiveness of training programs of the national team of Ukraine in canoe slalom for 2016-2019 according to the results of international competitions.
3. Analysis of training volumes and intensity of loads in micro-, meso- and macrocycles for the preparation of highly qualified canoe slalom athletes.
4. The method of individualization of the preparation of highly qualified canoe slalom athletes based on the kinetics of endogenous lactate.
5. Methods of mathematical statistics.

Results

Currently, when preparing training programs for training the Ukrainian team in canoe slalom, the following fixed assets are included in their content: special training (rowing in various intensity zones); general physical training (athletics running (km), fitness training, athletic training, sports, general developmental exercises). It is traditionally planned to perform kayaking and canoeing in five intensity zones, depending on the concentration of lactate formed in this process. (La):

The first zone is lactate less than 2 mmol / l. The second zone is lactate from 2 to 4 mmol / l. The third zone is lactate from 4 to 8 mmol / l. The fourth zone is lactate over 8 mmol / l.

The fifth zone is alactate, i.e., without lactate in the body.

Physical performance is based on certain physiological mechanisms, the study of which is important in monitoring the training of athletes who train endurance. Aerobic performance is determined by the functional reserves of the oxygen transporting system (cardiovascular, respiratory, circulatory) and tissue respiration systems. Anaerobic performance is determined by the power of intracellular anaerobic systems and muscle energy reserves. Depending on the power of work, classifications of muscle work were proposed. In the classification of Ferrari, H. G. et al 2017, four degrees of work power are distinguished, which correspond to four time zones.

The maximum power operation is performed in the zone, the duration of which is 10–20 s. Submaximal power work is carried out in an area lasting from 20 s to 5 min. In the zone from 5 minutes to 30 minutes, high power work is performed. Less powerful work is performed in the fourth zone (moderate power). The duration of work in the moderate power zone is over 30 minutes.

In the classification of Matveev, L.P. 1999, the first zone (maximum power zone) lasts no more than 15 s.

The second zone (submaximum power zone) is divided into two, where the duration of the first is from 15 s to 40 s, and the second from 40 s to 2 min. The fourth zone (high power zone) is from 2 minutes to 10 minutes. The fifth zone (moderate power zone) is over 10 minutes of rowing. In the first zone, work is mainly provided by the creatine phosphate mechanism of energy production (alactate phase of anaerobic metabolism). In the second zone (from 15 s to 40 s), glycolysis (lactate phase of anaerobic metabolism) plays the main role. In the third zone (from 40 s to 2 minutes), along with glycolysis, the aerobic mechanism of energy production is activated.

In the fourth zone, the aerobic mechanism plays a predominant role. In the fifth zone, the bulk of the work is carried out due to the aerobic mechanism of energy supply of physical activity. In relation to kayaking and canoeing, the following distances, according to Matveev, L.P. 2010, include the following distances: 3rd zone (submaximum power zone) - 500 m distance; 4th zone (high power zone) - distance of 1,000 m; 5th zone (zone of moderate power) - distance 5,000 m Messias, L. H. D., et al. 2015.

As is known Manchado-Gobatto, F. B., et al. 2014, the first intensity zone corresponds exclusively to the aerobic regime of energy supply of muscle activity. The second zone falls on the border of the threshold of anaerobic metabolism (TANM), but does not exceed it. In the third and fourth zones of muscle activity intensity, anaerobic energy supply processes predominate to a greater extent. The fifth rowing intensity zone involves training only the phosphate mechanism, the substrate of which is adenosine triphosphate (ATP).

The figure 1 shows the approximate dynamics of the volume of basic training tools for training canoe slalom athletes of the national team of Ukraine. The analysis of training programs showed that at the beginning of the preparatory period (October-November) the rowing volume in aerobic, aerobic-anaerobic and anaerobic modes gradually increases.

Figure 1. Dynamics of rowing volume and intensity in the annual training cycle

In December, the amount of training load in the aerobic and aerobic-anaerobic zones significantly decreases and its intensity increases in parallel. The volume of rowing in anaerobic mode increases by more than 2 times. Such dynamics of the training load corresponds to the general preparatory stage. In January, athletes do not use rowing due to the lack of natural conditions for training. In February, a special preparatory period begins in the national team. Gradually, the rowing volume in aerobic mode increases until April, then there is a certain decline, and the maximum load falls on June, that is, when the competitive stage begins. In June, there is a maximum rowing volume in the aerobic-anaerobic mode (an average of 340 km). However, the rowing volume in anaerobic mode in April - May decreases on average from 75 km to 60 km, and in June - July to 50 km per month or less. Such a construction of the training process corresponds to the traditional principle of undulation from the theory of periodization of sports training L. P. Matveev 2010.

In the course of the study, it was revealed that the total rowing volume in various intensity zones was on average 5115.26 ± 283.49 km (100%). As for the ratio of volumes in different zones of muscle work intensity, the rowing volume in the zone of moderate power averaged 2383.36 ± 142.74 km. With such rowing intensity, when the concentration of lactate does not exceed 2 mmol / l, the energy supply of muscle activity, that is, physical work, is carried out exclusively due to the aerobic mechanism. Under such conditions, the heart rate (HR) of athletes is within 70–80% of the maximum, and the intensity of the training load corresponds to 80–90% of the level of the anaerobic threshold (ANT). It should be noted that the share of rowing in this power zone is quite large and averages 46.6% of the total rowing volume.

It was determined that the rowing volume in the submaximal power-1 zone, when the boat's moving speed is within 70–80% of the maximum, is on average 1931.52 ± 89.56 km (37.7% of the total volume). With such rowing intensity, the lactate concentration is in the range of 2–4 mmol / l, heart rate is at the level of 80–90% of the maximum, and the intensity of the training load corresponds to 90–100% of the anaerobic threshold level (ANT). That is, the energy supply of muscle activity in this case is also carried out mainly due to the aerobic mechanism.

The annual rowing volume in the submaximal power-2 zone of the national team rowers is significantly reduced, compared to the previous two, and is approximately 480.71 ± 48.26 km, or 9.4% of the total volume of rowing in the water. Rowing intensity exceeds ANT by 10%, boat speed reaches 80–85% of maximum. The athletes' heart rate rises to 95% of the maximum value, and muscle activity at this intensity is provided not only by the aerobic, but also by the anaerobic mechanism, therefore the lactate concentration increases from 4 mmol / l to 8 mmol / l.

Rowing volume in a large power zone is an average of 229.48 ± 37.72 km, or 4.5% of the total mileage. In this case, the speed of the boat reaches 90–100% of the maximum, and the concentration of lactate exceeds 8 mmol / l.

The smallest rowing volume of the examined athletes falls on the maximum intensity zone. The speed of the boat in distance reaches its maximum. Such work takes an average of 92.63 ± 16.39 km per year (1.8%). This is usually rowing on segments from 20 to 100 meters.

Considering the annual amount of means of general physical preparation (GPP) for kayaking and canoeing, one can also draw a definite conclusion. So, the annual volume of GPP in them averages 191.25 ± 18.57 hours. Of these, an average of 70.67 ± 8.58 hours, or 36.6% of the total number of hours, is devoted to training. Athletic training takes the second place in the physical preparatio structure, which is given on average - 60.28 ± 6.36 hours, or 31.4%. In third place in the annual cycle is cross-country running. It amounts to an average of 31.27 ± 4.51 hours, or 16.3% of the total amount of funds for the general physical preparation. The least time is given for general developmental exercises - 30.05 ± 4.38 hours (15,7 %).

Dicussion

Thus, we can conclude that the construction of the annual training cycle of canoe slalom athletes of the main team of Ukraine is only partially based on the ideas of L.P. Matveev 2010, Garcia-Pallares, J., 2010, Macdermid, P. W., and Fink, P. W. 2017. This mainly concerns the construction of a general preparatory phase. At the beginning of the preparatory period (October), rowers perform a large volume of rowing in aerobic (294.0 km) and aerobic-anaerobic modes (162.0 km). In November - December, the rowing volume in the aerobic zone is reduced by more than 2 times, while the rowing volume in the aerobic-anaerobic and anaerobic modes increases. That is, in its structure, the training process of rowers of the basic composition of the national team of the Ukraine corresponds to the classical concepts of constructing the general preparatory stage of athletes according to the model of the annual cycle Messias, L. H. D., et al. 2015, Bonetti, D.L., & Hopkins, W.G. 2010, Okun D., Mulyk K. 2017.

In January, adult athletes were engaged only in general physical training. At the beginning of the specially preparatory stage (February - March), the rowing volume naturally increases in the aerobic-anaerobic mode. Moreover, it is more than the volume of rowing in aerobic mode. Then, the athletes gradually increase the volume of rowing in anaerobic mode, reaching a maximum by the beginning of the competitive stage (June). In June, the rowing volume in the aerobic regime also increases sharply. This situation is not typical for the competitive stage. However, an analysis of the performance of Ukrainian rowers at the World Championships on August 19–23, 2017 in Italy (Milan) (Bielik, V., Lendvorský, L., Lengvarský, L., Lopata, P., Petriska, R., & PelikáNová, J. 2018), where 4 medals were won (1 silver and 3 bronze) showed that the optimal model for the volume and intensity of the load was chosen in the annual training cycle.

The analysis of the structure and content of the annual training cycle of the main national team of Ukraine in canoe slalom showed that the construction of the training process requires a constant search and adjustment of the volume and intensity of the load. Such an approach can significantly increase the efficiency of the training process of rowers of the national team of Ukraine, which will contribute to the growth of their sportsmanship and competitive performance.

Conclusions

The analysis of the structure and content of the annual training cycle of the national team of Ukraine in canoe slalom showed that the construction of the training process at the specially preparatory and competitive stages requires a serious adjustment of the distribution of the volumes and intensity of the main means of training, taking into account the scientifically based patterns of development sport preparedness. This approach can significantly increase the effectiveness of the training process of rowers, which will contribute to the growth of their sportsmanship and competitive performance.

The total mileage of water rowing averages 5115 km. Of these: rowing in the moderate power zone - 2383 km (46.6%); rowing of submaximal power - 2419 km (47.1%); in the zone of high power - 229 km (4.5%); in the zone of maximum power - 92 km (1.8%).

The optimal amount of general physical fitness equipment used by kayakers and canoes in an annual training cycle is an average of 191 hours. Of these, 70 hours (36.6%) - training, 60 hours (31.4%) - athletic training, 31 hours (16.3%) - cross-country running and 30 hours (15.7%) - general developmental exercises;

The created model of the volume and intensity of training loads can be considered basic in building the training process of highly qualified canoe slalom athletes.

Conflict of interests. The authors declare that no conflict of interest, which can be perceived so that it can harm the impartiality of the article.

Financing sources. This article didn't get the financial support from the state, public or commercial organization.

References:

- Bielik, V., Lendvorský, L., Lengvarský, L., Lopata, P., Petriska, R., & PelikáNová, J. (2018). Road to the Olympics: Physical fitness of medalists of the canoe Sprint Junior European and World championship events over the past 20 years. *Journal of Sports Medicine and Physical Fitness*, 58(6), 768-777
- Bonetti, D.L., & Hopkins, W.G. (2010). Variation in performance times of elite flat-water canoeists from race to race. *International Journal of Sports Physiology and Performance*, 5(2), 210-217.
- Braghin, F., Cheli, F., Melzi, S. & Sabbioni, E. (2008). Race driver model. *Computers and structures*. 86, (13-14), 1503-1516. Название журнала
- Brooks, G. A. (2018). The Science and Translation of Lactate Shuttle Theory. *Cell metabolism*, 27(4), 757–785. United States. <https://doi.org/10.1016/j.cmet.2018.03.008>
- Ferrari, H. G., Messias, L. H. D., Reis, I. G. M., Gobatto, C. A., Sousa, F. A. B., Serra, C. C. S., & Manchado-Gobatto, F. B. (2017). Aerobic Evaluation in Elite Slalom Kayakers Using a Tethered Canoe System: A New Proposal. *International journal of sports physiology and performance*, 12(7), 864–871. United States. <https://doi.org/10.1123/ijsp.2016-0272>
- Garcia-Pallares, J., Garcia-Fernandez, M., Sanchez-Medina, L., & Izquierdo, M. (2010). Performance changes in

- world-class kayakers following two different training periodization models. *European journal of applied physiology*, 110(1), 99–107. Germany. <https://doi.org/10.1007/s00421-010-1484-9>
- Harwood, C.G., & Knight, C.J. (2015). Parenting in youth sport: A position paper on parenting expertise. *Psychology of Sport and Exercise*, 16, 24–35. doi: 10.1016/j.psychsport.2014.03.001
- Holt, N.L., Tamminen, K.A., Black, D.E., Mandigo, J.L., & Fox, K.R. (2009). Youth sport parenting styles and practices. *Journal of Sport & Exercise Psychology*, 31, 37–59.
- Hunter, A. (2009). Canoe slalom boat trajectory while negotiating an upstream gate. *Sports Biomechanics*, 8(2), 105–113.
- Hunter, A., Cochrane, J. & Sachlikidis, A. (2007). Canoe slalom -competition analysis reliability. *Sports Biomechanics*, 6(2), 155–170
- Macdermid, P. W., and Fink, P. W. (2017). The validation of a paddle power meter for slalom kayaking. *Sports Med. Int. Open* 1, E50–E57. doi: 10.1055/s-0043-100380
- Manchado-Gobatto, F. B., Arnosti Vieira, N., DalchecoMessias, L. H., Ferrari, H. G., Borin, J. P., de Carvalho Andrade, V., & Terezani, D. R. (2014). Anaerobic threshold and critical velocity parameters determined by specific tests of canoe slalom: Effects of monitored training. *Science & Sports*, 29(4), e55–e58. <https://doi.org/10.1016/j.scispo.2014.04.006>
- Matveev, L.P. (1999). *Fundamentals of the general theory of sports and the athletes' training system*. Moskov: OlympicLiterature.
- Matveev, L.P. (2010). *General theory of sports and its applied aspects: a textbook for universities of physical education*. Moskov: OlympicLiterature.
- Messias, L. H. D., De Barros Sousa, F. A., Dos Reis, I. G. M., Ferrari, H. G., Gobatto, C. A., Serra, C. C. S., et al. (2018). Novel paddle stroke analysis for elite slalom kayakers: relationship with force parameters. *PLoS One* 13:e0192835. doi: 10.1371/journal.pone.0192835
- Messias, L. H. D., dos Reis, I. G. M., Ferrari, H. G., & de Barros Manchado-Gobatto, F. (2014). Physiological, psychological and biomechanical parameters applied in canoe slalom training: a review. *International Journal of Performance Analysis in Sport*, 14(1), 24–41. Routledge. <https://doi.org/10.1080/24748668.2014.11868700>
- Messias, L. H. D., Ferrari, H. G., Reis, I. G. M., Scariot, P. P. M., & Manchado-Gobatto, F. B. (2015). Critical Velocity and Anaerobic Paddling Capacity Determined by Different Mathematical Models and Number of Predictive Trials in Canoe Slalom. *Journal of Sports Science & Medicine*, 14(1), 188–193.
- Messias, L. H. D., Sousa, F. A. de B., Dos Reis, I. G. M., Ferrari, H. G., Gobatto, C. A., Serra, C. C. S., Papoti, M., et al. (2018). Novel paddle stroke analysis for elite slalom kayakers: Relationship with force parameters. *PLoS one*, 13(2), e0192835. <https://doi.org/10.1371/journal.pone.0192835>
- Messias, L. H., Ferrari, H. G., Reis, I. G., Scariot, P. P., and Manchado-Gobatto, F. B. (2015). Critical velocity and anaerobic paddling capacity determined by different mathematical models and number of predictive trials in canoe slalom. *J. Sports Sci. Med*, 14, 188–193.
- Messias, L.H.D, Masselli dos Reis, I.G., Ferrari, H.G. & Manchado-Gobatto, F. (2014) Physiological, psychological and biomechanical parameters applied in canoe slalom training: a review. *International Journal of Performance Analysis in Sport*, 14, 24–41.
- Okun D., Mulyk K. (2017) Investigation of the relationship between the indicators of physical preparedness and the basic *technique* elements of young water-slalom athletes. *Slobozhanskyi herald of science and sport*, 2017, No5(61), pp. 69-71
- Okun, D. (2016) Comparative analysis of special preparedness young water-slalom. *Slobozhanskyi herald of science and sport*, 2016, No 1(51), pp. 38-40
- Sigmund, M., Rozsypal, R., Kudlacek, M., Kratochvil, J., & Sigmundová, D. (2016). Influence of one-year sport activities on the changes in morphological parameters and somatotypes in the current junior members of the Czech national whitewater slalom team. *Journal of Physical Education and Sport*, 16(1), 118.
- Zeimers, G., Anagnostopoulos, C., Zintz, T., & Willem, A. (2019). Organisational learning for corporate social responsibility in sport organisations. *European Sport Management Quarterly*, 19(1), 80–101.