EXPERIMENTAL STUDIES ON ACTIVE METABOLITES OF *CALOTROPIS GIGANTEA* FOR EVALUATION OF POSSIBLE ANTIFUNGAL PROPERTIES

Mandepudi David^{1,2*}, RavuruBharath Kumar³, Mandepudi Bhavani²

¹Faculty of Biotechnology, Jawaharlal Nehru Technological University-Hyderabad, Hyderabad, Andhra Pradesh, India

²Department of Biotechnology, School of Engineering, Sir Padampat Singhania University, Udaipur-313601, Rajasthan, India

³School of Biotechnology, Vignan University, Guntur – 522213, Andhra Pradesh, India

*Corresponding Author Email: david.mandepudi@gmail.com

Article Received on: 18/03/13 Revised on: 01/04/13 Approved for publication: 11/05/13

DOI: 10.7897/2230-8407.04553

IRJP is an official publication of Moksha Publishing House. Website: www.mokshaph.com

© All rights reserved.

ABSTRACT

The active metabolites like phytochemicals from the medicinal plants were under exploration for the development of novel and biodegradable effective drugs as an alternative to the ineffective contemporary medicine. In the similar lines, the *Calotropis gigantea* leaves, buds and flowers were extracted for the active metabolites using ethanol, methanol, chloroform and aqueous solvents and then these solvent extracts of 100µl (25µg/ml) were tested separately for the in vitro antifungal activity against the test pathogenic fungi viz. *Aspergillus niger*, *Rhizoctonia oryzae-sativae*, *Fusarium solani* and *Trichoderma viride*. In the results ethanol buds extract has shown the highest growth reducing activity (67.82%) on *Fusarium solani* where the positive control, clotrimazole (10µg/ml) recorded 58% growth reduction. Similarly aqueous buds extract has shown ~66% growth reduction on *Rhizoctonia oryzae-sativae* where Clotrimazole, the positive control has shown the maximum growth reduction capability (~73%); however all the aqueous extracts exhibited the growth reduction capabilities on all the test fungi. On the contrary methanol leaves extract increased the growth of *Aspergillus niger* by 85%; and similarly chloroform leaves and buds extracts exhibited the growth supporting activity on *Rhizoctonia oryzae-sativae* (32%) and *Trichoderma viride* (34%) respectively. This study brings out the mixed results of *Calotropis gigantea* solvent extracts as antifungal drugs and hence makes a caution on the knowledgeable and judicious use of these extracts as a medicine as evident among the folklores. Further it is also advised to make advanced studies in the similar line for the effective utilization of this *Calotropis* species in either directions to serve the mankind.

Keywords: Antifungal, *Calotropis*, Solvent extracts, Active metabolites, Phytochemicals

INTRODUCTION

Plant based active metabolites are under intensive exploration around the world among the scientific community¹⁻⁵ to supplement the contemporary medicinal practices with the plant and evidence based drugs as they have minimal side effects and folklore reputation. In this pursuit several plant based drugs and remedies were reported and still there exists vast potential to identify and explore the possible medicine from the herbal plants. The secondary metabolites were the most drug like molecules targeted in several investigations and they were proved to be fruitful in developing the potential drugs⁶. However the efforts in this line were limited mostly to the safest herbal medicines in terms of handling and processing⁷⁻⁸, so there can also be an alternative approach to choose the other plant sources like *Calotropis* species to explore the potential antimicrobial active metabolites⁹⁻¹² that may provide the latent drugs for animals, plants and even humans¹³. This *Calotropis* species having two variants viz. *Calotropis procera* and *Calotropis gigantea*, known for its poisonous nature, were least explored milk weeds with lot of secondary active metabolites in the form of latex in leaves, buds and to some extent in the flowers. Among these two variants *Calotropis procera*¹⁴⁻²⁰ has been explored to some extent for the medicinal properties but the *Calotropis gigantea* had the less exposure in the scientific investigations. So the present investigation reports the experimental studies to elucidate the possible antifungal properties of *Calotropis gigantea* aqueous and non-aqueous leaves, buds and flowers extracts on selected pathogenic fungi.

MATERIALS AND METHODS

Extracts of *Calotropis gigantea*

The solvent extracts of *Calotropis gigantea* leaves, buds & flowers were prepared with ethanol, methanol, chloroform

and aqueous solvents by soxhlet solvent extraction method and estimated the residual extract concentrations²¹. These solvent extracts of ethanol leaves (EL), ethanol buds (EB), ethanol flowers (EF), methanol leaves (ML), methanol buds (MB), methanol flowers (MF), chloroform leaves (CL), chloroform buds (CB), chloroform flowers (CF), Aqueous leaves (AL), Aqueous buds (AB) and Aqueous flowers (AF) were stored in the refrigerator until use.

Test Fungal Cultures

The fungal cultures *Rhizoctonia oryzae-sativae* (MTCC 2162), *Aspergillus niger* (MTCC 1344), *Fusarium solani* (MTCC 2671), and *Trichoderma viride* (MTCC 793) were procured from Microbial Type Culture Collection and Gene Bank (MTCC), Chandigarh, India and used them as the test fungi for the evaluation of *Calotropis gigantea* solvent extracts. All these fungal cultures were periodically subculture and maintained in the Microbiology laboratory, Department of Biotechnology, School of Engineering, Sir Padampat Singhania University, Udaipur, Rajasthan, India.

Inoculum and Media Preparation

The fungal spores were dispersed aseptically in 1% sterile saline solution to get the spore inoculum of 10⁵cfu/ml and used it as inoculum for the respective fungal cultures, in triplicates. The potato dextrose (PD) broth cultures inoculated with 100 µl of solvent extract with residual concentration of 25µg/ml from each sample separately except the control and the positive control (broth media added with 100µl of 10µg/ml clotrimazole, as the standard antifungal drug) with respect to each test fungus were sterilized by autoclaving at 121⁰C for 15min. and subsequently used these prepared sterile PD media for the tests.

Figure 1: Effects of *Calotropis gigantea* methanol extracts on the growth and survival of *Rhizoctonia oryzae-sativae* (RZ), *Fusarium solani* (FS), *Aspergillus niger* (AN) and *Trichoderma viride* (TV)

Figure 2: Effects of *Calotropis gigantea* ethanol extracts on the growth and survival of *Rhizoctonia oryzae-sativae* (RZ), *Fusarium solani* (FS), *Aspergillus niger* (AN) and *Trichoderma viride* (TV)

Figure 3: Effects of *Calotropis gigantea* chloroform extracts on the growth and survival of *Rhizoctonia oryzae-sativae* (RZ), *Fusarium solani* (FS), *Aspergillus niger* (AN) and *Trichoderma viride* (TV)

Figure 4: Effects of *Calotropis gigantea* aqueous extracts on the growth and survival of *Rhizoctonia oryzae-sativae* (RZ), *Fusarium solani* (FS), *Aspergillus niger* (AN) and *Trichoderma viride* (TV)

Broth Culturing of Test Fungi

The prepared sterile PD broth media were inoculated with 100µl of the respective prepared fungal spore suspension (10^5 cfu/ml) aseptically including the positive controls and negative controls. These spore inoculated media were incubated in an orbital shaker incubator at 120rpm, 30°C for 48 hrs.

Estimation of Biomass

The fungal biomass^{5,22} grown upto 48 hrs., were filtered through the pre-weighed Whatman filter papers separately and the residual biomass on the filter papers for each culture

were dried in the oven at 60°C for overnight and then the final dry weight was measured along with the filter paper. The differential fungal dry mass were calculated on the basis of control for each sample and used them to plot the growth reduction(%) profiles of *Calotropis gigantea* solvent extracts against the test pathogenic fungal organisms as shown in the Figures 1-4.

RESULTS

The effect of methanol extracts of *Calotropis gigantea* as shown in Fig-1, has brought out the buds extract as the highest growth reducing (67.82%) extract with respect to

Fusarium solani; otherwise also, the leaves extract as well as the flowers extract has the comparable growth reduction²³⁻²⁴ capacity (~52%) with that of the positive control, clotrimazole (58%). Similarly, the buds extract has recorded the second best growth reducing methanol extract (41.56%) on *Trichoderma viride*. But then both leaves and flowers extracts have shown mild growth reduction effects on *Trichoderma viride*. On the contrary, none of the methanol extracts, including controls, has positive growth reduction effects on *Aspergillus niger* and further they were found to support the growth and especially leaves extract has showed the increase in growth to about 85%. The ethanol extracts of *Calotropis gigantea* seem to be having almost no growth reduction capabilities (Fig. 2) on any of the test fungal organisms, owing to the fact that the methanol, negative control, has exhibited the maximum reduction capability on all the test fungi except *Fusarium solani*, where conversely ethanol flowers extracts exhibited the highest growth promoting capacity followed by ethanol buds and leaves extracts respectively. Similarly the ethanol buds extract has the growth promoting capability for soil born *Trichoderma viride*, whereas the ethanol leaves and flowers extracts has mild growth reducing capabilities on this fungus.

The chloroform extracts of *Calotropis gigantea*, like that of ethanol extracts, were found to have no growth reducing capabilities on these test fungi except *Fusarium solani* as shown in the Fig-3; where the chloroform leaves & flowers extracts have the slight growth reduction (~20%) capabilities on *Fusarium solani*, whereas the positive control clotrimazole had the 58.21% of growth reduction capacity. On the other hand the chloroform leaves and buds extracts have the growth promoting capabilities on *Rhizoctonia oryzae-sativae* (32%) and *Trichoderma viride* (34%) respectively.

The aqueous *Calotropis gigantea* extracts as shown in the Fig-4, exhibited growth reduction capabilities on all test fungi except *Trichoderma viride*, where only aqueous leaves extract has mild growth reduction capability (~5%) and other aqueous buds & flowers extracts supported the growth (~10%). Similarly, aqueous extracts have demonstrated (40-50) % of growth reduction on *Aspergillus niger*. Further these aqueous extracts have shown (20-30) % growth reduction on *Fusarium solani*²⁵ and (30-40) % growth reduction on *Rhizoctonia oryzae-sativae*.

DISCUSSION

The aqueous extracts of *Calotropis gigantea* produced growth reduction on *Rhizoctonia oryzae-sativae* (30-40)%, *Fusarium solani* (20-30)% and *Aspergillus niger* (40-50) %; which imply that *Calotropis gigantea* produces water soluble metabolites that could restrict the growth of fungus and protect itself from the decay. Further methanol extracts and aqueous extracts have demonstrated higher and moderate growth reduction potential on *Fusarium solani*, respectively; whereas chloroform extracts had mild growth reduction potential on it and on the contrary, ethanol extracts had growth supporting effects on *Fusarium solani*. Such kind of mixed response may be due to the proportionate distribution of respective metabolites for restriction and provision of plant based nutrients for these fungi. Similar results were also observed in the case of *Aspergillus niger*, whereas methanol buds & leaves extracts exhibited moderate & mild growth reduction potential on *Trichoderma viride*, but all other extracts favored the growth of *Trichoderma viride*. These results also suggest the proportionate distribution of

Calotropis gigantea active metabolites which may help to protect the plant during the vegetative or propagative phase and some metabolites may favor the growth of fungi during the degradation of wornout or shed plant materials²⁶. In addition, it may also be inferred that the support of *Calotropis gigantea* extracts for the growth of *Trichoderma viride* helps in reduction of soil born fungal diseases as *Trichoderma viride* was also found to act as a biological control against soil born fungal diseases²⁷⁻²⁸. Perhaps this was the reason for the use of *Calotropis gigantea* leaves as the green manure in the seedling bed to grow the seedlings and as manure for coconut plantations²⁹.

CONCLUSIONS

The studied *Calotropis gigantea* solvent extracts effects on the growth of pathogenic test fungal organisms revealed that, aqueous extracts of leaves, buds and flowers had the growth reduction potential except on *Trichoderma viride*, which may be the strategy of the plant to protect itself from these pathogenic fungi during the vegetative phase. These results are instrumental to understand the traditional practice of *Calotropis gigantea* aqueous extracts fresh leaves & latex preparations and their use as a medicine among many ethnic groups in the world. On the other hand the non-aqueous ethanol, methanol & chloroform extracts had the mixed distribution of the active metabolites from the *Calotropis gigantea* leaves, buds & flowers and hence necessitates the knowledge and caution in using these plant materials as part of the medicine and such knowledge has been with the traditional healers among the ethnic groups around the world. So, it would be of interest for scientific community and the society at large to have similar studies on the active metabolites in the extracts of *Calotropis* species to understand the traditional drug practices and to explore the possibilities of potential drugs for contemporary medical practice.

ACKNOWLEDGEMENT

Dr. M Lakshmi Narasu, Professor & Dr. Archana Giri, Assistant Professor, Faculty of Biotechnology, Jawaharlal Nehru Technological University, Hyderabad; Dr. Archana Gajbhiye, Head, Department of Biotechnology, Sir Padampat Singhania University, Udaipur were acknowledged for their suggestions and critical comments on the research work. Mr. Ashok Ghosh, President, Ms. Rinu Ghosh, Vice President, Prof. P.C. Deka, Vice Chancellor and Prof. Achintya Chaudary, Dean, School of Engineering of Sir Padampat Singhania University, Udaipur were also greatly acknowledged and thanked for their support and approvals for the research work.

REFERENCES

1. Koehn FE, Carter GT. The evolving role of natural products in drug discovery. Nat. Rev. Drug Discov., 2005; 4:206-220. <http://dx.doi.org/10.1038/nrd1657> PMID:15729362
2. Balunas MJ, Kinghorn AD. Drug discovery from medicinal plants. Life Sci., 2005; 78:431-441. <http://dx.doi.org/10.1016/j.lfs.2005.09.012> PMID:16198377
3. Anupam G, Bidus KD, Soroj KC, Goutam C. Antibacterial potentiality and phytochemical analysis of mature leaves of *Polyalthia longifolia* (Magnoliales: Annonaceae). The South Pacific Journal of Natural Science 2008; 26:68-72. <http://dx.doi.org/10.1071/SP08011>
4. Wakirwa JH, Ibrahim P, Madu SJ. Phytochemical screening and in vitro antimicrobial analysis of the ethanol stem bark extract of *Jatropha curcas* Linn. (Euphorbiaceae). International Research Journal of Pharmacy 2013; 4(3):97-100. <http://dx.doi.org/10.7897/2230-8407.04317>
5. Javid IP. Antifungal activity and phytochemical screening of plant species from northern India. World journal of pharmacy and pharmaceutical sciences 2012; 1(2):810-819.
6. Balandrin MF, Klocke JA, Wurtele ES, Bollinger WH. Natural plant chemicals: Sources of industrial and medicinal materials. Science 1985; 228:1154-60. <http://dx.doi.org/10.1126/science.3890182> PMID:3890182

7. Majaw S, Moirangthem J. Quantitative and qualitative analysis of *Clerodendron colebrookianum* Walp. Leaves and Zingiber cassumunar Roxb. Rhizomes. *Ethnobotanical Leaflets* 2009; 13:578-89.
8. Veronese P, Li X, Niu X, Weller SC, Bressan RA, Hasegawa PM. Bioengineering mint crop improvement. *Plant Cell, Tissue and Organ Culture* 2001; 64:133-144. <http://dx.doi.org/10.1023/A:1010649207445>
9. Cordell GA. Biodiversity and drug discovery: A symbiotic relationship. *Phytochemistry* 2000; 55:463-480. [http://dx.doi.org/10.1016/S0031-9422\(00\)00230-2](http://dx.doi.org/10.1016/S0031-9422(00)00230-2)
10. Maureen KK, Rana AC, Dixit VK. *Calotropis* species: A Comprehensive Review. *Phcog Mag Plant Rev.*, 2005; 1(2):48-52.
11. Sharma GK. *Calotropis* procera and *Calotropis gigantea*. *Indian journal of veterinary science* 1934; 4:63-74.
12. Mueen AK, Rana KAC, Dixit VK. Studies on *Calotropis* species for antimycocardial infections activity in albino rats. *Indian Drugs* 2002; 39(11):613-615.
13. Falguni KS, Parabiah MH. Ethnobotanical studies and validation of lead: a case study on evaluation of *Calotropis* sp. on dermal fungal infections. *International journal of pharmacy & life sciences* 2011; 2(6):797-800.
14. Nighat B, Sharma B, Pandey RS. Evaluation of insecticidal efficacy of *Calotropis* procera and *Annona squamosa* ethanol extracts against *Musca domestica*. *Journal of Biofertilizers & biopesticides* 2010; 1(1):1-6.
15. Shanti SK, Nagamalleswari SK, Chaitra D, Yasodamma N. Traditional use of *Calotropis* procera R. Br. against migraine. *Current science* 2011; 101(9):1120-1121.
16. Dewan S, Sangraula H, Kumar VL. Preliminary studies on the analgesic activity of latex of *Calotropis* procera. *Journal of Ethnopharmacology* 2000; 73:307-311. [http://dx.doi.org/10.1016/S0378-8741\(00\)00272-5](http://dx.doi.org/10.1016/S0378-8741(00)00272-5)
17. Kumar S, Dewan S, Sangraula H, Kumar VL. Anti-diarrhoeal activity of latex of *Calotropis* procera. *Journal of Ethnopharmacology* 2001; 76:115-118. [http://dx.doi.org/10.1016/S0378-8741\(01\)00219-7](http://dx.doi.org/10.1016/S0378-8741(01)00219-7)
18. Kumar VL, Shivka YM. In vivo and in vitro effect of latex of *Calotropis* procera on gastrointestinal smooth muscles. *Journal of Ethnopharmacology* 2004; 93:377-379. <http://dx.doi.org/10.1016/j.jep.2004.04.013> PMID:15234780
19. Roy S, Sehgal R, Padhy BM, Kumar VL. Antioxidant and protective effect of latex of *Calotropis* procera against alloxan-induced diabetes in rats. *Journal of Ethnopharmacology* 2005; 102:470-473. <http://dx.doi.org/10.1016/j.jep.2005.06.026> PMID:16054794
20. Rajesh R, Raghavendra CD, Gowda A, Nataraju BL, Dhananjaya K, Kemparaju K, Vishwanath BS. Procoagulant activity of *Calotropis gigantea* latex associated with fibrin(ogen)olytic activity. *Toxicon* 2005; 46(1):84-92. <http://dx.doi.org/10.1016/j.toxicon.2005.03.012> PMID:15922393
21. David M, Bharath KR, Bhavani M. Experimental investigation for pest resistant properties of *Calotropis gigantea*. *International Journal of Pharmaceutical Applications* 2012; 3(3):380-386.
22. Chandrasekaran M, Venkatesalu V. Antibacterial and antifungal activity of *Syzygium jambolanum* seeds. *J Ethnopharmacol.*, 2004; 91:105-108. <http://dx.doi.org/10.1016/j.jep.2003.12.012> PMID:15036477
23. Bashir U, Tahira JJ. Evaluation of *Eucalyptus camaldulensis* against *Fusarium solani*. *Int. J. Agric. Biol.*, 2012; 14:675-677.
24. Kirtikar KR, Basu BD. *Indian Medicinal Plants: Plates*. Vol. III, 2nd edn. Dehradun (India): Bishen Singh Mahendra Pal Singh Publishers and Distributors; 1999. p.191-92, 420-22, 993-94, 2045-47.
25. Abu-Taleb AM, Deeb KE, Al-Otibi FO. Bioactivity of some plant extracts against *Drechslera bisepata* and *Fusarium solani*. *Journal of Food* 2011; 9:769-774.
26. Jeyaseelan EC, Vinuja T, Pathmanathan K, Jeyadevan JP. Control of plant pathogenic fungi using organic solvent extracts of leaf, flower and fruit of *Lawsonia inermis* L. *International Journal of Pharmaceutical & Biological Archives* 2012; 3(4):783-788.
27. Brown HL, Bruce A. Assessment of the biocontrol potential of a *Trichoderma viride* isolate part I: Establishment of field and fungal cellar trials. *International Biodeterioration and Biodegradation* 1999; 44:219-223. [http://dx.doi.org/10.1016/S0964-8305\(99\)00082-7](http://dx.doi.org/10.1016/S0964-8305(99)00082-7)
28. Brown HL, Bruce A, Staines HJ. Assessment of the biocontrol potential of a *Trichoderma viride* isolate part II: protection against soft rot and basidiomycete decay. *International Biodeterioration and Biodegradation* 1999; 44:225-231. [http://dx.doi.org/10.1016/S0964-8305\(99\)00083-9](http://dx.doi.org/10.1016/S0964-8305(99)00083-9)
29. http://agritech.tnau.ac.in/itk/IndigenousTechKnowledge_Oper.html (Retrieved on 27th May 2013)

Cite this article as:

Mandepudi David, Ravuru Bharath Kumar, Mandepudi Bhavani. Experimental studies on active metabolites of *Calotropis gigantea* for evaluation of possible antifungal properties. *Int. Res. J. Pharm.* 2013; 4(5):250-254

Source of support: Nil, Conflict of interest: None Declared