Research Article

COMPARISON OF LIQUISOLID AND INCLUSION COMPLEXATION TECHNIQUE FOR DISSOLUTION RATE ENHANCEMENT OF VALSARTANHitesh Jain^{1*}, T. Y. Pasha², C. S. Bais¹, Anil Bhandari¹¹Jodhpur National University, Faculty of Pharmacy, Jodhpur, Rajasthan, India²Department of Chemistry, Parul Institute of Pharmacy and Research, Baroda, Gujarat, India

*Corresponding Author Email: hitesh_hitachi@rediffmail.com

Article Received on: 28/05/14 Revised on: 20/06/14 Approved for publication: 17/07/14

DOI: 10.7897/2230-8407.0507123**ABSTRACT**

The objective of this study is to screening of suitable non volatile liquid vehicle for the formulation of liquisolid tablets such as propylene glycol, polyethylene glycol 400 and tween 80 by using mathematical equations for enhancement of dissolution rate of drug and comparison of liquisolid technique with inclusion complex of β -cyclodextrin. Different liquisolid compacts were prepared using a mathematical model for calculating required quantities of powder and liquid ingredients to produce an acceptably flow able and compressible admixture. The prepared liquisolid systems were evaluated for their pre compression and post compression parameters. The drug-excipient interactions studies carried out by Infrared spectroscopy (IR) and X- ray diffraction (XRD). The maximum solubility of valsartan was found to be in propylene glycol. The liquisolid formulations containing propylene glycol as a non volatile solvent showed higher drug release. The IR studies confirmed that there was no interaction between the drug and excipients. The XRD analysis confirmed formation of a solid solution inside the compact matrix. All the pre compression and post compression properties of the liquisolid compacts were within the acceptable limits. Liquisolid technique can be used to improve the dissolution rate and bioavailability of low soluble drugs.

Keywords: Liquisolid compacts, Liquid medication, Mathematical model, Liquid load factor, β -cyclodextrin

INTRODUCTION

The BCS class II drugs for which the dissolution profile must be clearly defined and reproducible shows high absorption number (A_n) and low dissolution number (D_n). Drugs in this class are expected to have a variable dissolution profile due to the formulation and in vivo variables that, in turn, affect the absorption¹. The poor dissolution rate of such water-insoluble drug is a major impediment to the development of pharmaceutical dosage forms. The oral absorption of drugs is most often controlled by dissolution in the gastrointestinal tract. During the past few years many techniques have been developed such as drug micronization²⁻⁷ or nanosuspension,⁸⁻¹⁰ solid dispersions, co precipitation, lyophilization, micro encapsulation, use of pro drug, drug derivetization processes and inclusion of drug solutions into soft gelatin capsules to improve the solubility and bioavailability. Formulation of liquisolid compact (LS) is a novel "Powder Solution Technology" which makes use of liquid medications admixed with suitable carriers and coating materials and formulated into a moderately flowing, dry looking, non adherent and compressible powder forms have increased the drug dissolution rate profiles¹¹⁻¹⁴. Due to their significantly increased wetting properties and surface of drug available for dissolution, liquisolid compacts of water-insoluble substances may be expected to display enhanced drug release properties, and consequently, improved bioavailability¹⁵⁻¹⁸. Valsartan is an angiotensin II receptor antagonist used in the management of hypertension¹⁹⁻²¹. The drug is rapidly absorbed following oral administration with a bioavailability of about 23 %²². It is a weekly acidic drug that has tetrazole derivative containing acid and carboxylic acid groups with pKa values of 4.73 and 3.9 respectively. Peak plasma concentrations occur in 2 to 4 hours and its plasma half life is about 7.5 hours²³. Therefore, improvement in its solubility and dissolution rate may lead to enhancement in bioavailability.

MATERIAL AND METHODS

Valsartan was obtained as a gift sample from Alembic research center, Baroda. Crospovidon, Avicel pH 101, Aerosil 200, tween, polyethylene glycol (PEG), propylene glycol (PG), sodium hydroxide, potassium dihydrogen orthophosphate were purchased from S.D. Fine Chemicals Ltd. (Mumbai, India). All other chemicals, reagents and solutions used were of analytical grade.

Saturation solubility studies

To select the best non-volatile solvent to dissolve valsartan, solubility studies of valsartan were carried out in six different non-volatile solvents, i.e. PEG200, PEG400, tween 20, tween 40, tween 80 and propylene glycol. Saturated solutions were prepared by adding excess drug to the vehicles and shaking on the incubator shaker for 48 h at 25°C. Accurately weighed quantities of the filtered supernatants were further diluted with methanol and analyzed spectrophotometrically at 250 nm for their drug content. From these results, the solubility of valsartan in the respective liquid vehicle was calculated. Each experiment was carried out in triplicate²⁴.

Formulation of conventional tablet and Liquisolid compacts

Depending on the liquid vehicle and drug concentration in the liquid medication used, different liquid load factors²⁵ (L_f) ranging from 0.286 to 0.822 (w/w) were employed in liquisolid preparations as shown in Table 1. Calculated quantities of valsartan and non volatile solvents were accurately weighed and mixed at 30°C. The mixture was sonicated for 10 minutes to obtain a clear solution. Required quantities of carrier (Avicel PH 102) and coating materials (Aerosil 200) were incorporated to the admixture of drug vehicle and mixed completely. The optimum concentration of crospovidone was added to prepared liquisolid systems. The formulated powder was compressed into tablets using rotary

press. Conventional tablets of Valsartan were prepared by direct compression containing 40 mg drug with Avicel PH 102, Aerosil 200²⁶.

Preparation of Inclusion Complexes

The drug and β -cyclodextrin complexes (1:1, 1:2 and 1:3) were prepared by kneading technique as shown in Table 2. In this method required amount of drug and β -cyclodextrin (β CD) was taken and was transferred to a mortar pestle. The mixture was size reduced by continuous stirring with pestle. Water-ethanol mixture (3:1) ratio was added to the above physical mixture and continuously stirred until the slurry mass was formed. Slurry mass was collected and dried in a hot air oven for 2 h at 50°C, dried mass was collected and further dried in desiccators over silica gel for 24 h to remove all the excess residual solvents. The dried mass was collected and passed through 60# mesh, and was compressed with a tablet compression machine.²⁷⁻³⁰

Pre compression Studies of Liquisolid System

Flow Properties of Liquisolid System

The liquisolid formulations were evaluated for the following properties like angle of repose, bulk density, tapped density, Hausner's ratio, Carr's index.

Infrared Spectra Analysis

IR spectra of the liquisolid system were recorded by the KBr pellet method using the Fourier transform infrared spectrophotometer. A baseline correction was made using dried potassium bromide, and then the spectrum of pure valsartan and liquisolid systems were obtained.

X-Ray Powder Diffraction (XRD)

Polymorphic changes in the drug are important since they might affect the dissolution rate and bioavailability; therefore, it was necessary to study the polymorphic changes of valsartan in liquisolid compacts. XRD spectra of optimized sample was recorded using a high-power powder x-ray diffractometer (Ru-200B, Pune, India) with Cu as target at a scan speed of 4°/min. The samples were analyzed at a 2θ angle range of 2–45°. The operating voltage and current were 40 kV and 55 mA respectively³¹⁻³³.

Post compression Studies of Liquisolid Compacts

The prepared liquisolid compacts were evaluated for carrying out various tests such as a drug content, friability, weight variation, hardness and disintegration. All tests were carried out according to the USP compendia specifications^{34,35}.

In vitro Dissolution Studies

Dissolution studies of liquisolid tablets were carried out in USP Apparatus II (Paddle type) (Electro Lab). Tablets were placed in dissolution vessel containing 900 ml simulated gastric fluid maintained at $37 \pm 0.5^\circ\text{C}$ and stirred at 50 rpm. Aliquots of 5 ml were withdrawn at specified time intervals for 60 minutes and replaced with an equal volume of fresh dissolution medium. The samples were analyzed spectrophotometrically at 250 nm.

Comparison of the dissolution data of pure drug, DCT, LS4 and FC3

The comparison between the dissolution rate of drug was carried out between pure drug, DCT, LS compact (LS4), and inclusion complex (FC3) to determine the optimized formulation. The Figure 6 depicts the comparative drug

release of pure drug, DCT, and optimized formulation of LS4 and FC3, thus the increased dissolution rate was found to be for the LS4, this might be due to the highest solubility of valsartan drug in propylene glycol, and also may be due to the increased wettability of the drug molecules, and also the carrier coating material ratio.

Stability Study

After determining the drug content and release studies, the LS4 formulation was charged for the accelerated stability studies according to ICH guidelines ($40 \pm 2^\circ\text{C}$ and $75 \pm 5\%$ RH) for a period of 6 months in a stability chamber. The optimized formulation was placed in USP type-I flint vials and hermetically closed with bromobutyl rubber plugs and sealed with aluminum caps³⁶. The samples were withdrawn at 3 months and 6 months and evaluated for various test parameters and compared with initial results.

RESULT AND DISCUSSION

Vehicle selection

The solubility of valsartan was determined in a six different non volatile solvents as shown in Table 3. The drug solubility was found to be maximum in propylene glycol.

Flow Properties of the valsartan Liquisolid System

The flow properties of the liquisolid powder system are influenced by physical, mechanical, and environmental factors. The different flow properties were employed as shown in Table 4. All liquisolid formulations showed the acceptable flow properties due the presence of avicel PH 101 as a carrier material.

IR Spectra Analysis

The IR spectrum of valsartan showed its characteristic IR absorption peaks at 3437, 2964, 1936, 1729, 1602, 1471, 1205 and 759 cm^{-1} . The LS mixture containing different non volatile solvent showed the same characteristic peaks. These spectral observations indicated no interaction between the drug and excipients as shown in Figure 1 to 4

X-Ray Diffractometry (XRD)

The diffraction spectrum of pure valsartan showed that the drug was of crystalline in nature as demonstrated by numerous, distinct peaks at 2θ of 5.6° , 10.5° , 12.9° , 18.8° , 20.6° . The spectrum liquisolid system showed that some peaks of pure valsartan were absent and intensity of peaks was reduced as shown in Figure 5. The result indicates that the drug in complex form and liquisolid system is amorphous as compared to the pure drug. Hence, increased dissolution of the drug was observed.

Quality Control Studies

All the liquisolid formulations complied the Pharmacopoeial required specifications for the weight variation test, friability content and uniformity tests as shown in Table 5; the average hardness of liquisolid formulations ranging from 3.43 ± 0.15 to $6.26 \pm 0.11\text{ cm}^2$. Disintegration time was found to be in the range of 1.71 ± 0.121 to 5.07 ± 0.273 minutes.

In vitro Dissolution Studies

All the liquisolid formulations showed greater drug release in comparison with the pure drug, directly compressed tablet and tablets prepared from inclusion complexes due to more solubility in the non volatile solvents as shown in Figure 6 and 7

Table 1: Composition of different liquisolid compacts

Liquisolid system	Liquid vehicle	Excipient ratio ()	Liquid load Factor () #	Avicel pH 101 (Q) ‡ (mg)	Aerosil 200 (q) ^ (mg)	Unit dose weight (mg)
LS1	PG	5	0.822	0.243	0.048	0.348
LS2	PG	10	0.491	0.407	0.041	0.512
LS3	PG	15	0.380	0.526	0.035	0.631
LS4	PG	20	0.325	0.615	0.030	0.719
LS5	PEG	5	0.657	0.304	0.060	0.424
LS6	PEG	10	0.331	0.604	0.060	0.739
LS7	Tween 80	5	0.811	0.247	0.049	0.353
LS8	Tween 80	10	0.417	0.479	0.047	0.594
LS9	Tween 80	15	0.286	0.700	0.046	0.825

*An appropriate amount of liquid medication containing 40 mg of valsartan was incorporated in each tablet containing 5% of Crospovidon as super disintegrants; ‡Amount of carrier () material, ^ Amount of coating () material, #Liquid load factor is defined as = /

Table 2: Formulation of tablet from inclusion complexes

Formulation ^a (complex system)	Valsartan (mg)	β -CD (mg)	Lactose (mg)	Talc (mg)	Magnesium Stearate (mg)	Tablet Weight (mg)
FC1 (1:1)	40	40	196	6	3	300
FC2 (1:2)	40	80	156	6	3	300
FC3 (1:3)	40	120	116	6	3	300

^aAll formulations contain 5 % Crospovidone as super disintegrant, 2 % talc and 1 % magnesium stearate

Table 3: Solubility of valsartan in different solvents (Mean $\pm$ SD, n = 3)

Solvent	Solubility (% w/w)
Propylene glycol	113.33 $\pm$ 1.52
Tween 80	85 $\pm$ 2.65
Tween 40	73.66 $\pm$ 1.53
Tween 20	66.66 $\pm$ 2.08
PEG 400	69.33 $\pm$ 1.53
PEG 200	56.67 $\pm$ 3.21

Table 4: Pre compression studies of the liquisolid compacts (Mean $\pm$ SD, n = 3)

Formulation Code	Bulk Density (g/cm ³)	Tapped Density (g/cm ³)	Hausner's Ratio	Carr's Index (%)	Angle of repose
LS1	0.354 $\pm$ 0.004	0.410 $\pm$ 0.008	1.16 $\pm$ 0.022	13.88	32.50 $\pm$ 0.40
LS2	0.373 $\pm$ 0.011	0.423 $\pm$ 0.004	1.15 $\pm$ 0.030	13.56	30.65 $\pm$ 0.43
LS3	0.395 $\pm$ 0.004	0.449 $\pm$ 0.006	1.13 $\pm$ 0.023	12.09	36.87 $\pm$ 0.56
LS4	0.435 $\pm$ 0.017	0.477 $\pm$ 0.004	1.09 $\pm$ 0.056	08.73	24.75 $\pm$ 0.57
LS5	0.343 $\pm$ 0.005	0.421 $\pm$ 0.003	1.22 $\pm$ 0.026	18.57	33.56 $\pm$ 0.31
LS6	0.377 $\pm$ 0.005	0.431 $\pm$ 0.006	1.14 $\pm$ 0.032	12.37	33.83 $\pm$ 0.29
LS7	0.345 $\pm$ 0.003	0.417 $\pm$ 0.004	1.20 $\pm$ 0.002	14.24	38.87 $\pm$ 0.65
LS8	0.377 $\pm$ 0.004	0.431 $\pm$ 0.003	1.14 $\pm$ 0.023	12.66	39.99 $\pm$ 0.47
LS9	0.381 $\pm$ 0.002	0.434 $\pm$ 0.005	1.14 $\pm$ 0.019	12.27	38.91 $\pm$ 0.71

Table 5: Post compression studies of the liquisolid compacts (Mean $\pm$ SD, n = 3)

Formulation Code	Hardness (kg/cm ²)	Friability (%)	Disintegration time (min) n = 6	Content Uniformity (%)	Weight variation
LS1	3.77 $\pm$ 0.15	0.252	4.05 $\pm$ 0.105	100.25 $\pm$ 0.22	Pass
LS2	4.33 $\pm$ 0.31	0.201	3.28 $\pm$ 0.204	100.06 $\pm$ 0.08	Pass
LS3	5.43 $\pm$ 0.20	0.150	3.25 $\pm$ 0.243	99.51 $\pm$ 0.42	Pass
LS4	4.06 $\pm$ 0.11	0.110	1.71 $\pm$ 0.121	100.25 $\pm$ 0.34	Pass
LS5	4.20 $\pm$ 0.36	0.293	4.30 $\pm$ 0.329	98.23 $\pm$ 0.67	Pass
LS6	6.26 $\pm$ 0.11	0.139	3.73 $\pm$ 0.258	99.53 $\pm$ 0.43	Pass
LS7	3.43 $\pm$ 0.15	0.364	5.07 $\pm$ 0.273	97.21 $\pm$ 0.87	Pass
LS8	5.53 $\pm$ 0.30	0.183	4.37 $\pm$ 0.459	95.63 $\pm$ 0.60	Pass
LS9	5.70 $\pm$ 0.26	0.135	4.08 $\pm$ 0.366	97.41 $\pm$ 0.27	Pass

Table 6: Effect of aging on formulation LS4 (Mean $\pm$ SD, n = 3)

Test	Initial	40°C/75%RH	
		3 months	6 months
Hardness (kg/cm ²)	4.06 $\pm$ 0.11	4.20 $\pm$ 0.36	4.03 $\pm$ 0.15
Friability (%)	0.110	0.111	0.113
Disintegration time (min) n=3	1.71 $\pm$ 0.121	2.92 $\pm$ 0.264	2.70 $\pm$ 0.155
Drug content (%)	100.25 $\pm$ 0.34	100.18 $\pm$ 1.36	100.06 $\pm$ 0.087
% Drug Dissolved (60 min)	80.87	80.18	79.5

Figure 1: Spectral analysis of valsartan pure drug by FTIR

Figure 2: Spectral analysis of LS physical mixture of PG by FTIR

Figure 3: Spectral analysis of LS physical mixture of PEG by FTIR

Figure 4: Spectral analysis of LS physical mixture of Tween 80 by FTIR

Figure 5: X-ray diffraction pattern of valsartan and formulation LS4

Figure 6: Dissolution profile of valsartan liquisolid compacts

Figure 7: Comparative dissolution profile of pure drug, DCT, LS4 and FC3

CONCLUSION

Powder solution technology is one of the promising approaches to increase dissolution. XRD results demonstrated valsartan had almost entirely converted from crystalline to amorphous state. The liquisolid formulation LS4 showed highest dissolution rate when compared with pure drug, directly compressed tablet and formulation prepared by complexation. This was due to increase in wetting properties and surface area of drug available for dissolution media. Aging studies had no effect on dissolution profile.

REFERENCES

- Amidon GL, Lennernas H, Shah VP, et al. A theoretical basis for a biopharmaceutical classification system: The correlation of *in vitro* drug product dissolution and *in vivo* bioavailability. *Pharmaceutical Research* 1995; 12: 413-420. <http://dx.doi.org/10.1023/A:1016212804288>
- Rasenack N, Hartenhauer HB, Muller W. Micro crystals for dissolution rate enhancement of poorly water-soluble drugs. *International Journal of Pharmaceutics* 2003; 254: 137-145. [http://dx.doi.org/10.1016/S0378-5173\(03\)00005-X](http://dx.doi.org/10.1016/S0378-5173(03)00005-X)
- Rogers TL, Johnston KP, Williams RO. Micronized powders of a poorly water-soluble drug produced by a spray-freezing into liquid emulsion process. *European Journal of Pharmaceutics and Bio Pharmaceutics* 2003; 55: 161-172. [http://dx.doi.org/10.1016/S0939-6411\(02\)00193-5](http://dx.doi.org/10.1016/S0939-6411(02)00193-5)
- Kayrak D, Akman U, Hortaesu O. Micronization of ibuprofen by RESS. *Journal Supercritical Fluids* 2002; 26: 17-31. [http://dx.doi.org/10.1016/S0896-8446\(02\)00248-6](http://dx.doi.org/10.1016/S0896-8446(02)00248-6)
- Sarkari M, Brown J, Chen XX, Swinnea SR, Williams O, Johnston KP. Enhanced drug dissolution using evaporative precipitation into aqueous solution. *International Journal of Pharmaceutics* 2002; 243: 17-31. [http://dx.doi.org/10.1016/S0378-5173\(02\)00072-8](http://dx.doi.org/10.1016/S0378-5173(02)00072-8)
- Ticehurst MD, Basford PA, Dallman CI, Lukas TM, Marshall PV, Nichols G, et al. Characterization of the influence of micronization on the crystallinity and physical stability of revatropate hydrobromide. *International Journal of Pharmaceutics* 2000; 193: 247-259. [http://dx.doi.org/10.1016/S0378-5173\(99\)00347-6](http://dx.doi.org/10.1016/S0378-5173(99)00347-6)
- Feely JC, York P, Sumbly BS. Determination of surface properties and flow characteristics of salbutamol sulphate, before and after micronization. *International Journal of Pharmaceutics* 1998; 172: 89-96. [http://dx.doi.org/10.1016/S0378-5173\(98\)00179-3](http://dx.doi.org/10.1016/S0378-5173(98)00179-3)
- Elamin AA, Sebhatu T, Ahlneck C. The use of amorphous model substances to study mechanically activated materials in the solid state. *International Journal of Pharmaceutics* 1995; 199: 25-36. [http://dx.doi.org/10.1016/S0378-5173\(94\)00364-B](http://dx.doi.org/10.1016/S0378-5173(94)00364-B)
- Chaumcil JC. Micronization: A method for improving the bioavailability of poorly soluble drugs. *Experimental and Clinical Pharmacology* 1998; 20: 211-215.
- Parrott EL, Swarbrick J. *Encyclopedia of Pharmaceutical Technology*, 4th ed. Marcel Decker, CRS Press; 1995.
- Yalkowsky S, Rubino JT. Solubilization by cosolvents I: Organic solutes in propylene glycol-water mixtures. *Journal of Pharmaceutical Sciences* 1985; 74(4): 416. <http://dx.doi.org/10.1002/jps.2600740410>
- Sugawara M, Kadomura S, He X, Takekuma Y, Kohri N, Miyazaki K. The use of an *in vitro* dissolution and absorption system to evaluate oral absorption of two weak bases in pH-independent controlled-release formulations. *European Journal of Pharmaceutics and Bio pharmaceutics* 2005; 26(1): 1-8.
- Youn YS, Jung JY, Oh SH, Yoo SD, Lee KC. Improved intestinal delivery of salmon calcitonin by Lys 18-amine specific PEGylation: Stability, permeability, pharmacokinetic behavior and *in vivo* hypocalcemic efficacy. *Journal of Controlled Release* 2006; 114: 334-342. <http://dx.doi.org/10.1016/j.jconrel.2006.06.007>
- Spireas S, Sadu S, Grover R. *In vitro* release evaluation of hydrocortisone liquisolid tablets. *Journal of Pharmaceutical Sciences* 1998; 87(7): 867-872. <http://dx.doi.org/10.1021/js970346g>
- Spireas S, Wang T, Grover R. Effect of powder substrate on the dissolution properties of methcrotiazide liquisolid compacts. *Drug Delivery and Industrial Pharmacy* 1999; 25(2): 163-168. <http://dx.doi.org/10.1081/DDC-100102156>
- Nokhodchi A. The effect of type and concentration of vehicles on the dissolution rate of a poorly soluble drug from liquisolid compacts. *Journal of Pharmaceutical Sciences* 2005; 8(1): 18-25.
- Javadzadeh Y, Siah Shadbad MR, Barzegar Jalali M, Nokhodchi A. Enhancement of dissolution rate of piroxicam using liquisolid compacts. *IL Farmaco* 2005; 60: 361-365. <http://dx.doi.org/10.1016/j.farmac.2004.09.005>
- Spireas S, Sadu S. Enhancement of prednisolone dissolution properties using liquisolid compacts. *International Journal of Pharmaceutics* 1998; 166: 177-188. [http://dx.doi.org/10.1016/S0378-5173\(98\)00046-5](http://dx.doi.org/10.1016/S0378-5173(98)00046-5)
- Leidig M, Bambauer R, Kirchert EJ, Szabà T, Handrock R, Leinung D, et al. Efficacy, safety and tolerability of valsartan 80 mg compared to irbesartan 150 mg in hypertensive patients on long-term hemodialysis. *Clinical Nephrology* 2008; 69(6): 425-32. <http://dx.doi.org/10.5414/CNP69425>
- Baumhäkel M, Müller U, Böhm M. Valsartan improves symptoms and quality of life in patients with chronic heart failure. *MMW Fortschritte der Medizin* 2008; 150(1): 48-53.
- Flynn JT, Meyers KE, Neto JP, De Paula R, Zurowska A, Bagga A, et al. Efficacy and safety of the angiotensin receptor blocker valsartan in children with hypertension aged 1 to 5 years. *Hypertension* 2008; 52(2): 222-8. <http://dx.doi.org/10.1161/HYPERTENSIONAHA.108.111054>
- Kshirsagar SJ, Bhalekar MR, Madgulkar AR, Sable PN, Gupta SR. Dissolution improvement of poor water soluble drug valsartan and improving flow properties of solid dispersion. *Latin American Journal of Pharmacy* 2010; 29: 393-400.
- Goodman LS, Gilman A. *The pharmacological basis of therapeutics*. 10th ed. New-York; 2001.
- Gubbi S, Ravindra J. Formulation and characterization of atorvastatin calcium liquisolid compacts. *Asian Journal of Pharmaceutics* 2010; 5(2): 50-60.
- Fahmy R, Kassem M. Enhancement of famotidine dissolution rate through liquisolid tablets formulation: *in vitro* and *in vivo* evaluation. *European Journal of Pharmaceutics and Bio pharmaceutics* 2008; 69: 993-1003. <http://dx.doi.org/10.1016/j.ejpb.2008.02.017>
- Louis D, Tayel S, Soliman I. Improvement of dissolution properties of Carbamazepine through application of the liquisolid tablet technique. *European Journal of Pharmaceutics and Bio pharmaceutics* 2008; 69: 342-347. <http://dx.doi.org/10.1016/j.ejpb.2007.09.003>
- Ammara HO, Salama HA, Ghorab M, Mahmoud AA. Inclusion complexation of glimepiride in dimethyl- β -cyclodextrin. *Asian Journal of Pharmaceutics* 2007; 2(2): 44-55.
- Soniwala MM, Patel PR, Ansuri MS, Parikh RK, Gohel MC. Various approaches in dissolution enhancement of Rofecoxib. *Indian Journal of Pharmaceutics* 2005; 67: 61-63.
- Vij M, Garse H, Dand N, Kadam V, Hirlekar R. Effect of preparation method on complexation of cefdinir with β -cyclodextrin. *Journal of Inclusion Phenomena and Macrocyclic Chemistry* 2009; 11: 1-9.
- Magnúsdóttir A, Másson M, Loftsson T. Cyclodextrins. *Journal of Inclusion Phenomena and Macrocyclic Chemistry* 2002; 44: 213-218. <http://dx.doi.org/10.1023/A:1023079322024>
- Fiese EF, Hagen TY. *Pre formulation In: Lachman, L, Lieberman HA. The theory and practice of Industrial Pharmacy*. 3rded. Varghese Publishing House, New Delhi; 2009. p. 182.
- Ansel HC, Allen LV, Popovich NG. *Pharmaceutical dosage forms and drug delivery systems*, Lippincott Williams and Wilkins, Philadelphia; 1999.
- Brittain HG. *Physical Characterization of Pharmaceutical Solids*, 4thed. Marcel Decker, CRS Press; 1995. <http://dx.doi.org/10.1201/b14204>
- Yunxia B, Sunada H, Yonezawa Y, Danjo K. Evaluation of rapidly disintegrating tablets prepared by direct compression method. *Drug Delivery and Industrial Pharmacy*; 1999. p. 571- 581.
- Zhao N, Augsburger L. Functionality comparison of 3 classes of super disintegrants in promoting aspirin tablet disintegration and dissolution. *American Association of Pharmaceutical Scientists*; 2005. p. 79.
- Venkatesh GS, Averineni R, Usha Y, Armugam K. Enhanced Dissolution and Bioavailability of Glucilazide Using Solid Dispersion Techniques. *International Journal of Drug Delivery* 2010; 49-57.

Cite this article as:

Hitesh Jain, T. Y. Pasha, C. S. Bais, Anil Bhandari. Comparison of liquisolid and inclusion complexation technique for dissolution rate enhancement of valsartan. *Int. Res. J. Pharm.* 2014; 5(7):604-609 <http://dx.doi.org/10.7897/2230-8407.0507123>