Review Article

THE EFFECTS OF *HYPERICUM* (HYPERICACEA) SPECIES ON MICROORGANISMS: A REVIEW

Yasemin Akgöz*

Department of Biology, Science Faculty, Dicle University, 21280 Diyarbakır/Turkey

*Corresponding Author Email: ysmnak@hotmail.com

Article Received on: 19/04/15 Revised on: 07/05/15 Approved for publication: 20/06/15

DOI: 10.7897/2230-8407.06781

ABSTRACT

Hypericum L. is a medicinal plant which is commonly found on earth, has many varieties and the most frequently studied regarding its antimicrobial, antifungal and antiviral effects because of its biological and chemical content. The most recognizable genus of *Hypericum* is *Hypericum perforatum* (common St. John's Wort: SJW). The plants of that genus comprise several different chemical compounds. These compounds have been studied in many chemical studies due to their active ingredients. The antimicrobial activity of this group of species has also been reported in numerous studies. In this study, the effect of more than 100 species of *Hypericum* genus of bacteria, fungi and viruses has been identified.

Keywords: *Hypericum*; antimicrobial; antifungal; antiviral.

INTRODUCTION

Hypericum L. genus, which is a member of *Guttifera* (Hypericaceae) family, contains 484 species that have been classified into 36 taxonomic sections in all over the continents except Antarctica¹. Members of the *Hypericum* genus are perennial, herbaceous or scrubby, and can be naturally found West Asia, Europe, Siberia, North America, Australia, Cyprus, North Africa, Madeira and the Azores and is naturalized in many parts of the world^{2,3}. The length of *Hypericum*, whose distribution can reach 2400 m in dry, calcareous and stony soils, light forests, meadows, marshes and shorelines, rocky places, roadsides, and non-cultivated fields can be up to 60 cm typically range from 40 to 80 cm in height^{4,5}. From the time of the ancient Greeks down through the Middle Ages, the plant was considered to be imbued with magical powers and was used to ward off evil and protect against disease⁶. Recently there has been increasing interest in the genus *Hypericum*, because it is a source of a variety of compounds with different biological activities⁷. The most popular species of this genus, *Hypericum perforatum* (common St. John's Wort: SJW), is one of the aromatic and medicinal plants which is widely used in many different treatments for various diseases^{8,10}. The commercially available *H. perforatum* derived products include phytopharmaceuticals and nutraceuticals, teas, tinctures, juices, and oily macerates⁹.

Chemical Properties of *Hypericum perforatum*

In the previous studies on this genus, the compounds isolated from SJW have been naphthodianthro derivatives (hypericin, pseudohypericin, protohypericin, and protopseudohypericin), flavonols (hyperoside, rutin, quercitrin, isoquercitrin, and quercetin) biflavons (biapigenin and amentoflavone), the phloroglucinol derivatives (hyperforin, adhyperforin), xanthenes, the phenolic acids (caffeic acid, chlorogenic acid, ferulic acid, hyperfolin, *p*-cumaric acid, *p*-hydroxybenzoic acid, vanillic acid), essential oils, sterols, beta-sitosterol, vitamin C, and vitamin A^{2,11-14}. In addition to these compounds, imanine and novoiimanine compounds which have antimicrobial activity have also been identified in SJW¹⁵. Nowadays, it is widely known that many of these compounds possess antibacterial activity against some pathogen

microorganisms. Apart from the compounds mentioned above, many other biologically active compounds from the same family have been isolated and acyl phloroglucinol particles, which have antimicrobial activity, have been identified^{16,17}.

Biological Effects of *Hypericum* Species

In phytochemical studies, compounds of *Hypericum* demonstrated antimicrobial¹⁸, anticancer¹⁹, antidepressant²⁰, antiviral^{21,22}, antioxidant^{23,24}, cytotoxic¹⁸, antifungal and antimalarial²⁵ activities. Benzopyran of xanthenes and flavonoid derived from these compounds are known to have antimicrobial activity against various fungi and bacteria^{12,18}.

Medical Uses of *Hypericum* Species

Many kinds of diseases have been treated with herbal remedies since ancient times²⁶. During the second half of the 20th century, the acceptance of traditional medicine as an alternative form of health care and the development of microbial resistance to the classical antibiotics led researchers to investigate the antimicrobial activity of several medicinal plants²⁷. Herbal remedies are still being used extensively in many countries²⁶. *Hypericum* is a well-known plant in herbal medicine due to its different species which have therapeutic effect. SJW has been used for some medical treatment such as removing wounds and burns in skin, sciatica, shingles, eczema, urinary incontinence in children, menopausal disorders, premenstrual syndromes, central nervous system disorders, mild and moderate depression, nerve damages, fear, worry, anxiety, joint inflammation (arthritis), bladder irritation, common cold, diabetes mellitus, dyspepsia, and has been found to be quite effective particularly in wounds, buckling, and cramps^{8,9,28-30}. Some *Hypericum* species are still popular wound healing agents in the treatment of some infectious diseases^{17,31}. Nowadays, the use of the commercial extracts taken from this plant as antidepressants. Pharmaceutical companies, particularly in Europe, prepare standard formulations of this herb that are taken by millions of people. Worldwide annual sales of products made from SJW presently exceeded several billion dollars in the world³².

The aim of this study was to review important studies about *Hypericum* species. This comprehensive study is important for future studies.

Antiviral effect of *Hypericum* Species

A number of compounds extracted from various species of higher plants have shown antiviral activity. Examples included tannins, flavones, alkaloids, that displayed *in vitro* activity against numerous viruses³³. Many compounds have been isolated from plants of this family and have had their antiviral activity studied³⁴. Hypericin and pseudohypericin extracted from SJW and may have therapeutic benefits viral infection and retroviral diseases²⁸. Both *in vitro* and *in vivo* studies, hypericin significantly inhibited viruses such as HIV (Human immunodeficiency virus)³⁵, Influenza A³⁶, Human cytomegalovirus (HCMV)^{33,37}, the DNA-virus, i.e. *Herpes Simplex* (type 1 and type 2)^{33,34,37,38}. The viral battery consisted of one DNA-virus, i.e. *Herpes simplex* type 1 (HSV-1), and RNA-viruses, i.e. *Poliomyelitis*, *Coxsackie*, *Semliki forest*, *Measles* and *Vesicular stomatitis virus* (VSV)³⁰.

Table 1 shows published articles about antiviral effects of some *Hypericum* species on different types of viruses such as *Coxsackie B2 virus*, *Measles edmonston A*, *Poliomyelitis virus*, *Semliki forest virus*, *Vesicular stomatitis virus*, *Human cytomegalovirus* (HCMV), *Human immunodeficiency virus* (HIV1), *Herpes simplex virus* (HSV1), *Influenza A virus* (IAV), *Feline immunodeficiency virus* (FIV) and *Hepatitis C. H. revolutum* and *H. perforatum* are the most studied species of *Hypericum* for different virus analyses. The antiviral effect of *Hypericum* species, the most preferred virus was HSV, which has been analysed with ten different *Hypericum* species (*H. capitatum*, *H. connatum*, *H. hookerianum*, *H. kotschyianum*, *H. mysorensense*, *H. neurocalycinum*, *H. perforatum*, *H. revolutum*, *H. salsugineum*, *H. scabrum*) and it has been followed by HIV1 with four different *Hypericum* species (*H. capitatum*, *H. connatum*, *H. perforatum*, *H. scabrum*). Eight *Hypericum* species (*H. caprifoliatum*, *H. hookerianum*, *H. japonicum*, *H. kotschyianum*, *H. mysorensense*, *H. neurocalycinum*, *H. polyanthemum*, *H. salsugineum*) have been analysed on only one type of virus.

Antibacterial and Antifungal Effect of *Hypericum* Species

Antibiotic resistance has become a global concern. In recent years there is increasing incidence of multiple resistance in human pathogenic microorganisms, largely due to the indiscriminate use of commercial antimicrobial drugs commonly employed in the treatment of infectious diseases³⁹. Multidrug-resistant *Staphylococcus aureus* (MRSA) infections, particularly those caused by methicillin-resistant *S. aureus*, have been a major threat to public health in hospitals and the community in the past decade⁴⁰. This has forced scientist to search for new antimicrobial substances from various sources like medicinal plants^{26,41}. Food spoilage is one of the most important issues facing the food industry. In fact, food-borne illnesses are a global problem, even in developing and developed countries. Food spoilage or deterioration is predominantly caused by the growth of microorganisms⁴¹. In many studies, many pathogenic microorganisms such as *Escherichia coli*, *Staphylococcus* sp., *Listeria monocytogenes*, *Campylobacter jejuni*, *Candida* sp., *Zygosaccharomyces* sp., *Fusarium* sp., *Aspergillus* sp., *Rhizopus* sp., *Penicillium* sp. and *Salmonella* sp. have been reported as the main factor in food spoilage and foodborne illnesses (food poisoning).

Hypericum genus has been found to yield compounds having the antimicrobial activity. Inhibitorial effects of SJW on Gram-positive bacteria have also been observed and attributed to hyperforin, the main acylphloroglucinol isolated from this genus^{40,42}. Phloroglucinol derivatives, found frequently in the lipophilic fractions of several *Hypericum* species, have demonstrated

antibacterial activities against microorganisms such as *S. aureus*, *Bacillus cereus*, *B. subtilis* and *Nocardia gardenen*.

In this study, as shown in Table 2, Gram-positive bacteria such as *Bacillus* sp., *Clostridium perfringens*, *Corynebacterium* sp., *Listeria monocytogenes*, *Micrococcus* sp., *Mycobacterium* sp., *Staphylococcus* sp., *Nocardia*, *Sarcina lutea*, *Streptococcus* sp., *Enterococcus* sp. and Gram-negative bacteria such as *Acinetobacter baumannii*, *Aeromonas hydrophila*, *Agrobacterium tumefaciens*, *Alcaligenes faecalis*, *Azotobacter chroococcum*, *Bordetella bronchiseptica*, *Branhamella catarrhalis*, *Campylobacter* sp., *E. coli*, *Erwinia carotovora*, *Enterobacter* sp., *Helicobacter pylori*, *Klebsiella* sp., *Proteus vulgaris*, *Pseudomonas* sp., *Shigella flexneri*, *Serratia marcescens*, *Salmonella* sp. and *Yersinia enterocolitica* have been commonly used for determination of anti-microbial and anti-fungal effects of some *Hypericum* species.

Table 2 also shown that In 93 different species of *Hypericum* have antibacterial effects on 31 different bacteria and 91 different *Hypericum* species except *H. amblycalyx* and *H. olivieri* have anti-staphylococcal effect. The *Staphylococcus* species have been identified as the leading bacteria which are mostly studied in terms of their interaction with this genus. Subsequently, 64 different species of *Hypericum* have been studied on *E. coli*, which is the second widely researched bacteria and the third researched is *Pseudomonas*. However, in bacteria species like *Alcaligenes faecalis*, *Azotobacter chroococcum*, *Campylobacter* sp., *Nocardia* sp. and *Vibrio* sp. only one study has been carried out with relation to one species of *Hypericum*. At the end of these studies, it has been revealed that *H. scabrum* and *H. perforatum* have become the top species about the effect of antibacterial studies.

Today, most of these microorganisms is known to cause many diseases in the inside or the outside the body of all living things⁴³. The detection of antimicrobial activity of *H. perforatum* has led into further studies on different species of *Hypericum*. Extracts of *H. brasiliense*, *H. scabrum*, *H. hircinum*, *H. hookerianum*, *H. canariense*, *H. maculatum*, *H. mysorensense* and *H. patulum*, and the antimicrobial activities of pure compounds obtained from these were investigated. Many extracts have been reported to be active and many pure compounds have showed either weak activity or non-active properties^{44,45}.

Antimicrobial activities of four different *Hypericum* species and *Hypericum calycinum*'s extracts taken from leaves and petals towards eight different microorganisms have been studied, and this study has proved that *H. calycinum*'s acetone extract has the highest rate of antimicrobial activity on *Staphylococcus aureus* and *Mycobacterium smegmatis*⁴⁶.

In addition, the effects of antibacterial and antifungal activities of essential oil extracted from *H. alpinum*, *H. barbatum*, *H. rumeliacum*, *H. maculatum*, *H. perforatum*, *H. hirsutum* have been examined by other researchers. The highest activity has been observed in the essential oil extracted from *H. barbatum*, and *H. alpinum* and *H. hirsutum* haven't showed any impact against *P. mirabilis*, *P. aeruginosa* and *C. Albicans*⁴⁷.

One of the most comprehensive researches on *Hypericum* has been carried out by Gibbons *et al.*⁴⁸. In their study, the impact of the chloroform and methanol extracts of 34 different species of *Hypericum* on *Staphylococcus* has been analysed. According to the results, it has been emphasized that *Hypericum* genus is of a great potential for anti-staphylococcal effect.

Table 1: Effects of different *Hypericum* species on Viruses

Hypericum Species	VIRUSES											References
	Cox (Coxsackie virus)	Measles Edmonston A	Poliomyelitis virus	Semliki forest virus	Vesicular stomatitis virus	HCMV	HIV1	HSV1	IAV	FIV	Hepati C	
H. caprifoliatum, H. polyanthemum										x		[37]
H. capitatum							x	x				[34]
H. connatum							x	x		x		[11, 33, 38]
H. hookerianum H. mysorens								x				[32]
H. japonicum									x			[51]
H. kotschy anum, H. neurocalycinum, H. salsugineum								x				[36]
H. perforatum						x	x	x	x		x	[9, 12, 23, 30, 35, 41, 46, 52-58]
H. revolutum	x	x	x	x	x			x				[59]
H. scabrum							x	x				[7, 25, 34, 40, 60, 85, 93-95]

Table 2: The effects of different *Hypericum* species on Bacteria

Hypericum Species	BACTERIA																														References		
	<i>Acinetobacter baumannii</i>	<i>Aeromonas hydrophila</i>	<i>Agrobacterium tumefaciens</i>	<i>Alcaligenes faecalis</i>	<i>Azotobacter chroococcum</i>	<i>Bacillus</i> sp.	<i>Bordetella bronchiseptica</i>	<i>Branhamella catarrhalis</i>	<i>Campylobacter</i> sp.	<i>Clostridium perfringens</i>	<i>Corynebacterium</i> sp.	<i>Escherichia coli</i>	<i>Enterococcus</i> sp.	<i>Erwinia carotovora</i>	<i>Enterobacter</i> sp.	<i>Helicobacter pylori</i>	<i>Klebsiella</i> sp.	<i>Listeria monocytogenes</i>	<i>Micrococcus</i> sp.	<i>Mycobacterium</i> sp.	<i>Nocardia</i> sp.	<i>Proteus vulgaris</i>	<i>Pseudomonas</i> sp.	<i>Sarcinulutea</i>	<i>Shigella</i> sp.	<i>Staphylococcus</i> sp.	<i>Serratia marcescens</i>	<i>Streptococcus</i> sp.	<i>Salmonella</i> sp.	<i>Vibriosp.</i>		<i>Yersinia enterocolitica</i>	
<i>H. acmosepalum</i>																										x							[47, 61]
<i>H. addingtonii</i>																										x							[47]
<i>H. aethiopicum</i>						x						x					x							x	x					x			[26, 46, 62-64]
<i>H. alpinum</i>			x			x						x							x			x	x	x		x				x			[46]
<i>H. amblycalyx</i>																x																	[4]
<i>H. androsaemum</i>												x	x							x				x			x			x		x	[43, 47, 66]
<i>H. arnoldianum</i>																										x							[47]
<i>H. ascyron</i>																										x							[67]
<i>H. barbatum</i>			x			x						x							x				x	x	x		x				x		[30, 46]
<i>H. beanii</i>																											x						[39, 47]
<i>H. bellum</i> ssp. <i>latisepalum</i>																										x							[47]
<i>H. calycinum</i>						x						x	x								x					x							[43, 68]
<i>H. canariense</i>						x	x					x					x		x	x				x		x							[8, 69]
<i>H. caprifoliatum</i>						x						x							x	x						x							[11, 69, 70]
<i>H. capitatum</i>						x		x		x		x														x							[34]
<i>H. carinatum</i>						x						x														x							[11, 70]

[9, 12, 23, 29, 30, 35,
41, 43, 46, 54, 55, 57,
66, 68, 79, 89]

The volatile compounds extracted from different species of *Hypericum* have been tested for their fungicidal activities on *Fusarium*^{49,50}, *C. albicans*, *Saccaromyces cerevisiae*^{11,26}, *Cladosporium cucumerinum*⁵¹, *Verticillium albo-atrum*, *Alternaria solani*, *Rhizoctonia solani*⁴⁹ by several authors.

In Table 3, antifungal activities of different *Hypericum* genus on 23 different fungi *Alternaria*, *Aspergillus* sp., *Candida* sp., *Cladosporium* sp., *Cryptococcus* sp., *Epidermophyton* sp., *Fusarium* sp., *Penicillium* sp., *Rhizoctonia solani*, *Rhodotorula rubra*, *Khuyveromyces fragilis*, *Microsporium* sp., *Ureaplasma urealyticum*, *Trichomonas vaginalis*, *Trichophyton*, *Gardnerella vaginalis*, *Oligella ureolytica*, *Neisseria gonorrhoeae*, *Saccharomyces* sp. and *Verticillium albo-atrum* have been analysed. Ten different *Hypericum* species have been studied in terms of their activity mechanism toward only one type of fungi. The most studied fungus has become *Candida* sp. and has followed by *Aspergillus* sp.

CONCLUSION

Hypericum species have become a popular area for many scientific researches for a long time. It has been estimated that it would be a significant material for both biological and chemical studies thanks to its availability and widespread distribution on earth for many years more. In this study, a large number of chemical studies on *Hypericum* genus in many different areas have been reviewed. It has been also expected that this attraction would remain as it is now for the researchers in the future; many new compounds, their properties and their effects would be discovered throughout the new studies, and it would be subject to different areas. Because, with their practical structure, plants will continue to be important resources in the future, for the development of new chemotherapeutic agents¹⁰⁴.

REFERENCES

- Crockett SL, Robson NKB. Taxonomy and Chemotaxonomy of the Genus *Hypericum*. Med Aromat Plant Sci Biotechnol 2011;5(1): 1–13.
- Saddiqe, Z, Naeem I, Maimoona A. A review of the antibacterial activity of *Hypericum perforatum* L. Journal of Ethnopharmacology 2010; 131(3): 511–521. <http://dx.doi.org/10.1016/j.jep.2010.07.034>
- Meral G, Karabay N. In vitro antibacterial activities of three *Hypericum* species from West Anatolia. Turkish Electronic Journal of Biotechnology Special issue 2002; 6–10.
- Davis PH. Flora of Turkey and the East Aegean Islands Vol:1 Edinburgh at the University Pres, 1965; 567.
- Robson NKB, Strid A. *Hypericum* L in: strid a, editor Mountain Flora of Greece, Cambridge University Pres, 1986; 594.
- Hobbs, C., St. John's Wort (*Hypericum perforatum* L.): a review. 1998: <http://www.christopherhobbs.com/website/library/articles/articlefiles/stjohnswort01.html>.
- Kizil G, Toker Z, Ozen HC, Aytekin C. The antimicrobial activity of essential oils of *Hypericum scabrum*, *Hypericum scabroides* and *Hypericum triquetrifolium*. Phytotherapy Research 2004; 18(4), 339–341. <http://dx.doi.org/10.1002/ptr.1460>
- Rabanal RM, Arias A, Prado B, Hernández-Pérez M, Sánchez-Mateo, CC. Antimicrobial studies on three species of *Hypericum* from the Canary Islands. J Ethnopharmacol 2002; 81(2), 287–92. [http://dx.doi.org/10.1016/S0378-8741\(02\)00083-1](http://dx.doi.org/10.1016/S0378-8741(02)00083-1)
- Milosevic T, Solujic S, Sukdolak S. In vitro study of ethanolic extract of *Hypericum perforatum* on growth and sporulation of some bacteria and fungi. Turk J Biol 2007; 31, 237–241.
- Dordević A, Zlatković B, Lazarević J, Palić R. A detailed chemical composition and antimicrobial activity of *Hypericum richeri* Vill subsp Grisebachii (Boiss) Nyman essential oil from Serbia. Journal of Medicinal Plants Research 2011; 5(23), 5486–5492.
- Dall'Agnol R, Ferraz A, Bernardi AP. Antimicrobial activity of some *Hypericum* species. Phytomedicine 2003; 10(6–7), 511–6. <http://dx.doi.org/10.1078/094471103322331476>
- Conforti F, Stati GA, Tundis R, Bianchi A, Agrimonti C, Sacchetti G, Andreotti E, Menichini F, Poli F. Comparative chemical composition and variability of biological activity of methanolic extracts from *Hypericum perforatum* L. Natural Product Research 2005; 19(3), 295–303. <http://dx.doi.org/10.1080/14786410410001715596>
- Bilia AR, Gallori S, Vincieri FF. St John's Wort and depression-efficiency, safety and tolerability. Life Sciences 2002; 70, 3077–3096. [http://dx.doi.org/10.1016/S0024-3205\(02\)01566-7](http://dx.doi.org/10.1016/S0024-3205(02)01566-7)
- Weiss RF. Herbal Medicine, England, Beaconsfield Puplichers, 1991; 295.
- Aramaki Y, Chiba K, Tada M. Spiro-lactones, hyperolactone AD from *Hypericum chinense*. Phytochemistry 1995; 38, 1419–1421. [http://dx.doi.org/10.1016/0031-9422\(94\)00862-N](http://dx.doi.org/10.1016/0031-9422(94)00862-N)
- Rocha L, Marston A, Potterat O, Kaplan MAC, Stoeckli EH, Hostettmann K. Antibacterial phloroglucinols and flavonoids from *Hypericum brasiliense*. Phytochemistry 1995; 40, 1447–1452. [http://dx.doi.org/10.1016/0031-9422\(95\)00507-4](http://dx.doi.org/10.1016/0031-9422(95)00507-4)
- Jayasuriya H, McChesney JD, Swanson SM, Pezzuto JM. Antimicrobial and cytotoxic activity of rottlerin-type compounds from *Hypericum drummondii*. J Nat Prod 1989; 52 (2), 325–331. <http://dx.doi.org/10.1021/np50062a018>
- Agostinis P, Vantieghem A, Merlevede W, De Witte PAM. Hypericin in cancer treatment: more light on the way. Int J Biochem Cell Biol 2002; 34 (3), 221–241. [http://dx.doi.org/10.1016/S1357-2725\(01\)00126-1](http://dx.doi.org/10.1016/S1357-2725(01)00126-1)
- Butterwerck V, Jürgenliemk G, Nahrstedt A, Winterhoff H. Flavonoids from *Hypericum perforatum* show antidepressant activity in the forced swimming test. Planta Medica 2000; 66, 3–6. <http://dx.doi.org/10.1055/s-2000-11119>
- Jacobson JM, Feinman L, Liebes L, Ostrow N, Koslowski V, Tobia A, Cabana BE, Lee D, Spritzler J, Prince AM. Pharmacokinetics, safety, and antiviral effects of hypericin, a derivative of St John's wort plant, in patients with chronic hepatitis C virus infection. Antimicrob Agents Chemother 2001; 45(2), 517–24. <http://dx.doi.org/10.1128/AAC.45.2.517-524.2001>
- Meruelo D, Lavie G, Lavie D. Therapeutic agents with dramatic antiretroviral activity and little toxicity at effective doses: aromatic polycyclic diones hypericin and pseudohypericin. Proc Natl Acad Sci 1988; 85, 5230–5234. <http://dx.doi.org/10.1073/pnas.85.14.5230>
- Cakir A, Mavi A, Yildirim A, Duru ME, Harmandar M, Kazaz C. Isolation and characterization of antioxidant phenolic compounds from the aerial parts of *Hypericum hyssopifolium* L. by activity-guided fractionation. J Ethnopharm 2003; 87(1), 73–83. [http://dx.doi.org/10.1016/S0378-8741\(03\)00112-0](http://dx.doi.org/10.1016/S0378-8741(03)00112-0)
- Ali M, Arfan M, Ahmad H, Zaman K, Khan F, Amarowicz R. Comparative antioxidant and antimicrobial activities of phenolic compounds extracted from five *Hypericum* species. Food Technol Biotechnol 2011; 49(2), 205–213.
- De'coster L, Stoeckli-Evan H, Msonthi JD, Hostettmann K. A new antifungal chromene and a related dichromene from *Hypericum revolutum*. Planta Med 1986; 55, 429. <http://dx.doi.org/10.1055/s-2007-969238>
- Unal EL, Mavi A, Kara AA, Cakir A, Şengül M, Yildirim A. Antimicrobial and antioxidant activities of some plants used as remedies in Turkish. Traditional Medicine 2008; 46(3), 207–224.
- Al-Bakri AG, Afifi FU. Evaluation of antimicrobial activity of selected plant extracts by rapid XTT colorimetry and bacterial enumeration. Journal of Microbiological Methods 2007; 68(1), 19–25. <http://dx.doi.org/10.1016/j.mimet.2006.05.013>
- Cakir A, Kordali S, Kilic H, Kaya E. Antifungal properties of essential oil and crude extracts of *Hypericum linarioides* Bosse.

- Biochemical Systematics and Ecology 2005; 33, 245–256. <http://dx.doi.org/10.1016/j.bse.2004.08.006>
28. Dulger B, Hacıoglu N. Antibacterial activity of three endemic *Hypericum* species used in folkloric medicine against methicilin-resistant *Staphylococcus aureus*. Biotechnology & Biotechnological Equipment Special Edition 2009; 23(2).
 29. Males Z, Brantner AH, Sović K, Pilepić KH, Plazibat M. Comparative phytochemical and antimicrobial investigations of *Hypericum perforatum* L subsp *perforatum* and *Hypericum perforatum* subsp *angustifolium* (DC) Gaudin. Acta Pharm 2006; 56, 359–367.
 30. Radulović N, Jovanović V, Stojanović G, Šmelcerović A, Spieller M, Asakawa Y. Screening of *in vitro* antimicrobial and antioxidant activity of nine *Hypericum* species from the Balkans. Food Chemistry 2007; 103, 15–21. <http://dx.doi.org/10.1016/j.foodchem.2006.05.062>
 31. Klemow KM, Bartlow A, Crawford J, Kocher N, Shah J, Ritsick M. Medical Attributes of St. John's Wort (*Hypericum perforatum*). in Herbal Medicine: Biomolecular and Clinical Aspects., I.F.F. Benzie and S. Wachtel-Galor, Editors. 2011; CRC Press. <http://dx.doi.org/10.1201/b10787-12>
 32. Vijayan P, Raghu C, Ashok G, Dhanaraj SA, Suresh B. Antiviral activity of medicinal plants of Nilgiris. Indian J Med Res 2004; 120, 24–29.
 33. Fritz D, Venturi C, Cargnin S, Schripsema J, Roehe P, Montanha J, Von Poser G. Herpes virus inhibitory substances from *Hypericum connatum* Lam, a plant used in southern Brazil to treat oral lesions. Journal of Ethnopharmacology 2007; 113, 517–520. <http://dx.doi.org/10.1016/j.jep.2007.07.013>
 34. Sokmen A, Jones BM, Erturk M. Antimicrobial activity of extracts from the cell cultures of some Turkish medicinal plants. Phytother Res 1999; 13, 355–357. [http://dx.doi.org/10.1002/\(SICI\)1099-1573\(199906\)13:4<355::AID-PTR454>3.3.CO;2-5](http://dx.doi.org/10.1002/(SICI)1099-1573(199906)13:4<355::AID-PTR454>3.3.CO;2-5)
 35. Pu X, Liang J, Wang X, Xu T, Hua L, Shang R, Liu Y, Xing Y. Anti-influenza virus effect of *Hypericum perforatum* L extract. Virologica Sinica 2009; 24(1), 19–27. <http://dx.doi.org/10.1007/s12250-009-2983-x>
 36. Duman R. Antiherpetic activity of some endemic *Hypericum* species in Turkey. African Journal of Biotechnology 2012; 11(5), 1240–1244.
 37. Schmitt AC, Ravazzolo APÜ, Von Poser GL. Investigation of some *Hypericum* species native to Southern of Brazil for antiviral activity. Journal of Ethnopharmacology 2001; 77(2-3), 239–245. [http://dx.doi.org/10.1016/S0378-8741\(01\)00314-2](http://dx.doi.org/10.1016/S0378-8741(01)00314-2)
 38. Westh, H., Zinn C.S., Rosdahl V.T. An international multicenter study of antimicrobial consumption and resistance in *Staphylococcus aureus* isolates from 15 hospitals in 14 countries. Microbial Drug Resistance 2004; 10: 169–176. <http://dx.doi.org/10.1089/1076629041310019>
 39. Ka Po Shiu W, Gibbons S. Anti-staphylococcal acylphloroglucinols from *Hypericum beanii*. Phytochemistry 2006; 67, 2568–2572. <http://dx.doi.org/10.1016/j.phytochem.2006.09.037>
 40. Pirbalouti AG, Malekpoor F, Hamed B. Ethnobotany and antimicrobial activity of medicinal plants of Bakhtiari Zagross mountains. Iran Journal of Medicinal Plants Research 2012; 6(5), 675–679.
 41. Avato P, Raffo F, Guglielmi G, Vitali C, Rosato A. Extracts from St John's Wort and their antimicrobial activity. Phytother Res 2004; 18, 230–232. <http://dx.doi.org/10.1002/ptr.1430>
 42. Kayser FH, Bienz K, Eckert J, Zinkernagel MR. Medical Microbiology-Georg Thieme Verlag 2004; 70469 Stuttgart, Germany.
 43. Mukherjee PK, Saritha GS, Suresh B. Antibacterial spectrum of *Hypericum hookerianum*. Fitoterapia 2001; 72(5), 558–560. [http://dx.doi.org/10.1016/S0367-326X\(00\)00331-2](http://dx.doi.org/10.1016/S0367-326X(00)00331-2)
 44. Mukherjee PK, Saritha GS, Suresh B. Antimicrobial potential of two different *Hypericum* species available in India. Phytotherapy Research 2002; 16(7), 692–695 <http://dx.doi.org/10.1002/ptr.1016>
 45. Sakar MK, Tamer AU. Antimicrobial activity of different extracts from *Hypericum* species. Fitoterapia 1990; 61, 464–466.
 46. Saroglou V, Marin PD, Rancic A, Veljic Skaltsa, MH. Composition and antimicrobial activity of the essential oil of six *Hypericum* species from Serbia. Biochemical Systematics and Ecology 2007; 35(3), 146–152. <http://dx.doi.org/10.1016/j.bse.2006.09.009>
 47. Gibbons S, Ohlendorf B, Johnsen I The genus *Hypericum* a valuable resource of anti-staphylococcal leads. Fitoterapia 2002; 73(4), 300–304.
 48. Cakir A, Kordali S, Zengin H, Izumi S, Hirata T. Composition and antifungal activity of essential oils isolated from *Hypericum hyssopifolium* and *Hypericum heterophyllum*. Flavour Fragr Jour 2004; 19, 62–68. <http://dx.doi.org/10.1002/ffj.1279>
 49. Fraternali D, Bertoli A, Giamperi L, Bucchini A, Ricci D, Menichini F, Trinciarelli E, Pistelli L. Antifungal evaluation of *Hypericum triquetrifolium* polar extracts against *Fusarium* spp. Natural Product Communications 2006; 1(12), 1117 – 1122.
 50. Rocha L, Marston A, Auxiliadora M, Kaplan C, Evans H S, Thull U, Testa B, Hostettmann K. An antifungal γ -pyrone and xanthenes with monoamine oxidase inhibitory activity from *Hypericum brasiliense*. Phytochemistry 1994; 36: 1381–1385. [http://dx.doi.org/10.1016/S0031-9422\(00\)89727-7](http://dx.doi.org/10.1016/S0031-9422(00)89727-7)
 51. Liu N, Hu XL, Meng YR, Zhu YT, Huang BS, Lin PZ. Effect of anti-influenza virus H3N2 of *Hypericum japonicum* *in vivo*. Zhong Yao Cai 2008; 31(7), 1022–1024.
 52. Yoon SJ, Kim JH, Kim TW, Choi UK. Screening of the antimicrobial activity against *Helicobacter pylori* from herb plants. Journal of the Korean Society for Applied Biological Chemistry 2005; 48(2), 161–165.
 53. Spritzler J, Prince AM (2001) Pharmacokinetics, safety, and antiviral effects of hypericin, a derivative of St John's Wort plant, in patients with chronic Hepatitis C virus infection. Antimicrob Agents Chemother 45, 517–524.
 54. Borchardt JR, Wyse DL, Sheaffer CC, Kauppi KL. Antimicrobial activity of native and naturalized plants of Minnesota and Wisconsin. Journal of Medicinal Plants Research 2008; 2(5), 098–110.
 55. Birt DF, Widrechner MP, Hammer KDP, Hillwig ML, Wei J, Kraus GA, Murphy PA, McCoy J, Wurtele ES, Neighbors JD, Wiemer DF, Maury WJ, Price JP. *Hypericum* in infection: Identification of anti-viral and anti-inflammatory constituents. Pharm Biol 2009; 47(8), 774–782.
 56. Mortensen T, Shen S, Shen F, Walsh M, Sims KR, Miller CD. Investigating the effectiveness of St John's Wort herb as an antimicrobial agent against. Mycobacteria Phytotherapy Research 2012; 26: 9, 1327–1333.
 57. Axarlis S, Mentis A, Demetrios C, Mitaku S, Skaltsounis AL, Marselos M, Malamas M. Antiviral *in vitro* activity of *Hypericum perforatum* L extract on the Human cytomegalovirus (HCMV). Phytotherapy Research 1998; 12(7), 507–511.
 58. Serkedjieva J, Manolova N, Zgórnjak-Nowosielska I, Zawilińska B, Grzybek J. Antiviral activity of the infusion (SHS-174) from flowers of *Sambucus nigra* L, aerial parts of *Hypericum perforatum* L, and roots of *Saponaria officinalis* L against Influenza and Herpes simplex viruses. Phytotherapy Research 1990; 4(3), 97–100.
 59. Cos P, Hermans N, Bruyne TD, Apers S, Sindambiwe JB, Berghe DV, Pieters L, Vlietinck AJ. Further evaluation of Rwandan medicinal plant extracts for their antimicrobial and antiviral activities. Journal of Ethnopharmacology 2002; 79(2), 155–163.
 60. Pirbalouti AG, Rahnama GH, Malekpoor F, Broujeni HR. Variation in antibacterial activity and phenolic content of *Hypericum scabrum* L populations. Journal of Medicinal Plants Research 2011; 5(17), 4119–4125.

61. Osman K, Evangelopoulos D, Basavannacharya C, Gupta A, McHugh TD. An antibacterial from *Hypericum acmosepalum* inhibits ATP-dependent MurE ligase from *Mycobacterium tuberculosis*. International Journal of Antimicrobial Agents 2012; 39, 124-129.
62. Van Vuuren S. Validating the antimicrobial activity of African traditional medicines. Planta Med 2010; 76.
63. Van Vuuren SF. The antimicrobial activity of medicinal plants to treat sexually transmitted infections (STI's). Planta Med 2009; 75.
64. Mthethwa NS. Antimicrobial activity testing of traditionally used plants for treating wounds and sores at Ongoye area KwaZulu-Natal, South Africa. University of Zululand Institutional Repository. Masters of science 2009; 200021480, 145.
65. Dulger B, Gonuz A. Antibacterial activity of the endemic *Hypericum kazdagensis*. Fitoterapia 2005; 76, 237–239.
66. Mazandarani M, Yassaghi S, Rezaei MB, Mansourian AR, Ghaemi, EO. Ethnobotany and Antibacterial Activities of Two Endemic Species of *Hypericum* in North-East of Iran. Asian Journal of Plant Sciences 2007; 6(2), 354-358.
67. Mu Q, Gibbons S, Stavri M, Smith E, Zhou F, Hu C. Antibacterial effects of *Hypericum ascyron* and *Hypericum japonicum* against multidrug-resistant *Staphylococcus aureus*. Pharmaceutical Biology 2006; 44(3), 157–159.
68. Maggi F, Cecchini C, Cresci A, Coman MM. Chemical composition and antimicrobial activity of the essential oils from several *Hypericum* taxa (Guttiferae) growing in central Italy (Appennino Umbro-Marchigiano). Chemistry & Biodiversity 2010; 7(2), 447–466.
69. Dall'Agnol R, Ferraz A, Bernardi AP, Albring D, Nör C, Schapoval EES, Von Poser GL. Bioassay-guided Isolation of antimicrobial benzopyrans and phloroglucinol derivatives from *Hypericum* species. Phytother Res 2005; 19, 291–293.
70. Fenner R, Sortino M, Rates SM, Dall'Agnol R, Ferraz A, Bernardi AP, Albring D, Nör C, Von Poser G, Schapova E, Zacchino, S. Antifungal activity of some Brazilian *Hypericum* species. Phytomedicine 2005; 12(3), 236-40.
71. Akbar S, Al-Yahya MA. Screening of Saudi plants for phytoconstituents, pharmacological and antimicrobial properties *Hypericum chrysostictum*. Australian Journal of Medical Herbalism 2011; 23:2, 1033-8330.
72. Fratianni F, Nazzaro F, Marandino A, Rosario Fusco M, Coppola R, De Feo V, De Martino L. Biochemical composition, antimicrobial activities, and anti-quorum-sensing activities of ethanol and ethyl acetate extracts from *Hypericum connatum* Lam (Guttiferae). Journal of Medicinal Food 2013; 16(5): 454-459. <http://dx.doi.org/10.1089/jmf.2012.0197>
73. Ladeira AM, Silva G, Raggi L, Young MC, Agripino DG. Chemical composition and antimicrobial activities of the essential oil of *Hypericum cordatum* (VellConc) N Robson (Hypericaceae). Journal of Essential Oil Research 2009; 21, 558-560.
74. Schwob I, Bessiere JM, Dherbomez M, Viano J. Composition and antimicrobial activity of the essential oil of *Hypericum coris*. Fitoterapia 2002; 73, 511–513.
75. Sassi A, Skhiri FH, Aouni M. Investigation of some medicinal plants from Tunisia for antimicrobial activities. Pharmaceutical Biology 2007; 45(5), 421–428.
76. Sarkisian A, Janssen M, Matta H, Henry GE, LaPlante KL, Rowley DC. Inhibition of bacterial growth and biofilm production by constituents from *Hypericum* spp. Phytotherapy Research, early view 2011.
77. Ghasemi Y, Khalaj A, Mohagheghzadeh A, Khosravi AR, Morowvat MH. Compositions and antimicrobial activities of the essential oils and extract of *Hypericum elongatum*. Journal of Applied Sciences 2007; 7, 2671-2675.
78. Dulger B, Hacıoglu, N, Dulger G. Antimicrobial activity of endemic *Hypericum havvae* from Turkey. Asian Journal of Chemistry 2008; 20(5), 3889-3892.
79. Pistelli L, Bertoli A, Zucconelli S, Morelli I, Panizzi, L, Menichini, F. Antimicrobial activity of crude extracts and pure compounds of *Hypericum hircinum*. Fitoterapia 2000; 71, 138-40.
80. Maggi F, Cecchini C, Cresci A, Coman MM, Tirillini B, Sagratini, G. Chemical composition and antimicrobial activity of *Hypericum hircinum* L. subsp. majus essential oil. Chemistry of Natural Compounds 2010; 46(1), 125-129.
81. Mehrabadi Yari L, Hosein M, Surmaghi S, Amin G, BadamiN, Emami M, Asgari T. Phytochemical and antimicrobial investigation of Taleghan plants species. Planta Med 2006; 72.
82. Toker Z, Kızıl G, Ozen HC, Kızıl M, Ertekin A.S. Compositions and antimicrobial activities of the essential oils of two *Hypericum* species from Turkey. Fitoterapia 2006; 77, 57–60.
83. Mukherjee PK, Saritha GS, Suresh B. Antibacterial spectrum of *Hypericum hookerianum*. Fitoterapia 2001; 72, 558-560.
84. JinBao Y, RongQing N, DongFeng W, CaiTang L, Jing L. Experimental study on resistance of *Hypericum japonicum* Thunb to *Helicobacter pylori*. China Journal of Traditional Chinese Medicine and Pharmacy 2009; 24(6), 821-823.
85. Baris D, Kızıl M, Aytekin C, Kızıl G, Yavuz M, Ceken B, Ertekin AS. In vitro antimicrobial and antioxidant activity of ethanol extract of three *Hypericum* and three *Achillea* species from Turkey. International Journal of Food Properties 2011; 14, 339-355.
86. Gudzić B, Djokovic D, Vajs V, PalićR, Stojanovic G. Composition and antimicrobial activity of the essential oil of *Hypericum maculatum*. Crantz Flavour and Fragrance Journal 2002; 17(5), 392–394.
87. Pulok K, Mukherjee GS, Saritha B. Antimicrobial potential of two different *Hypericum* species available in India. Phytotherapy Research 2002; 16(7), 692–695.
88. Nariman F, Eftekhari F, Habibi Z, Massarrat S, Malekzadeh R. Antibacterial activity of twenty Iranian plant extracts against clinical isolates. Iranian Journal of Basic Medical Sciences 2009; 12(2), 105-111.
89. Stamatis G, Kyriazopoulos P, Golegou S, Basayiannis A, Skaltsas S, Skaltsa H. In vitro anti-*Helicobacter pylori* activity of Greek herbal medicines. J Ethnopharmacol J Ethnopharmacol 2003; 88 (2-3), 175-179.
90. Herrera RM, Pérez M, Herrera MDA, López-García R, Rabanal RM, Arias A. Antimicrobial activity of extracts from plants endemic to the Canary Islands. Phytotherapy Research 1996; 10(4), 364–366.
91. Đorđević A, Lazarević J, Šmelcerović A, Stojanović G. The case of *Hypericum rochelii* Griseb & Schenk and *Hypericum umbellatum* A Kern essential oils: Chemical composition and antimicrobial activity. Journal of Pharmaceutical and Biomedical Analysis 2013; 77, 145–148.
92. Couladis M, Chinou IB, Tzakou O, Petrakis PV. Composition and antimicrobial activity of the essential oil of *Hypericum rumeliacum* subsp. *apollinis* (Boiss & Heldr). Phytotherapy Research 2003; 17, 152–154.
93. Dulger B, Gonuz A, Bilen S. Antimicrobial studies on three *Hypericum* Species from Turkey. South African Journal of Botany 2005; 1.
94. Sökmen A, Vardar-Ünlü G, Darıcı N, Şahin S. Antimicrobial activities of methanolic extracts of various plants growing in the Sivas district. Turkish Journal of Infection 2000; 14(2), 253-256.
95. Shafaghat A. Antioxidant, antimicrobial activities and fatty acid components of flower, leaf, stem and seed of *Hypericum scabrum*. Nat Prod Commun 2011; 6(11), 1739-42.
96. Ebrahimi A, Asgharian S, Habibian S. Antimicrobial activities of isolated endophytes from some Iranian native medicinal plants. Iranian Journal of Pharmaceutical Sciences 2010; 6(3), 217-222.

97. Pirbalouti AG, Jahanbazi P, Enteshari S, Malekpoor F, Hamed B. Antimicrobial activity of some Iranian medicinal plants. Arch Biol Sci 2010; 62(3), 633-642.
98. Akhbari M, Batooli H, Mozdianfard M. Comparative study of composition and biological activities of SDE prepared essential oils from flowers and fruits of two *Hypericum* species from central Iran. Natural Product Research 2012; 26(3), 193-202.
99. Ćorđević A, Šmelcerović A, Veličković D, Jovanović VS, Mitić V, Kostić D, Palić R. Antimicrobial and antioxidant activities of essential oil and crude extracts of *Hypericum tetrapterum* Fries (Hypericaceae). Journal of Medicinal Plants Research 2010; 4(14), 1441-1445.
100. Pistelli L, Bertoli A, Morelli I, Menichini F, Musmanno RA, Di Maggio T, Coratz G. Chemical and antibacterial evaluation of *Hypericum triquetrifolium* Turra. Phytotherapy Research 2005; 19, 787-791.
101. Rouis Z, Abid N, Koudja S, Yangui T, Elaissi A, Cioni PL, Flamini G, Aouni M. Evaluation of the cytotoxic effect and antibacterial, antifungal, and antiviral activities of *Hypericum triquetrifolium* Turra essential oils from Tunisia. BioMed Central Complementary and Alternative Medicine 2013; 13:24. <http://dx.doi.org/10.1186/1472-6882-13-24>
102. Nogueira T, Medeiros MA, Marcelo-Curto MJ, García-Pérez BE, Luna-Herrera J, Cêu Costa M. Profile of antimicrobial potential of fifteen *Hypericum* species from Portugal. Industrial Crops and Products 2013; 47, 126-131.
103. Akgöz Y, Toker Z. Antioxidant and Antimicrobial Effects of the *Hypericum retusum* Aucher plants extracts prepared in various solvents. Fresenius Environmental Bulletin 2013; 2A: 22, 493-1-499.
104. Akgöz Y. Biological and antioxidant activities of Silene Sp. Int. Res. J. Pharm. 2014; 5(11):810-813 <http://dx.doi.org/10.7897/2230-8407.0511166>

Cite this article as:

Yasemin Akgöz. The effects of *Hypericum* (Hypericaceae) species on microorganisms: A review. Int. Res. J. Pharm. 2015; 6(7):390-399 <http://dx.doi.org/10.7897/2230-8407.06781>

Source of support: Nil, Conflict of interest: None Declared

Disclaimer: IRJP is solely owned by Moksha Publishing House - A non-profit publishing house, dedicated to publish quality research, while every effort has been taken to verify the accuracy of the content published in our Journal. IRJP cannot accept any responsibility or liability for the site content and articles published. The views expressed in articles by our contributing authors are not necessarily those of IRJP editor or editorial board members.